

PATENT
[MOBT:217]
[38-21(15646)B US]

APPLICATION FOR UNITED STATES LETTERS PATENT

for

PREPARATION OF DEALLERGENIZED PROTEINS AND PERMUTEINS

by

Murtaza F. Alibhai

James D. Astwood

Charles F. McWherter and

Hugh A. Sampson

EXPRESS MAIL MAILING LABEL	
NUMBER	<u>E1 5212 77946 U 6</u>
DATE OF DEPOSIT	<u>1-5-01</u>
I hereby certify that this paper or fee is being deposited with the United States Postal Service "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to: Assistant Commissioner for Patents, Washington D.C. 20231.	
<u><i>Elizabeth Hays</i></u> Signature	

FIELD OF THE INVENTION

The invention relates generally to non-naturally occurring novel proteins which have insecticidal properties, and more specifically to the design, preparation, and use of proteins that have been deallergized while maintaining their insecticidal properties. Deallergized patatin proteins include variants that have had allergenic sequences modified, and permuteins that have had their amino acid sequences rearranged at one or more breakpoints.

BACKGROUND OF THE INVENTION

Insecticidal proteins

The use of natural products, including proteins, is a well known method of controlling many insect, fungal, viral, bacterial, and nematode pathogens. For example, endotoxins of *Bacillus thuringiensis* (*B.t.*) are used to control both lepidopteran and coleopteran insect pests. Genes producing these endotoxins have been introduced into and expressed by various plants, including cotton, tobacco, and tomato. There are, however, several economically important insect pests such as boll weevil (BWV), *Anthonomus grandis*, and corn rootworm (CRW), *Diabrotica* spp. that are not as susceptible to *B.t.* endotoxins as are insects such as lepidopterans. In addition, having other, different gene products for control of insects which are susceptible to *B.t.* endotoxins is important, if not vital, for resistance management.

It has been recently discovered that the major storage protein of potato tubers, patatins (Gaillard, T., *Biochem. J.* 121: 379-390, 1971; Racusen, D., *Can. J. Bot.*, 62: 1640-1644, 1984; Andrews, D.L., *et al.*, *Biochem. J.*, 252: 199-206, 1988), will control various insects, including western rootworm (WCRW, *Diabrotica virgifera*), southern corn rootworm (SCRW, *Diabrotica undecimpunctata*), and boll weevil (BWV, *Anthonomus grandis*) (U.S. Patent No. 5,743,477). Patatins are lethal to some larvae and will stunt the growth of survivors so that maturation is prevented or severely delayed, resulting in no reproduction. These proteins, have nonspecific lipid acyl hydrolase activity and studies have shown that the enzyme activity is essential for its insecticidal

activity (Strickland, J.A., *et al.*, *Plant Physiol.*, 109: 667-674, 1995; U.S. Patent No. 5,743,477). Patatins can be applied directly to the plants or introduced in other ways well known in the art, such as through the application of plant-colonizing microorganisms, which have been transformed to produce the enzymes, or by the plants themselves after similar transformation.

In potato, the patatins are found predominantly in tubers, but also at much lower levels in other plant organs (Hofgen, R. and Willmitzer, L., *Plant Science*, 66: 221-230, 1990). Genes that encode patatins have been previously isolated by Mignery, G.A., *et al.* (*Nucleic Acids Research*, 12: 7987-8000, 1984; Mignery, G.A., *et al.*, *Gene*, 62: 27-44, 1988; Stiekema, *et al.*, *Plant Mol. Biol.*, 11: 255-269, 1988) and others. Patatins are found in other plants, particularly solanaceous species (Ganal, *et al.*, *Mol. Gen. Genetics*, 225: 501-509, 1991; Vancanneyt, *et al.*, *Plant Cell*, 1: 533-540, 1989) and recently Zea mays (WO 96/37615). Rosahl, *et al.* (*EMBO J.*, 6: 1155-1159, 1987) transferred it to tobacco plants, and observed expression of patatin, demonstrating that the patatin genes can be heterologously expressed by plants.

Patatin is an attractive for use in planta as an insect control agent, but unfortunately a small segment of the population displays allergic reactions to patatin proteins, and in particular to potato patatin, as described below.

Food allergens

There are a variety of proteins that cause allergic reactions. Proteins that have been identified as causing an allergic reaction in hypersensitive patients occur in many plant and animal derived foods, pollens, fungal spores, insect venoms, insect feces, and animal dander and urine (King, H.C., *Ear Nose Throat J.*, 73(4): 237-241, 1994 ; Astwood, J.D., *et al.*, *Clin. Exp. Allergy*, 25: 66-72, 1995; Astwood, J.D. and Fuchs R.L., *Monographs in allergy Vol. 32: Highlights in food allergy*, pp. 105-120, 1996; Metcalfe, D.D., *et al.*, *Critical Reviews in Food Science and Nutrition*, 36S: 165-186, 1996). The offending proteins of many major sources of allergens have been characterized by clinical and molecular methods. The functions of allergenic proteins *in vivo* are diverse, ranging from enzymes to regulators of the cell cytoskeleton.

To understand the molecular basis of allergic disease, the important IgE binding epitopes of many allergen proteins have been mapped (Elsayed, S. and Apold, J., *Allergy* 38(7): 449-459, 1983; Elsayed, S. *et al.*, *Scand J. Clin. Lab. Invest. Suppl.* 204: 17-31 1991; Zhang, L., *et al.*, *Mol. Immunol.* 29(11): 1383-1389, 1992). The optimal peptide length for IgE binding has been reported to be between 8 and 12 amino acids. Conservation of epitope sequences is observed in homologous allergens of disparate species (Astwood, J.D., *et al.*, *Clin. Exp. Allergy*, 25: 66-72, 1995). Indeed, conservative substitutions introduced by site-directed mutagenesis reduce IgE binding of known epitopes when presented as peptides.

Food allergy occurs in 2-6 % of the population. Eight foods or food groups (milk, eggs, fish, crustacea, wheat, peanuts, soybeans, and tree nuts) account for 90% of allergies to foods. Nevertheless, over 160 different foods have been reported to cause adverse reactions, including potato (Hefle, S., *et al.*, *Crit. Rev. in Food Sci. Nutr.*, 36S: 69-90, 1996).

Mode of action of allergens

Regardless of the identity of the allergen, it is theorized that the underlying mechanism of allergen response is the same. Immediate hypersensitivity (or anaphylactic response) is a form of allergic reaction which develops very quickly, i.e., within seconds or minutes of exposure of the patient to the causative allergen, and is mediated by B lymphocyte IgE antibody production. Allergic patients exhibit elevated levels of IgE, mediating hypersensitivity by priming mast cells which are abundant in the skin, lymphoid organs, in the membranes of the eye, nose and mouth, and in the respiratory tree and intestines. The IgE in allergy-suffering patients becomes bound to the IgE receptors of mast cells. When this bound IgE is subsequently contacted by the appropriate allergen, the mast cell is caused to degranulate and release various substances such as histamine into the surrounding tissue (Church *et al.* In: Kay, A.B. ed., *Allergy and Allergic Diseases*, Oxford, Blackwell Science, pp. 149-197, 1997).

It is the release of these substances which is responsible for the clinical symptoms typical of immediate hypersensitivity, namely contraction of smooth muscle in the airways or in the intestine, the dilation of small blood vessels, and the increase in their

permeability to water and plasma proteins, the secretion of thick sticky mucus, and (in the skin) the stimulation of nerve endings that result in itching or pain. Immediate hypersensitivity is, at best, a nuisance to the sufferer; at worst it can present very serious problems and can in rare cases even result in death.

Allergic reactions to potato

Food allergy to potato is considered rare in the general population (Castells, M.C., *et al.*, *Allergy Clin. Immunol.*, 8: 1110-1114, 1986; Hannuksela, M., *et al.*, *Contact Dermatitis*, 3: 79-84, 1977; Golbert, T.M., *et al.*, *Journal of Allergy*, 44: 96-107, 1969). Approximately 200 individuals have participated in published clinical accounts of potato allergy (Hefle, S. *et al.*, *Critical Reviews in Food Science and Nutrition*, 36S: 69-90, 1996). A number of IgE binding proteins have been identified in potato tuber extracts (see Table 1), however the amino acid sequence and function of these proteins has not been determined (Wahl, R., *et al.*, *Intl. Arch. Allergy Appl. Immunol.*, 92: 168-174, 1990).

Table 1: Studies of potato tuber IgE-binding proteins (allergens)

Study	Protein Characteristics
(Castells, M.C. <i>et al.</i> <i>J. Allergy Clin. Immunol.</i> 78, 1110-1114, 1986)	Unknown 14 to 40 kDa
(Wahl, R. <i>et al.</i> <i>Intl. Arch. Allergy Appl. Immunol.</i> 92: 168-174, 1990)	Unknown 42/43 kDa
	Unknown 65 kDa
	Unknown 26 kDa
	Unknown 20 kDa
	Unknown 14 kDa
	Unknown < 14 kDa (~ 5 kDa)
(Ebner, C. <i>et al.</i> in: Wuthrich, B. & Ortolani, C. (eds.), <i>Highlights in food allergy. Monographs in Allergy, Volume 32</i> Basel, Karger, pp. 73-77, 1996)	Unknown 42/43 kDa
	Unknown 23 kDa
	Unknown ~ 16 kDa
	Unknown < 14 kDa (~ 5 kDa)

Improved safety from the use of hypoallergenic proteins

Patatin has been identified as an allergenic protein (Seppala, U. *et al.*, *J. Allergy Clin. Immunol.* 103:165-171, 1999). Accordingly, potato allergic subjects may not be

1 able to safely consume products containing unmodified patatin protein, such as crops to
2 which foliar applications of patatins have been applied, or crops which have been
3 engineered to express patatin. In addition, proliferation of food allergens in the food
4 supply is considered hazardous (Metcalf, D.D., *et al.*, *Critical Reviews and Food*
5 *Science and Nutrition*, 36S: 165-186, 1996). There are additional concerns regarding the
6 use of potentially allergenic food proteins where workers might be exposed to airborne
7 particulates, initiating a new allergic response (Moneret-Vautrin, D.A., *et al.*, *Rev. Med.*
8 *Interne.*, 17(7): 551-557, 1996).

9 Permuteins

10 Novel proteins generated by the method of sequence transposition resembles that
11 of naturally occurring pairs of proteins that are related by linear reorganization of their
12 amino acid sequences (Cunningham, *et al.* *Proc. Natl. Sci., U.S.A.*, 76: 3218-3222, 1979;
13 Teather, *et al.*, *J. Bacteriol.*, 172: 3837-3841, 1990; Schimming, *et al.*, *Eur. J. Biochem.*,
14 204: 13-19, 1992; Yamiuchi, *et al.*, *FEBS Lett.*, 260: 127-130, 1991; MacGregor, *et al.*,
15 *FEBS Lett.*, 378: 263-266, 1996). The first *in vitro* application of sequence
16 rearrangement to proteins was described by Goldenberg and Creighton (Goldenberg and
17 Creighton, *J. Mol. Biol.*, 165: 407-413, 1983). A new N-terminus is selected at an
18 internal site (breakpoint) of the original sequence, the new sequence having the same
19 order of amino acids as the original from the breakpoint until it reaches an amino acid
20 that is at or near the original C-terminus. At this point the new sequence is joined, either
21 directly or through an additional portion or sequence (linker), to an amino acid that is at
22 or near the original N-terminus, and the new sequence continues with the same sequence
23 as the original until it reaches a point that is at or near the amino acid that was N-terminal
24 to the breakpoint site of the original sequence, this residue forming the new C-terminus
25 of the chain. This approach has been applied to proteins which range in size from 58 to
26 462 amino acids and represent a broad range of structural classes (Goldenberg and
27 Creighton, *J. Mol. Biol.*, 165: 407-413, 1983; Li and Coffino, *Mol. Cell. Biol.*, 13: 2377-
28 2383, 1993; Zhang, *et al.*, *Nature Struct. Biol.*, 1: 434-438, 1995; Buchwalder, *et al.*,
29 *Biochemistry*, 31: 1621-1630, 1994; Protasova, *et al.*, *Prot. Eng.*, 7: 1373-1377, 1995;
30 Mullins, *et al.*, *J. Am. Chem. Soc.*, 116: 5529-5533, 1994; Garrett, *et al.*, *Protein Science*,

5: 204-211, 1996; Hahn, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 91: 10417-10421, 1994;
Yang and Schachman, *Proc. Natl. Acad. Sci. U.S.A.*, 90: 11980-11984, 1993; Luger, *et al.*, *Science*, 243: 206-210, 1989; Luger, *et al.*, *Prot. Eng.*, 3: 249-258, 1990; Lin, *et al.*, *Protein Science*, 4: 159-166, 1995; Vignais, *et al.*, *Protein Science*, 4: 994-1000, 1995; Ritco-Vonsovici, *et al.*, *Biochemistry*, 34: 16543-16551, 1995; Horlick, *et al.*, *Protein Eng.*, 5: 427-431, 1992; Kreitman, *et al.*, *Cytokine*, 7: 311-318, 1995; Viguera, *et al.*, *Mol. Biol.*, 247: 670-681, 1995; Koebnik and Kramer, *J. Mol. Biol.*, 250: 617-626, 1995; Kreitman, *et al.*, *Proc. Natl. Acad. Sci.*, 91: 6889-6893, 1994).

There exists a need for the development of plant expressible insecticidal proteins which possess minimal or no allergenic properties.

SUMMARY OF THE INVENTION

Novel protein sequences, and nucleic acid sequences encoding them are disclosed. The proteins maintain desirable enzymatic and insecticidal properties while displaying reduced or eliminated allergenicity.

Allergenic epitopes are identified by scanning overlapping peptide sequences with an immunoreactivity assay. Alanine scanning and 'rational substitution' is performed on identified peptide sequences to determine specific amino acids which contribute to antibody binding, and presumably, to the allergenic properties of the whole protein. Individual mutations are introduced into the whole protein sequence by methods such as site directed mutagenesis of the encoding nucleic acid sequence to delete or modify the allergenic sequences.

Glycosylation target residues are identified within amino acid sequences of proteins which have demonstrated allergy eliciting properties. Glycosylation target amino acid residues are rationally substituted with other amino acid residues to eliminate glycosylation and to provide a variant deglycosylated protein. The variant protein may then exhibit reduced allergen eliciting properties and may also exhibit reduced binding to IgE within serum of patients observed to be allergic to said glycosylated protein.

Permutins of the deallergized protein sequences can be constructed to further reduce or eliminate allergic reactions. The encoding nucleic acid sequence is modified to

1 produce a non-naturally occurring protein having a linear amino acid sequence different
2 from the naturally occurring protein sequence, while maintaining enzymatic and
3 insecticidal properties. The permutein is preferably produced in plant cells, and more
4 preferably produced at a concentration which is toxic to insects ingesting the plant cells.

5 Methods for reducing, eliminating, or decreasing allergen eliciting properties of a
6 protein are specifically contemplated herein. Such methods comprise steps including
7 identifying one or more patients exhibiting an allergic sensitivity to an allergen eliciting
8 protein and obtaining a sample of serum from the patient; exposing the patient serum to a
9 first set of synthetic overlapping peptides which represent the allergen eliciting protein in
10 order to identify such peptides which exhibit epitopes which bind to IgE present within
11 the allergic patients' serum and wherein the IgE present in the serum has a specific
12 affinity for the said allergen eliciting protein; producing a second set of peptides which
13 are variant peptides based on the first set of peptides which were identified to bind
14 specifically to IgE present in patient serum, wherein the second set variant peptides
15 exhibit alanine scanning or rational scanning amino acid substitutions which exhibit
16 reduced, decreased, or eliminated IgE binding when compared to the first set non-variant
17 peptides, and wherein such substitutions which reduce, eliminate or decrease IgE binding
18 are identified as result effective substitutions; and modifying the amino acid sequence of
19 the allergen eliciting protein to contain one or more of said result effective substitutions,
20 wherein the modified protein is a variant of the allergen eliciting protein which lacks
21 allergen eliciting protein or exhibits reduced allergen eliciting properties, and wherein the
22 variant of the allergen eliciting protein comprising one or more result effective
23 substitutions exhibits reduced, decreased, or totally eliminated binding of IgE present
24 within said patients' serum.

25 The novel proteins can be used in controlling insects, as nutritional supplements,
26 in immunotherapy protocols, and in other potential applications. Transgenic plant cells
27 and plants containing the encoding nucleic acid sequence can be particularly beneficial in
28 the control of insects, and as a nutritional/immunotherapy material.

DESCRIPTION OF THE FIGURES

The following figures form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention can be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

Figure 1 illustrates the alignment of potato patatin PatA (acyl lipid hydrolase) with patatin (acyl lipid hydrolase) homologs and related amino acid sequences, the homologs and related sequences being from both dicot and monocot plant species.

Figure 2 illustrates IgE binding to overlapping peptide sequences.

Figure 3 illustrates construction of nucleic acid sequences encoding patatin permutein proteins, and in this figure for illustrative purposes a breakpoint at position 247 is shown.

DESCRIPTION OF THE SEQUENCE LISTINGS

The following description of the sequence listing forms part of the present specification and is included to further demonstrate certain aspects of the present invention. The invention can be better understood by reference to one or more of these sequences in combination with the detailed description of specific embodiments presented herein.

SEQ ID NO:1	DNA sequence encoding a patatin (acyl lipid hydrolase) protein
SEQ ID NO:2	potato patatin protein sequence
SEQ ID NO:3	thermal amplification primer
SEQ ID NO:4	thermal amplification primer
SEQ ID NO:5	thermal amplification product
SEQ ID NO:6	Pre-cleavage patatin protein produced in <i>Pichia pastoris</i>
SEQ ID NO:7	Post-cleavage patatin protein produced in <i>Pichia pastoris</i>
SEQ ID NO:8	Y106F mutagenic primer
SEQ ID NO:9	Y129F mutagenic primer
SEQ ID NO:10	Y185F mutagenic primer

SEQ ID NO:11	Y193F mutagenic primer
SEQ ID NO:12	Y185F and Y193F mutagenic primer
SEQ ID NO:13	Y270F mutagenic primer
SEQ ID NO:14	Y316F mutagenic primer
SEQ ID NO:15	Y362F mutagenic primer
SEQ ID NO:16-104	Peptide scan sequences of a patatin protein
SEQ ID NO:105-241	Alanine and rational scan sequences of selected patatin peptides
SEQ ID NO:242	thermal amplification primer 27
SEQ ID NO:243	thermal amplification primer 48
SEQ ID NO:244	thermal amplification primer 47
SEQ ID NO:245	thermal amplification primer 36
SEQ ID NO:246	pMON37402 sequence encoding permutein protein
SEQ ID NO:247	Permutein protein encoded from pMON37402 sequence
SEQ ID NO:248	thermal amplification primer 58
SEQ ID NO:249	thermal amplification primer 59
SEQ ID NO:250	pMON37405 sequence encoding permutein protein
SEQ ID NO:251	Permutein protein encoded by pMON37405 sequence
SEQ ID NO:252	thermal amplification primer 60
SEQ ID NO:253	thermal amplification primer 61
SEQ ID NO:254	pMON37406 sequence encoding permutein protein
SEQ ID NO:255	Permutein protein encoded by pMON37406 sequence
SEQ ID NO:256	thermal amplification primer 62
SEQ ID NO:257	thermal amplification primer 63
SEQ ID NO:258	pMON37407 sequence encoding permutein protein
SEQ ID NO:259	Permutein protein encoded by pMON37407 sequence
SEQ ID NO:260	thermal amplification primer 60
SEQ ID NO:261	thermal amplification primer 65
SEQ ID NO:262	pMON37408 sequence encoding permutein protein
SEQ ID NO:263	Permutein protein encoded by pMON37408 sequence
SEQ ID NO:264	pMON40701 sequence encoding permutein protein

SEQ ID NO:265	Permutin protein encoded by pMON40701 sequence
SEQ ID NO:266	thermal amplification primer Syn1
SEQ ID NO:267	thermal amplification primer Syn2
SEQ ID NO:268	thermal amplification primer Syn3
SEQ ID NO:269	thermal amplification primer Syn4
SEQ ID NO:270	pMON40703 sequence encoding permutin protein
SEQ ID NO:271	Permutin protein encoded by pMON40703 sequence
SEQ ID NO:272	thermal amplification primer Syn10
SEQ ID NO:273	thermal amplification primer Syn11
SEQ ID NO:274	pMON40705 sequence encoding permutin protein
SEQ ID NO:275	Permutin protein encoded by pMON40705 sequence
SEQ ID NO:276-277	Permutin linker sequences
SEQ ID NO:278	Patatin isozyme PatA+ (including signal peptide)
SEQ ID NO:279	Patatin isozyme PatB+ (including signal peptide)
SEQ ID NO:280	Patatin isozyme PatFm (mature protein lacking signal peptide)
SEQ ID NO:281	Patatin isozyme PatIm (mature protein lacking signal peptide)
SEQ ID NO:282	Patatin isozyme PatL+ (including signal peptide)
SEQ ID NO:283	Rational substitution peptide
SEQ ID NO:284	Corn homolog peptide
SEQ ID NO:285	patatin homolog Pat17 DNA coding sequence and amino acid translation
SEQ ID NO:286	patatin homolog Pat17 amino acid sequence
SEQ ID NO:287	dicot patatin homolog amino acid sequence pentin1_phb
SEQ ID NO:288	dicot patatin homolog amino acid sequence 5c9_phb
SEQ ID NO:289	maize patatin homolog amino acid sequence corn1_pep
SEQ ID NO:290	maize patatin homolog amino acid sequence corn2_pep
SEQ ID NO:291	maize patatin homolog amino acid sequence corn3_pep
SEQ ID NO:292	maize patatin homolog amino acid sequence corn4_pep
SEQ ID NO:293	maize patatin homolog amino acid sequence corn5_pep

DEFINITIONS

The following definitions are provided in order to aid those skilled in the art in understanding the detailed description of the present invention. Some words and phrases may also be defined in other sections of the specification. No limitation should be placed on the definitions presented for the terms below, where other meanings are evidenced elsewhere in the specification in addition to those specified below.

“Allergen” refers to a biological or chemical substance that induces an allergic reaction or response. An allergic response can be an immunoglobulin E-mediated response.

Amino acid codes: A (Ala) = alanine; C (Cys) = cysteine; D (Asp) = aspartic acid; E (Glu) = glutamic acid; F (Phe) = phenylalanine; G (Gly) = glycine; H (His) = histidine; I (Ile) = isoleucine; K (Lys) = lysine; L (Leu) = leucine; M (Met) = methionine; N (Asn) = asparagine; P (Pro) = proline; Q (Gln) = glutamine; R (Arg) = arginine; S (Ser) = serine; T (Thr) = threonine; V (Val) = valine; W (Trp) = tryptophan; Y (Tyr) = tyrosine.

“Amplification” refers to increasing the number of copies of a desired molecule.

“Coding sequence”, “open reading frame”, and “structural sequence” refer to the region of continuous sequential nucleic acid base pair triplets encoding a protein, polypeptide, or peptide sequence.

“Codon” refers to a sequence of three nucleotides that specify a particular amino acid.

“Complementarity” refers to the specific binding of adenine to thymine (or uracil in RNA) and cytosine to guanine on opposite strands of DNA or RNA.

“Deallergenize” (render hypoallergenic) refers to the method of engineering or modifying a protein or the encoding DNA such that the protein has a reduced or eliminated ability to induce an allergic response with respect to the ability of the unmodified protein. A deallergized protein can be referred to as being hypoallergenic. The degree of deallergenization of a protein can be measured *in vitro* by the reduced binding of IgE antibodies.

“DNA segment heterologous to the promoter region” means that the coding DNA segment does not exist in nature in the same gene with the promoter to which it is now attached.

1 “DNA segment” refers to a DNA molecule that has been isolated free of total
2 genomic DNA of a particular species.

3 “Electroporation” refers to a method of introducing foreign DNA into cells that
4 uses a brief, high voltage DC (direct current) charge to permeabilize the host cells,
5 causing them to take up extra-chromosomal DNA.

6 “Encoding DNA” refers to chromosomal DNA, plasmid DNA, cDNA, or
7 synthetic DNA which encodes any of the enzymes discussed herein.

8 “Endogenous” refers to materials originating from within an organism or cell.

9 “Endonuclease” refers to an enzyme that hydrolyzes double stranded DNA at
10 internal locations.

11 “Epitope” refers to a region on an allergen that interacts with the cells of the
12 immune system. Epitopes are often further defined by the type of antibody or cell with
13 which they interact, e.g. if the region reacts with B-cells or antibodies (IgE), it is called a
14 B-cell epitope.

15 “Exogenous” refers to materials originating from outside of an organism or cell.
16 This typically applies to nucleic acid molecules used in producing transformed or
17 transgenic host cells and plants.

18 “Expressibly coupled” and “expressibly linked” refer to a promoter or promoter
19 region and a coding or structural sequence in such an orientation and distance that
20 transcription of the coding or structural sequence can be directed by the promoter or
21 promoter region.

22 “Expression” refers to the transcription of a gene to produce the corresponding
23 mRNA and translation of this mRNA to produce the corresponding gene product, i.e., a
24 peptide, polypeptide, or protein.

25 “Heterologous DNA” refers to DNA from a source different than that of the
26 recipient cell.

27 “Homologous DNA” refers to DNA from the same source as that of the recipient
28 cell.

29 “Identity” refers to the degree of similarity between two nucleic acid or protein
30 sequences. An alignment of the two sequences is performed by a suitable computer
31 program. A widely used and accepted computer program for performing sequence

alignments is CLUSTALW v1.6 (Thompson, et al. *Nucl. Acids Res.*, 22: 4673-4680, 1994). The number of matching bases or amino acids is divided by the total number of bases or amino acids, and multiplied by 100 to obtain a percent identity. For example, if two 580 base pair sequences had 145 matched bases, they would be 25 percent identical. If the two compared sequences are of different lengths, the number of matches is divided by the shorter of the two lengths. For example, if there were 100 matched amino acids between 200 and a 400 amino acid proteins, they are 50 percent identical with respect to the shorter sequence. If the shorter sequence is less than 150 bases or 50 amino acids in length, the number of matches are divided by 150 (for nucleic acid bases) or 50 (for amino acids), and multiplied by 100 to obtain a percent identity.

"IgE" (Immunoglobulin E) refers to a specific class of immunoglobulin secreted by B cells. IgE binds to specific receptors on Mast cells. Interaction of an allergen with mast cell-bound IgE may trigger allergic symptoms.

"Immunotherapy" refers to any type of treatment that targets the immune system. Allergy immunotherapy is a treatment in which a progressively increasing dose of an allergen is given in order to induce an immune response characterized by tolerance to the antigen/allergen, also known as desensitization.

"*In vitro*" refers to "in the laboratory" and/or "outside of a living organism".

"*In vivo*" refers to "in a living organism".

"Insecticidal polypeptide" refers to a polypeptide having insecticidal properties that adversely affects the growth and development of insect pests.

"Monocot" refers to plants having a single cotyledon (the first leaf of the embryo of seed plants); examples include cereals such as maize, rice, wheat, oats, and barley.

"Multiple cloning site" refers to an artificially constructed collection of restriction enzyme sites in a vector that facilitates insertion of foreign DNA into the vector.

"Mutation" refers to any change or alteration in a nucleic acid sequence. Several types exist, including point, frame shift, splicing, and insertion/deletions.

"Native" refers to "naturally occurring in the same organism". For example, a native promoter is the promoter naturally found operatively linked to a given coding sequence in an organism. A native protein is one naturally found in nature and untouched or not otherwise manipulated by the hand of man.

1 “Nucleic acid segment” is a nucleic acid molecule that has been isolated free of
2 total genomic DNA of a particular species, or that has been synthesized. Included with
3 the term “nucleic acid segment” are DNA segments, recombinant vectors, plasmids,
4 cosmids, phagemids, phage, viruses, etcetera.

5 “Nucleic acid” refers to deoxyribonucleic acid (DNA) and ribonucleic acid
6 (RNA).

7 Nucleic acid codes: A = adenosine; C = cytosine; G = guanosine; T = thymidine;
8 N = equimolar A, C, G, and T; I = deoxyinosine; K = equimolar G and T; R = equimolar
9 A and G; S = equimolar C and G; W = equimolar A and T; Y = equimolar C and T.

10 “Open reading frame (ORF)” refers to a region of DNA or RNA encoding a
11 peptide, polypeptide, or protein or capable of being translated to protein, or a region of
12 DNA capable of being transcribed into an RNA product.

13 “Plasmid” refers to a circular, extrachromosomal, self-replicating piece of DNA.

14 “Point mutation” refers to an alteration of a single nucleotide in a nucleic acid
15 sequence.

16 “Polymerase chain reaction (PCR)” refers to an enzymatic technique to create
17 multiple copies of one sequence of nucleic acid. Copies of DNA sequence are prepared
18 by shuttling a DNA polymerase between two oligonucleotides. The basis of this
19 amplification method is multiple cycles of temperature changes to denature, then re-
20 anneal primers, followed by extension to synthesize new DNA strands in the region
21 located between the flanking primers. Also known as thermal amplification.

22 “Probe” refers to a polynucleotide sequence which is complementary to a target
23 polynucleotide sequence in the analyte. An antibody can also be used as a probe to detect
24 the presence of an antigen. In that sense, the antigen binding domain of the antibody has
25 some detectable affinity for the antigen and binds thereto. The binding of the antibody to
26 the antigen can be measured by means known in the art, such as by chemiluminescence,
27 phosphorescence, fluorescence, colorimetric chemical deposition at the site of binding, or
28 otherwise.

29 “Promoter” or “promoter region” refers to a DNA sequence, usually found
30 upstream (5') to a coding sequence, that controls expression of the coding sequence by
31 controlling production of messenger RNA (mRNA) by providing the recognition site for

1 RNA polymerase and/or other factors necessary for start of transcription at the correct
2 site. As contemplated herein, a promoter or promoter region includes variations of
3 promoters derived by means of ligation to various regulatory sequences, random or
4 controlled mutagenesis, and addition or duplication of enhancer sequences. The
5 promoter region disclosed herein, and biologically functional equivalents thereof, are
6 responsible for driving the transcription of coding sequences under their control when
7 introduced into a host as part of a suitable recombinant vector, as demonstrated by its
8 ability to produce mRNA.

9 “Recombinant DNA construct” or “recombinant vector” refers to any agent such
10 as a plasmid, cosmid, virus, autonomously replicating sequence, phage, or linear or
11 circular single-stranded or double-stranded DNA or RNA nucleotide sequence, derived
12 from any source, capable of genomic integration or autonomous replication, comprising a
13 DNA molecule in which one or more DNA sequences have been linked in a functionally
14 operative manner. Such recombinant DNA constructs or vectors are capable of
15 introducing a 5' regulatory sequence or promoter region and a DNA sequence for a
16 selected gene product into a cell in such a manner that the DNA sequence is transcribed
17 into a functional mRNA which is translated and therefore expressed. Recombinant DNA
18 constructs or recombinant vectors can be constructed to be capable of expressing
19 antisense RNAs, in order to inhibit translation of a specific RNA of interest.

20 “Recombinant proteins”, also referred to as “heterologous proteins”, are proteins
21 which are normally not produced by the host cell.

22 “Regeneration” refers to the process of growing a plant from a plant cell (e.g.,
23 plant protoplast or explant).

24 “Regeneration” refers to the process of growing a plant from a plant cell (e.g.,
25 plant protoplast or explant).

26 “Regulatory sequence” refers to a nucleotide sequence located upstream (5'),
27 within, and/or downstream (3') to a DNA sequence encoding a selected gene product
28 whose transcription and expression is controlled by the regulatory sequence in
29 conjunction with the protein synthetic apparatus of the cell.

1 “Restriction enzyme” refers to an enzyme that recognizes a specific palindromic
2 sequence of nucleotides in double stranded DNA and cleaves both strands; also called a
3 restriction endonuclease. Cleavage typically occurs within the restriction site.

4 “Result-effective substitution” (RES) refers to an amino acid substitution within
5 an IgE-binding region (epitope) of a target protein which reduces or eliminates the IgE
6 binding by that epitope. Examples herein are directed to patatin protein and homologues,
7 however, as will be readily recognized by those skilled in the art, the method is more
8 broadly applicable to proteins other than patatins, and in particular is applicable to any
9 protein exhibiting allergen eliciting properties.

10 “Selectable marker” refers to a nucleic acid sequence whose expression confers a
11 phenotype facilitating identification of cells containing the nucleic acid sequence.
12 Selectable markers include those which confer resistance to toxic chemicals (e.g.
13 ampicillin resistance, kanamycin resistance), complement a nutritional deficiency (e.g.
14 uracil, histidine, leucine), or impart a visually distinguishing characteristic (e.g. color
15 changes or fluorescence).

16 “Transcription” refers to the process of producing an RNA copy from a DNA
17 template.

18 “Transformation” refers to a process of introducing an exogenous nucleic acid
19 sequence (e.g., a vector, recombinant nucleic acid molecule) into a cell or protoplast in
20 which that exogenous nucleic acid is incorporated into a chromosome or is capable of
21 autonomous replication.

22 “Transformed cell” is a cell whose DNA has been altered by the introduction of
23 an exogenous nucleic acid molecule into that cell.

24 “Transgenic cell” refers to any cell derived from or regenerated from a
25 transformed cell or derived from a transgenic cell. Exemplary transgenic cells include
26 plant calli derived from a transformed plant cell and particular cells such as leaf, root,
27 stem, e.g., somatic cells, or reproductive (germ) cells obtained from a transgenic plant.

28 “Transgenic plant” refers to a plant or progeny thereof derived from a transformed
29 plant cell or protoplast, wherein the plant DNA contains an introduced exogenous nucleic
30 acid sequence not originally present in a native, non-transgenic plant of the same species.

1 Alternatively, the plant DNA can contain the introduced nucleic acid sequence in a higher
2 copy number than in the native, non-transgenic plant of the same species.

3 "Translation" refers to the production of protein from messenger RNA.

4 "Vector" refers to a plasmid, cosmid, bacteriophage, or virus that carries foreign
5 DNA into a host organism.

6 "Western blot" refers to protein or proteins that have been separated by
7 electrophoresis, transferred and immobilized onto a solid support, then probed with an
8 antibody.

9 DETAILED DESCRIPTION OF THE INVENTION

10 Design of deallergenized patatin proteins

11 Deallergenizing a protein can be accomplished by the identification of allergenic
12 sites, followed by modification of the sites to reduce or eliminate the binding of
13 antibodies to the sites. The IgE-binding regions of patatin were previously unreported.
14 Mapping of the IgE epitopes was accomplished by synthesizing 10-mer peptides based on
15 the patatin 17 protein sequence (SEQ ID NO: 2) which overlap by six amino acids. As
16 potato proteins are denatured upon cooking potato products, it is expected that the 10-mer
17 peptides sufficiently mimic the unfolded full length protein for antibody binding
18 purposes. Peptides were identified based upon their ability to bind to IgE antibodies.
19 Individual amino acids within the identified peptides were changed to reduce or eliminate
20 binding to IgE present in sera from potato sensitive patients. These changes are termed
21 result-effective amino acid substitutions (RES). The RES can be subsequently introduced
22 into the full length protein by site directed mutagenesis of the encoding nucleic acid
23 sequence or other means known in the art. Similar strategies have been employed
24 elsewhere to determine the dominant IgE epitopes in a major peanut allergen (Stanley,
25 J.S., *et al.*, *Arch. Biochem. Biophys.*, 342(2): 244-253, 1997).

26 Certain amino acid residues important for allergenicity of patatin are identified.
27 Some of the designed patatin peptides wherein single amino acid residues were replaced
28 with alanine or phenylalanine, showed significantly reduced or no binding to sera from
29 potato sensitive patients.

1 A "deallergenized patatin" refers to a patatin protein differing in at least one of
2 the amino acid residues as defined by the result effective substitutions resulting in the
3 patatin protein having reduced reactivity towards sera from potato sensitive patients. The
4 deallergenized patatin preferably maintains insecticidal properties, and preferably
5 maintains its characteristic enzymatic profile.

6 Summary of method to deallergenize a patatin protein

- 7 ▪ Mapping of IgE epitopes by immunoassay of synthetic overlapping peptides
8 using sera from potato sensitive patients;
- 9 ▪ Identification of result-effective substitutions by alanine scanning and/or
10 rational scanning;
- 11 ▪ Modification of the amino acid sequence of patatin by site-directed
12 mutagenesis of the encoding nucleic acid sequence;
- 13 ▪ Evaluation of enzyme activity (esterase) and/or insecticidal activity of the
14 modified protein(s); and
- 15 ▪ Evaluation of the new protein(s) for allergenicity by IgE immunoassay.

16
17 Nucleic acid sequences encoding patatin have been cloned by several
18 investigators (e.g. Mignery, *et al.*, *Nucleic Acids Research*, 12: 7987-8000, 1984;
19 Mignery, *et al.*, *Gene*, 62: 27-44, 1988; WO 94/21805; Canadian Patent Application No.
20 2090552). These nucleic acid sequences can then be manipulated using site directed
21 mutagenesis to encode a hypoallergenic patatin. These nucleic acid sequences can than
22 be used to transform bacterial, yeast or plant cells, resulting in the production of
23 hypoallergenic patatin protein.

24 Deallergenized patatin proteins

25 For simplicity, individual amino acids are referred to by their single letter codes.
26 Correlation between the single letter codes, three letter codes, and full amino acid names
27 is presented in the definitions section above.

28 One embodiment of the invention is an isolated deallergenized patatin protein.
29 The protein is modified relative to the wild-type protein sequence such that they exhibit

1 reduced binding to anti-patatin antibodies such as those obtained from humans or animals
2 allergic to potatoes. The reduced binding is measured relative to the binding of the
3 unmodified patatin protein to the anti-patatin antibodies.

4 The deallergenized patatin protein can comprise SEQ ID NO:2 modified in one or
5 more of the following regions, or SEQ ID NO:7 modified in one or more of the following
6 regions. The single or multiple amino acid modifications reduce the binding of the
7 modified protein relative to the binding of the corresponding unmodified protein. The
8 regions for modification include amino acid positions 104-113 of SEQ ID NO:2 (85-94
9 of SEQ ID NO:7), 128-137 of SEQ ID NO:2 (109-118 of SEQ ID NO:7), 184-197 of
10 SEQ ID NO:2 (165-178 of SEQ ID NO:7), 264-277 of SEQ ID NO:2 (245-258 of SEQ
11 ID NO:7), 316-325 of SEQ ID NO:2 (297-306 of SEQ ID NO:7), and 360-377 of SEQ ID
12 NO:2 (341-358 of SEQ ID NO:7). The possible amino acid modifications include
13 replacing an amino acid with A, E, F, P, or S. The modifications replace one or more
14 amino acids in the identified regions, without increasing or decreasing the total number of
15 amino acids in the protein.

16 Preferably, the deallergenized patatin protein comprises SEQ ID NO:2 modified
17 by one or more changes, or SEQ ID NO:7 modified by one or more changes. SEQ ID
18 NO:7 differs from wild type SEQ ID NO:2 in that the first 22 amino acids of SEQ ID
19 NO:2 are replaced with EAE (Glu-Ala-Glu). For example, the changes to SEQ ID NO:2
20 or SEQ ID NO:7 can be: the Y corresponding to position 106 of SEQ ID NO:2 or
21 position 87 of SEQ ID NO:7 is replaced with F or A; the I corresponding to position 113
22 of SEQ ID NO:2 or position 94 of SEQ ID NO:7 is replaced with A; the Y corresponding
23 to position 129 of SEQ ID NO:2 or position 110 of SEQ ID NO:7 is replaced with F or
24 A; the K corresponding to position 137 of SEQ ID NO:2 or position 118 of SEQ ID NO:7
25 is replaced with A; the S corresponding to position 184 of SEQ ID NO:2 or position 165
26 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 185 of SEQ ID
27 NO:2 or position 166 of SEQ ID NO:7 is replaced with F or A; the A corresponding to
28 position 188 of SEQ ID NO:2 or position 169 of SEQ ID NO:7 is replaced with S; the T
29 corresponding to position 192 of SEQ ID NO:2 or position 173 of SEQ ID NO:7 is
30 replaced with A or P; the Y corresponding to position 193 of SEQ ID NO:2 or position
31 174 of SEQ ID NO:7 is replaced with F or A; the K corresponding to position 268 of

1 SEQ ID NO:2 or position 249 of SEQ ID NO:7 is replaced with A or E; the T
2 corresponding to position 269 of SEQ ID NO:2 or position 250 of SEQ ID NO:7 is
3 replaced with A; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of
4 SEQ ID NO:7 is replaced with F or A; the K corresponding to position 273 of SEQ ID
5 NO:2 or position 254 of SEQ ID NO:7 is replaced with A; the K corresponding to
6 position 313 of SEQ ID NO:2 or position 294 of SEQ ID NO:7 is replaced with E; the N
7 corresponding to position 314 of SEQ ID NO:2 or position 295 of SEQ ID NO:7 is
8 replaced with A; the N corresponding to position 315 of SEQ ID NO:2 or position 296 of
9 SEQ ID NO:7 is replaced with A; the Y corresponding to position 316 of SEQ ID NO:2
10 or position 297 of SEQ ID NO:7 is replaced with F or A; the Y corresponding to position
11 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F; the K
12 corresponding to position 367 of SEQ ID NO:2 or position 348 of SEQ ID NO:7 is
13 replaced with A; the R corresponding to position 368 of SEQ ID NO:2 or position 349 of
14 SEQ ID NO:7 is replaced with A; the F corresponding to position 369 of SEQ ID NO:2
15 or position 350 of SEQ ID NO:7 is replaced with A; the K corresponding to position 371
16 of SEQ ID NO:2 or position 352 of SEQ ID NO:7 is replaced with A; the L
17 corresponding to position 372 of SEQ ID NO:2 or position 353 of SEQ ID NO:7 is
18 replaced with A; and the L corresponding to position 373 of SEQ ID NO:2 or position
19 354 of SEQ ID NO:7 is replaced with A.

20 More preferably, SEQ ID NO:2 is modified by the following changes or SEQ ID
21 NO:7 is modified by the following changes: the Y corresponding to position 106 of SEQ
22 ID NO:2 or position 87 of SEQ ID NO:7 is replaced with F; the Y corresponding to
23 position 129 of SEQ ID NO:2 or position 110 of SEQ ID NO:7 is replaced with F; the Y
24 corresponding to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is
25 replaced with F; the Y corresponding to position 193 of SEQ ID NO:2 or position 174 of
26 SEQ ID NO:7 is replaced with F; the Y corresponding to position 270 of SEQ ID NO:2
27 or position 251 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 316
28 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is replaced with F; and the Y
29 corresponding to position 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is
30 replaced with F.

Most preferably, SEQ ID NO:2 is modified by the following changes or SEQ ID NO:7 is modified by the following changes: the Y corresponding to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 316 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is replaced with F; and the Y corresponding to position 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

Nucleic acids

An additional embodiment of the invention is an isolated nucleic acid molecule segment comprising a structural nucleic acid sequence which encodes a deallergenized patatin protein.

The structural nucleic acid sequence can generally encode any deallergenized patatin protein. The structural nucleic acid sequence preferably encodes a deallergenized patatin protein comprising SEQ ID NO:2 modified in one or more of the following regions, or SEQ ID NO:7 modified in one or more of the following regions. The single or multiple amino acid modifications reduce the binding of the modified protein relative to the binding of the corresponding unmodified protein. The regions for modification include amino acid positions 104-113 of SEQ ID NO:2 (85-94 of SEQ ID NO:7), 128-137 of SEQ ID NO:2 (109-118 of SEQ ID NO:7), 184-197 of SEQ ID NO:2 (165-178 of SEQ ID NO:7), 264-277 of SEQ ID NO:2 (245-258 of SEQ ID NO:7), 316-325 of SEQ ID NO:2 (297-306 of SEQ ID NO:7), and 360-377 of SEQ ID NO:2 (341-358 of SEQ ID NO:7). The possible amino acid modifications include replacing an amino acid with A, E, F, P, or S. The modifications replace one or more amino acids in the identified regions, without increasing or decreasing the total number of amino acids in the protein.

Alternatively, the structural nucleic acid sequence encodes SEQ ID NO:2 modified by one or more of the following changes or encoding SEQ ID NO:7 modified by one or more of the following changes: the Y corresponding to position 106 of SEQ ID NO:2 or position 87 of SEQ ID NO:7 is replaced with F or A; the I corresponding to position 113 of SEQ ID NO:2 or position 94 of SEQ ID NO:7 is replaced with A; the Y

1 corresponding to position 129 of SEQ ID NO:2 or position 110 of SEQ ID NO:7 is
2 replaced with F or A; the K corresponding to position 137 of SEQ ID NO:2 or position
3 118 of SEQ ID NO:7 is replaced with A; the S corresponding to position 184 of SEQ ID
4 NO:2 or position 165 of SEQ ID NO:7 is replaced with A; the Y corresponding to
5 position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F or A;
6 the A corresponding to position 188 of SEQ ID NO:2 or position 169 of SEQ ID NO:7 is
7 replaced with S; the T corresponding to position 192 of SEQ ID NO:2 or position 173 of
8 SEQ ID NO:7 is replaced with A or P; the Y corresponding to position 193 of SEQ ID
9 NO:2 or position 174 of SEQ ID NO:7 is replaced with F or A; the K corresponding to
10 position 268 of SEQ ID NO:2 or position 249 of SEQ ID NO:7 is replaced with A or E;
11 the T corresponding to position 269 of SEQ ID NO:2 or position 250 of SEQ ID NO:7 is
12 replaced with A; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of
13 SEQ ID NO:7 is replaced with F or A; the K corresponding to position 273 of SEQ ID
14 NO:2 or position 254 of SEQ ID NO:7 is replaced with A; the K corresponding to
15 position 313 of SEQ ID NO:2 or position 294 of SEQ ID NO:7 is replaced with E; the N
16 corresponding to position 314 of SEQ ID NO:2 or position 295 of SEQ ID NO:7 is
17 replaced with A; the N corresponding to position 315 of SEQ ID NO:2 or position 296 of
18 SEQ ID NO:7 is replaced with A; the Y corresponding to position 316 of SEQ ID NO:2
19 or position 297 of SEQ ID NO:7 is replaced with F or A; the Y corresponding to position
20 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F; the K
21 corresponding to position 367 of SEQ ID NO:2 or position 348 of SEQ ID NO:7 is
22 replaced with A; the R corresponding to position 368 of SEQ ID NO:2 or position 349 of
23 SEQ ID NO:7 is replaced with A; the F corresponding to position 369 of SEQ ID NO:2
24 or position 350 of SEQ ID NO:7 is replaced with A; the K corresponding to position 371
25 of SEQ ID NO:2 or position 352 of SEQ ID NO:7 is replaced with A; the L
26 corresponding to position 372 of SEQ ID NO:2 or position 353 of SEQ ID NO:7 is
27 replaced with A; and the L corresponding to position 373 of SEQ ID NO:2 or position
28 354 of SEQ ID NO:7 is replaced with A.

29 More preferably, the structural nucleic acid sequence encodes SEQ ID NO:2
30 modified by the following changes or SEQ ID NO:7 modified by the following changes:
31 the Y corresponding to position 106 of SEQ ID NO:2 or position 87 of SEQ ID NO:7 is

1 replaced with F; the Y corresponding to position 129 of SEQ ID NO:2 or position 110 of
2 SEQ ID NO:7 is replaced with F; the Y corresponding to position 185 of SEQ ID NO:2
3 or position 166 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 193
4 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is replaced with F; the Y
5 corresponding to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is
6 replaced with F; the Y corresponding to position 316 of SEQ ID NO:2 or position 297 of
7 SEQ ID NO:7 is replaced with F; and the Y corresponding to position 362 of SEQ ID
8 NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

9 Most preferably, the structural nucleic acid sequence encodes SEQ ID NO:2
10 modified by the following changes or SEQ ID NO:7 modified by the following changes:
11 the Y corresponding to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is
12 replaced with F; the Y corresponding to position 193 of SEQ ID NO:2 or position 174 of
13 SEQ ID NO:7 is replaced with F; the Y corresponding to position 270 of SEQ ID NO:2
14 or position 251 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 316
15 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is replaced with F; and the Y
16 corresponding to position 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is
17 replaced with F.

18 Recombinant vectors

19 An additional embodiment is directed towards recombinant vectors comprising a
20 structural nucleic acid sequence which encodes a deallergenized patatin protein. The
21 recombinant vector comprises operatively linked in the 5' to 3' orientation: a promoter
22 that directs transcription of a structural nucleic acid sequence; a structural nucleic acid
23 sequence, and a 3' transcription terminator.

24 The structural nucleic acid sequence can encode SEQ ID NO:2 modified in one or
25 more of the following regions, or SEQ ID NO:7 modified in one or more of the following
26 regions. The single or multiple amino acid modifications reduce the binding of the
27 modified protein relative to the binding of the corresponding unmodified protein. The
28 regions for modification include amino acid positions 104-113 of SEQ ID NO:2 (85-94
29 of SEQ ID NO:7), 128-137 of SEQ ID NO:2 (109-118 of SEQ ID NO:7), 184-197 of
30 SEQ ID NO:2 (165-178 of SEQ ID NO:7), 264-277 of SEQ ID NO:2 (245-258 of SEQ

1 ID NO:7), 316-325 of SEQ ID NO:2 (297-306 of SEQ ID NO:7), and 360-377 of SEQ ID
2 NO:2 (341-358 of SEQ ID NO:7). The possible amino acid modifications include
3 replacing an amino acid with A, E, F, P, or S. The modifications replace one or more
4 amino acids in the identified regions, without increasing or decreasing the total number of
5 amino acids in the protein.

6 Alternatively, the recombinant vector comprises operatively linked in the 5' to 3'
7 orientation: a promoter that directs transcription of a structural nucleic acid sequence; a
8 structural nucleic acid sequence encoding SEQ ID NO:2 modified by one or more of the
9 following changes or encoding SEQ ID NO:7 modified by one or more of the following
10 changes: the Y corresponding to position 106 of SEQ ID NO:2 or position 87 of SEQ ID
11 NO:7 is replaced with F or A; the I corresponding to position 113 of SEQ ID NO:2 or
12 position 94 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 129 of
13 SEQ ID NO:2 or position 110 of SEQ ID NO:7 is replaced with F or A; the K
14 corresponding to position 137 of SEQ ID NO:2 or position 118 of SEQ ID NO:7 is
15 replaced with A; the S corresponding to position 184 of SEQ ID NO:2 or position 165 of
16 SEQ ID NO:7 is replaced with A; the Y corresponding to position 185 of SEQ ID NO:2
17 or position 166 of SEQ ID NO:7 is replaced with F or A; the A corresponding to position
18 188 of SEQ ID NO:2 or position 169 of SEQ ID NO:7 is replaced with S; the T
19 corresponding to position 192 of SEQ ID NO:2 or position 173 of SEQ ID NO:7 is
20 replaced with A or P; the Y corresponding to position 193 of SEQ ID NO:2 or position
21 174 of SEQ ID NO:7 is replaced with F or A; the K corresponding to position 268 of
22 SEQ ID NO:2 or position 249 of SEQ ID NO:7 is replaced with A or E; the T
23 corresponding to position 269 of SEQ ID NO:2 or position 250 of SEQ ID NO:7 is
24 replaced with A; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of
25 SEQ ID NO:7 is replaced with F or A; the K corresponding to position 273 of SEQ ID
26 NO:2 or position 254 of SEQ ID NO:7 is replaced with A; the K corresponding to
27 position 313 of SEQ ID NO:2 or position 294 of SEQ ID NO:7 is replaced with E; the N
28 corresponding to position 314 of SEQ ID NO:2 or position 295 of SEQ ID NO:7 is
29 replaced with A; the N corresponding to position 315 of SEQ ID NO:2 or position 296 of
30 SEQ ID NO:7 is replaced with A; the Y corresponding to position 316 of SEQ ID NO:2
31 or position 297 of SEQ ID NO:7 is replaced with F or A; the Y corresponding to position

1 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F; the K
2 corresponding to position 367 of SEQ ID NO:2 or position 348 of SEQ ID NO:7 is
3 replaced with A; the R corresponding to position 368 of SEQ ID NO:2 or position 349 of
4 SEQ ID NO:7 is replaced with A; the F corresponding to position 369 of SEQ ID NO:2
5 or position 350 of SEQ ID NO:7 is replaced with A; the K corresponding to position 371
6 of SEQ ID NO:2 or position 352 of SEQ ID NO:7 is replaced with A; the L
7 corresponding to position 372 of SEQ ID NO:2 or position 353 of SEQ ID NO:7 is
8 replaced with A; and the L corresponding to position 373 of SEQ ID NO:2 or position
9 354 of SEQ ID NO:7 is replaced with A; and a 3' transcription terminator.

10 Most preferably, the vector comprises a structural nucleic acid sequence encoding
11 SEQ ID NO:2 modified by the following changes or SEQ ID NO:7 modified by the
12 following changes: the Y corresponding to position 106 of SEQ ID NO:2 or position 87
13 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 129 of SEQ ID
14 NO:2 or position 110 of SEQ ID NO:7 is replaced with F; the Y corresponding to
15 position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F; the Y
16 corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is
17 replaced with F; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of
18 SEQ ID NO:7 is replaced with F; the Y corresponding to position 316 of SEQ ID NO:2
19 or position 297 of SEQ ID NO:7 is replaced with F; and the Y corresponding to position
20 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

21 Most preferably, the vector comprises a structural nucleic acid sequence encoding
22 SEQ ID NO:2 modified by the following changes or SEQ ID NO:7 modified by the
23 following changes: the Y corresponding to position 185 of SEQ ID NO:2 or position 166
24 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 193 of SEQ ID
25 NO:2 or position 174 of SEQ ID NO:7 is replaced with F; the Y corresponding to
26 position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is replaced with F; the Y
27 corresponding to position 316 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is
28 replaced with F; and the Y corresponding to position 362 of SEQ ID NO:2 or position
29 343 of SEQ ID NO:7 is replaced with F.

Recombinant host cells

A further embodiment of the invention is directed towards recombinant host cells comprising a structural nucleic acid sequence encoding a deallergenized patatin protein. The recombinant host cell preferably produces a deallergenized patatin protein. More preferably, the recombinant host cell produces a deallergenized patatin protein in a concentration sufficient to inhibit growth or to kill an insect which ingests the recombinant host cell. The recombinant host cell can generally comprise any structural nucleic acid sequence encoding a deallergenized patatin protein.

The recombinant host cell can comprise a structural nucleic acid sequence encoding SEQ ID NO:2 modified in one or more of the following regions, or SEQ ID NO:7 modified in one or more of the following regions. The single or multiple amino acid modifications reduce the binding of the modified protein relative to the binding of the corresponding unmodified protein. The regions for modification include amino acid positions 104-113 of SEQ ID NO:2 (85-94 of SEQ ID NO:7), 128-137 of SEQ ID NO:2 (109-118 of SEQ ID NO:7), 184-197 of SEQ ID NO:2 (165-178 of SEQ ID NO:7), 264-277 of SEQ ID NO:2 (245-258 of SEQ ID NO:7), 316-325 of SEQ ID NO:2 (297-306 of SEQ ID NO:7), and 360-377 of SEQ ID NO:2 (341-358 of SEQ ID NO:7). The possible amino acid modifications include replacing an amino acid with A, E, F, P, or S. The modifications replace one or more amino acids in the identified regions, without increasing or decreasing the total number of amino acids in the protein.

Alternatively, the recombinant host cell comprises a structural nucleic acid sequence encoding SEQ ID NO:2 modified by one or more of the following changes or encoding SEQ ID NO:7 modified by one or more of the following changes: the Y corresponding to position 106 of SEQ ID NO:2 or position 87 of SEQ ID NO:7 is replaced with F or A; the I corresponding to position 113 of SEQ ID NO:2 or position 94 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 129 of SEQ ID NO:2 or position 110 of SEQ ID NO:7 is replaced with F or A; the K corresponding to position 137 of SEQ ID NO:2 or position 118 of SEQ ID NO:7 is replaced with A; the S corresponding to position 184 of SEQ ID NO:2 or position 165 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F or A; the A corresponding to position 188 of SEQ ID

1 NO:2 or position 169 of SEQ ID NO:7 is replaced with S; the T corresponding to position
2 192 of SEQ ID NO:2 or position 173 of SEQ ID NO:7 is replaced with A or P; the Y
3 corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is
4 replaced with F or A; the K corresponding to position 268 of SEQ ID NO:2 or position
5 249 of SEQ ID NO:7 is replaced with A or E; the T corresponding to position 269 of
6 SEQ ID NO:2 or position 250 of SEQ ID NO:7 is replaced with A; the Y corresponding
7 to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is replaced with F or
8 A; the K corresponding to position 273 of SEQ ID NO:2 or position 254 of SEQ ID NO:7
9 is replaced with A; the K corresponding to position 313 of SEQ ID NO:2 or position 294
10 of SEQ ID NO:7 is replaced with E; the N corresponding to position 314 of SEQ ID
11 NO:2 or position 295 of SEQ ID NO:7 is replaced with A; the N corresponding to
12 position 315 of SEQ ID NO:2 or position 296 of SEQ ID NO:7 is replaced with A; the Y
13 corresponding to position 316 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is
14 replaced with F or A; the Y corresponding to position 362 of SEQ ID NO:2 or position
15 343 of SEQ ID NO:7 is replaced with F; the K corresponding to position 367 of SEQ ID
16 NO:2 or position 348 of SEQ ID NO:7 is replaced with A; the R corresponding to
17 position 368 of SEQ ID NO:2 or position 349 of SEQ ID NO:7 is replaced with A; the F
18 corresponding to position 369 of SEQ ID NO:2 or position 350 of SEQ ID NO:7 is
19 replaced with A; the K corresponding to position 371 of SEQ ID NO:2 or position 352 of
20 SEQ ID NO:7 is replaced with A; the L corresponding to position 372 of SEQ ID NO:2
21 or position 353 of SEQ ID NO:7 is replaced with A; and the L corresponding to position
22 373 of SEQ ID NO:2 or position 354 of SEQ ID NO:7 is replaced with A.

23 More preferably, the recombinant host cell comprises a structural nucleic acid
24 sequence encoding SEQ ID NO:2 modified by the following changes or SEQ ID NO:7
25 modified by the following changes: the Y corresponding to position 106 of SEQ ID NO:2
26 or position 87 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 129 of
27 SEQ ID NO:2 or position 110 of SEQ ID NO:7 is replaced with F; the Y corresponding
28 to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F; the
29 Y corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is
30 replaced with F; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of
31 SEQ ID NO:7 is replaced with F; the Y corresponding to position 316 of SEQ ID NO:2

1 or position 297 of SEQ ID NO:7 is replaced with F; and the Y corresponding to position
2 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

3 Most preferably, the recombinant host cell comprises a structural nucleic acid
4 sequence encoding SEQ ID NO:2 modified by the following changes or SEQ ID NO:7
5 modified by the following changes: the Y corresponding to position 185 of SEQ ID NO:2
6 or position 166 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 193
7 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is replaced with F; the Y
8 corresponding to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is
9 replaced with F; the Y corresponding to position 316 of SEQ ID NO:2 or position 297 of
10 SEQ ID NO:7 is replaced with F; and the Y corresponding to position 362 of SEQ ID
11 NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

12 The recombinant host cell can generally be any type of host cell, and preferably is
13 a bacterial, fungal, or plant cell. The bacterial cell is preferably an *Escherichia coli*
14 bacterial cell. The fungal cell is preferably a *Saccharomyces cerevisiae*,
15 *Schizosaccharomyces pombe*, or *Pichia pastoris* fungal cell. The plant cell can be a
16 monocot, dicot, or conifer plant cell. The plant cell is preferably an alfalfa, banana,
17 canola, corn, cotton, cucumber, peanut, potato, rice, soybean, sunflower, sweet potato,
18 tobacco, tomato, or wheat plant cell. The recombinant host cell preferably further
19 comprises operatively linked to the structural nucleic acid sequence a promoter that
20 directs transcription of the structural nucleic acid sequence. The recombinant host cell
21 preferably further comprises operatively linked to the structural nucleic acid sequence a
22 3' transcription terminator and a polyadenylation site.

23 Recombinant plants

24 An additional embodiment of the invention is a recombinant plant comprising a
25 structural nucleic acid sequence encoding a deallergenized patatin protein. The
26 recombinant plant preferably produces a deallergenized patatin protein. More preferably,
27 the recombinant plant produces a deallergenized patatin protein in a concentration
28 sufficient to inhibit growth or to kill an insect which ingests plant tissue from the
29 recombinant plant.

1 The recombinant plant can comprise a structural nucleic acid sequence encoding
2 SEQ ID NO:2 modified in one or more of the following regions, or SEQ ID NO:7
3 modified in one or more of the following regions. The single or multiple amino acid
4 modifications reduce the binding of the modified protein relative to the binding of the
5 corresponding unmodified protein. The regions for modification include amino acid
6 positions 104-113 of SEQ ID NO:2 (85-94 of SEQ ID NO:7), 128-137 of SEQ ID NO:2
7 (109-118 of SEQ ID NO:7), 184-197 of SEQ ID NO:2 (165-178 of SEQ ID NO:7), 264-
8 277 of SEQ ID NO:2 (245-258 of SEQ ID NO:7), 316-325 of SEQ ID NO:2 (297-306 of
9 SEQ ID NO:7), and 360-377 of SEQ ID NO:2 (341-358 of SEQ ID NO:7). The possible
10 amino acid modifications include replacing an amino acid with A, E, F, P, or S. The
11 modifications replace one or more amino acids in the identified regions, without
12 increasing or decreasing the total number of amino acids in the protein.

13 Alternatively, the recombinant plant can comprise a structural nucleic acid
14 sequence encoding SEQ ID NO:2 modified by one or more of the following changes or
15 encoding SEQ ID NO:7 modified by one or more of the following changes: the Y
16 corresponding to position 106 of SEQ ID NO:2 or position 87 of SEQ ID NO:7 is
17 replaced with F or A; the I corresponding to position 113 of SEQ ID NO:2 or position 94
18 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 129 of SEQ ID
19 NO:2 or position 110 of SEQ ID NO:7 is replaced with F or A; the K corresponding to
20 position 137 of SEQ ID NO:2 or position 118 of SEQ ID NO:7 is replaced with A; the S
21 corresponding to position 184 of SEQ ID NO:2 or position 165 of SEQ ID NO:7 is
22 replaced with A; the Y corresponding to position 185 of SEQ ID NO:2 or position 166 of
23 SEQ ID NO:7 is replaced with F or A; the A corresponding to position 188 of SEQ ID
24 NO:2 or position 169 of SEQ ID NO:7 is replaced with S; the T corresponding to position
25 192 of SEQ ID NO:2 or position 173 of SEQ ID NO:7 is replaced with A or P; the Y
26 corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is
27 replaced with F or A; the K corresponding to position 268 of SEQ ID NO:2 or position
28 249 of SEQ ID NO:7 is replaced with A or E; the T corresponding to position 269 of
29 SEQ ID NO:2 or position 250 of SEQ ID NO:7 is replaced with A; the Y corresponding
30 to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is replaced with F or
31 A; the K corresponding to position 273 of SEQ ID NO:2 or position 254 of SEQ ID NO:7

1 is replaced with A; the K corresponding to position 313 of SEQ ID NO:2 or position 294
2 of SEQ ID NO:7 is replaced with E; the N corresponding to position 314 of SEQ ID
3 NO:2 or position 295 of SEQ ID NO:7 is replaced with A; the N corresponding to
4 position 315 of SEQ ID NO:2 or position 296 of SEQ ID NO:7 is replaced with A; the Y
5 corresponding to position 316 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is
6 replaced with F or A; the Y corresponding to position 362 of SEQ ID NO:2 or position
7 343 of SEQ ID NO:7 is replaced with F; the K corresponding to position 367 of SEQ ID
8 NO:2 or position 348 of SEQ ID NO:7 is replaced with A; the R corresponding to
9 position 368 of SEQ ID NO:2 or position 349 of SEQ ID NO:7 is replaced with A; the F
10 corresponding to position 369 of SEQ ID NO:2 or position 350 of SEQ ID NO:7 is
11 replaced with A; the K corresponding to position 371 of SEQ ID NO:2 or position 352 of
12 SEQ ID NO:7 is replaced with A; the L corresponding to position 372 of SEQ ID NO:2
13 or position 353 of SEQ ID NO:7 is replaced with A; and the L corresponding to position
14 373 of SEQ ID NO:2 or position 354 of SEQ ID NO:7 is replaced with A.

15 More preferably, the recombinant plant comprises a structural nucleic acid
16 sequence encoding SEQ ID NO:2 modified by the following changes or SEQ ID NO:7
17 modified by the following changes: the Y corresponding to position 106 of SEQ ID NO:2
18 or position 87 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 129 of
19 SEQ ID NO:2 or position 110 of SEQ ID NO:7 is replaced with F; the Y corresponding
20 to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F; the
21 Y corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is
22 replaced with F; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of
23 SEQ ID NO:7 is replaced with F; the Y corresponding to position 316 of SEQ ID NO:2
24 or position 297 of SEQ ID NO:7 is replaced with F; and the Y corresponding to position
25 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

26 Most preferably, the recombinant plant comprises a structural nucleic acid
27 sequence encoding SEQ ID NO:2 modified by the following changes or SEQ ID NO:7
28 modified by the following changes: the Y corresponding to position 185 of SEQ ID NO:2
29 or position 166 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 193
30 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is replaced with F; the Y
31 corresponding to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is

1 replaced with F; the Y corresponding to position 316 of SEQ ID NO:2 or position 297 of
2 SEQ ID NO:7 is replaced with F; and the Y corresponding to position 362 of SEQ ID
3 NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

4 The recombinant plant can generally be any type of plant. The plant can be a
5 monocot, dicot, or conifer plant. The plant is preferably an alfalfa, banana, canola, corn,
6 cotton, cucumber, peanut, potato, rice, soybean, sunflower, sweet potato, tobacco,
7 tomato, or wheat plant.

8 The recombinant plant preferably further comprises operatively linked to the
9 structural nucleic acid sequence a promoter that directs transcription of the structural
10 nucleic acid sequence. The recombinant plant preferably further comprises operatively
11 linked to the structural nucleic acid sequence a 3' transcription terminator and a
12 polyadenylation site.

13 Methods of preparation

14 Embodiments of the invention are further directed towards methods of preparing
15 recombinant host cells and recombinant plants useful for the production of deallergenized
16 patatin proteins.

17 A method of preparing a recombinant host cell useful for the production of
18 deallergenized patatin proteins can comprise selecting a host cell; transforming the host
19 cell with a recombinant vector; and obtaining recombinant host cells.

20 The recombinant vector comprises a structural nucleic acid sequence encoding
21 SEQ ID NO:2 modified in one or more of the following regions, or SEQ ID NO:7
22 modified in one or more of the following regions. The single or multiple amino acid
23 modifications reduce the binding of the modified protein relative to the binding of the
24 corresponding unmodified protein. The regions for modification include amino acid
25 positions 104-113 of SEQ ID NO:2 (85-94 of SEQ ID NO:7), 128-137 of SEQ ID NO:2
26 (109-118 of SEQ ID NO:7), 184-197 of SEQ ID NO:2 (165-178 of SEQ ID NO:7), 264-
27 277 of SEQ ID NO:2 (245-258 of SEQ ID NO:7), 316-325 of SEQ ID NO:2 (297-306 of
28 SEQ ID NO:7), and 360-377 of SEQ ID NO:2 (341-358 of SEQ ID NO:7). The possible
29 amino acid modifications include replacing an amino acid with A, E, F, P, or S. The

1 modifications replace one or more amino acids in the identified regions, without
2 increasing or decreasing the total number of amino acids in the protein.

3 Alternatively, the recombinant vector comprises a structural nucleic acid sequence
4 encoding SEQ ID NO:2 modified by one or more of the following changes or encoding
5 SEQ ID NO:7 modified by one or more of the following changes: the Y corresponding to
6 position 106 of SEQ ID NO:2 or position 87 of SEQ ID NO:7 is replaced with F or A; the
7 I corresponding to position 113 of SEQ ID NO:2 or position 94 of SEQ ID NO:7 is
8 replaced with A; the Y corresponding to position 129 of SEQ ID NO:2 or position 110 of
9 SEQ ID NO:7 is replaced with F or A; the K corresponding to position 137 of SEQ ID
10 NO:2 or position 118 of SEQ ID NO:7 is replaced with A; the S corresponding to
11 position 184 of SEQ ID NO:2 or position 165 of SEQ ID NO:7 is replaced with A; the Y
12 corresponding to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is
13 replaced with F or A; the A corresponding to position 188 of SEQ ID NO:2 or position
14 169 of SEQ ID NO:7 is replaced with S; the T corresponding to position 192 of SEQ ID
15 NO:2 or position 173 of SEQ ID NO:7 is replaced with A or P; the Y corresponding to
16 position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is replaced with F or A;
17 the K corresponding to position 268 of SEQ ID NO:2 or position 249 of SEQ ID NO:7 is
18 replaced with A or E; the T corresponding to position 269 of SEQ ID NO:2 or position
19 250 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 270 of SEQ ID
20 NO:2 or position 251 of SEQ ID NO:7 is replaced with F or A; the K corresponding to
21 position 273 of SEQ ID NO:2 or position 254 of SEQ ID NO:7 is replaced with A; the K
22 corresponding to position 313 of SEQ ID NO:2 or position 294 of SEQ ID NO:7 is
23 replaced with E; the N corresponding to position 314 of SEQ ID NO:2 or position 295 of
24 SEQ ID NO:7 is replaced with A; the N corresponding to position 315 of SEQ ID NO:2
25 or position 296 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 316
26 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is replaced with F or A; the Y
27 corresponding to position 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is
28 replaced with F; the K corresponding to position 367 of SEQ ID NO:2 or position 348 of
29 SEQ ID NO:7 is replaced with A; the R corresponding to position 368 of SEQ ID NO:2
30 or position 349 of SEQ ID NO:7 is replaced with A; the F corresponding to position 369
31 of SEQ ID NO:2 or position 350 of SEQ ID NO:7 is replaced with A; the K

1 corresponding to position 371 of SEQ ID NO:2 or position 352 of SEQ ID NO:7 is
2 replaced with A; the L corresponding to position 372 of SEQ ID NO:2 or position 353 of
3 SEQ ID NO:7 is replaced with A; and the L corresponding to position 373 of SEQ ID
4 NO:2 or position 354 of SEQ ID NO:7 is replaced with A.

5 More preferably, the vector comprises a structural nucleic acid sequence encoding
6 SEQ ID NO:2 modified by the following changes or SEQ ID NO:7 modified by the
7 following changes: the Y corresponding to position 106 of SEQ ID NO:2 or position 87
8 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 129 of SEQ ID
9 NO:2 or position 110 of SEQ ID NO:7 is replaced with F; the Y corresponding to
10 position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F; the Y
11 corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is
12 replaced with F; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of
13 SEQ ID NO:7 is replaced with F; the Y corresponding to position 316 of SEQ ID NO:2
14 or position 297 of SEQ ID NO:7 is replaced with F; and the Y corresponding to position
15 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

16 Most preferably, the vector comprises a structural nucleic acid sequence encoding
17 SEQ ID NO:2 modified by the following changes or SEQ ID NO:7 modified by the
18 following changes: the Y corresponding to position 185 of SEQ ID NO:2 or position 166
19 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 193 of SEQ ID
20 NO:2 or position 174 of SEQ ID NO:7 is replaced with F; the Y corresponding to
21 position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is replaced with F; the Y
22 corresponding to position 316 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is
23 replaced with F; and the Y corresponding to position 362 of SEQ ID NO:2 or position
24 343 of SEQ ID NO:7 is replaced with F.

25 The method can generally be used to prepare any type of recombinant host cell.
26 Preferably, the method can be used to prepare a recombinant bacterial cell, a recombinant
27 fungal cell, or a recombinant plant cell. The bacterial cell is preferably an *Escherichia*
28 *coli* bacterial cell. The fungal cell is preferably a *Saccharomyces cerevisiae*,
29 *Schizosaccharomyces pombe*, or *Pichia pastoris* fungal cell. The plant cell can be a
30 monocot, dicot, or conifer plant cell. The plant cell is preferably an alfalfa, banana,

1 canola, corn, cotton, cucumber, peanut, potato, rice, soybean, sunflower, sweet potato,
2 tobacco, tomato, or wheat plant cell.

3 An additional embodiment is directed towards methods for the preparation of
4 recombinant plants useful for the production of deallergenized patatin proteins. The
5 method can comprise selecting a host plant cell; transforming the host plant cell with a
6 recombinant vector; obtaining recombinant host cells; and regenerating a recombinant
7 plant from the recombinant host plant cells.

8 The recombinant vector comprises a structural nucleic acid sequence encoding
9 SEQ ID NO:2 modified in one or more of the following regions, or SEQ ID NO:7
10 modified in one or more of the following regions. The single or multiple amino acid
11 modifications reduce the binding of the modified protein relative to the binding of the
12 corresponding unmodified protein. The regions for modification include amino acid
13 positions 104-113 of SEQ ID NO:2 (85-94 of SEQ ID NO:7), 128-137 of SEQ ID NO:2
14 (109-118 of SEQ ID NO:7), 184-197 of SEQ ID NO:2 (165-178 of SEQ ID NO:7), 264-
15 277 of SEQ ID NO:2 (245-258 of SEQ ID NO:7), 316-325 of SEQ ID NO:2 (297-306 of
16 SEQ ID NO:7), and 360-377 of SEQ ID NO:2 (341-358 of SEQ ID NO:7). The possible
17 amino acid modifications include replacing an amino acid with A, E, F, P, or S. The
18 modifications replace one or more amino acids in the identified regions, without
19 increasing or decreasing the total number of amino acids in the protein.

20 Alternatively, the recombinant vector comprises a structural nucleic acid sequence
21 encoding SEQ ID NO:2 modified by one or more of the following changes or encoding
22 SEQ ID NO:7 modified by one or more of the following changes: the Y corresponding to
23 position 106 of SEQ ID NO:2 or position 87 of SEQ ID NO:7 is replaced with F or A; the
24 I corresponding to position 113 of SEQ ID NO:2 or position 94 of SEQ ID NO:7 is
25 replaced with A; the Y corresponding to position 129 of SEQ ID NO:2 or position 110 of
26 SEQ ID NO:7 is replaced with F or A; the K corresponding to position 137 of SEQ ID
27 NO:2 or position 118 of SEQ ID NO:7 is replaced with A; the S corresponding to
28 position 184 of SEQ ID NO:2 or position 165 of SEQ ID NO:7 is replaced with A; the Y
29 corresponding to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is
30 replaced with F or A; the A corresponding to position 188 of SEQ ID NO:2 or position
31 169 of SEQ ID NO:7 is replaced with S; the T corresponding to position 192 of SEQ ID

NO:2 or position 173 of SEQ ID NO:7 is replaced with A or P; the Y corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is replaced with F or A; the K corresponding to position 268 of SEQ ID NO:2 or position 249 of SEQ ID NO:7 is replaced with A or E; the T corresponding to position 269 of SEQ ID NO:2 or position 250 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is replaced with F or A; the K corresponding to position 273 of SEQ ID NO:2 or position 254 of SEQ ID NO:7 is replaced with A; the K corresponding to position 313 of SEQ ID NO:2 or position 294 of SEQ ID NO:7 is replaced with E; the N corresponding to position 314 of SEQ ID NO:2 or position 295 of SEQ ID NO:7 is replaced with A; the N corresponding to position 315 of SEQ ID NO:2 or position 296 of SEQ ID NO:7 is replaced with A; the Y corresponding to position 316 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is replaced with F or A; the Y corresponding to position 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F; the K corresponding to position 367 of SEQ ID NO:2 or position 348 of SEQ ID NO:7 is replaced with A; the R corresponding to position 368 of SEQ ID NO:2 or position 349 of SEQ ID NO:7 is replaced with A; the F corresponding to position 369 of SEQ ID NO:2 or position 350 of SEQ ID NO:7 is replaced with A; the K corresponding to position 371 of SEQ ID NO:2 or position 352 of SEQ ID NO:7 is replaced with A; the L corresponding to position 372 of SEQ ID NO:2 or position 353 of SEQ ID NO:7 is replaced with A; and the L corresponding to position 373 of SEQ ID NO:2 or position 354 of SEQ ID NO:7 is replaced with A.

More preferably, the vector comprises a structural nucleic acid sequence encoding SEQ ID NO:2 modified by the following changes or SEQ ID NO:7 modified by the following changes: the Y corresponding to position 106 of SEQ ID NO:2 or position 87 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 129 of SEQ ID NO:2 or position 110 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 316 of SEQ ID NO:2

or position 297 of SEQ ID NO:7 is replaced with F; and the Y corresponding to position 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

Most preferably, the vector comprises a structural nucleic acid sequence encoding SEQ ID NO:2 modified by the following changes or SEQ ID NO:7 modified by the following changes: the Y corresponding to position 185 of SEQ ID NO:2 or position 166 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 193 of SEQ ID NO:2 or position 174 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 270 of SEQ ID NO:2 or position 251 of SEQ ID NO:7 is replaced with F; the Y corresponding to position 316 of SEQ ID NO:2 or position 297 of SEQ ID NO:7 is replaced with F; and the Y corresponding to position 362 of SEQ ID NO:2 or position 343 of SEQ ID NO:7 is replaced with F.

The recombinant plant can generally be any type of plant. The plant can be a monocot, dicot, or conifer plant. The plant is preferably an alfalfa, banana, canola, corn, cotton, cucumber, peanut, potato, rice, soybean, sunflower, sweet potato, tobacco, tomato, or wheat plant.

Deallergenized patatin proteins can be prepared by isolating the deallergenized patatin protein from any one of the above described host cells or plants.

Deglycosylation

The examples herein provide evidence that glycosylation of can contribute to the allergenicity of a protein. Accordingly, rational substitution of amino acid residues likely to be the targets of glycosylation within a subject allergen protein may reduce or eliminate the allergenic properties of the protein without adversely affecting the enzymatic, insecticidal, antifungal or other functional properties of the protein.

Glycosylation commonly occurs as either N-linked or O-linked forms. N-linked glycosylation usually occurs at the motif Asn-Xaa-Ser/Thr, where Xaa is any amino acid except Pro (Kasturi, L. et al., *Biochem J.* 323: 415-519, 1997; Melquist, J.L. et al., *Biochemistry* 37: 6833-6837, 1998). O-linked glycosylation occurs between the hydroxyl group of serine or threonine and an amino sugar.

Site directed mutagenesis of selected asparagine, serine, or threonine may be used to reduce or eliminate the glycosylation of patatin proteins. A search of SEQ ID NO:2

1 for the Asn-Xaa-Ser/Thr motif reveals one occurrence at amino acid positions 202-204.
2 Mutagenization of the nucleic acid sequence encoding this region may result in a reduced
3 allergenicity of the encoded protein.

4 In order to test this conceptual approach to reducing allergenicity of patatin
5 proteins, two sets of experiments were performed: a) production of patatin proteins in
6 *Escherichia coli*, which do not glycosylate proteins; and b) production of patatin proteins
7 with an N202Q site directed mutation.

8 Antibodies obtained from patients HS-07 and G15-MON (not potato allergic) did
9 not show specific binding to wild type patatin, patatin produced in *E. coli*, or the N202Q
10 variant. Antibodies obtained from patient HS-01 (potato allergic) bound to wild type
11 patatin, but not to patatin produced in *E. coli* or the N202Q variant. Antibodies obtained
12 from patient HS-02 (potato allergic) bound strongly to wild type patatin, but extremely
13 weakly to patatin produced in *E. coli*, and binding to the N202Q variant resembled vector
14 controls. Antibodies obtained from patient HS-03 (potato allergic) bound to wild type
15 patatin, but not to patatin produced in *E. coli* or the N202Q variant. Antibodies obtained
16 from patient HS-05 (potato allergic) bound to wild type patatin, but very weakly to
17 patatin produced in *E. coli* and the N202Q variant. Antibodies obtained from patient HS-
18 06 (potato allergic) strongly bound wild type patatin, the N202Q variant, and to patatin
19 produced in *E. coli*. These results strongly suggest that glycosylation is at least partially
20 responsible for the antigenic properties of patatin proteins, and that site directed
21 mutagenesis may be used to reduce or eliminate specific antibody binding. Mutagenesis
22 at position 202 of SEQ ID NO:2 may be useful for reducing or eliminating specific
23 antibody binding.

24 Permuteins

25 The positions of the internal breakpoints described in the following Examples are
26 found on the protein surface, and are distributed throughout the linear sequence without
27 any obvious bias towards the ends or the middle. Breakpoints occurring below the
28 protein surface can additionally be selected. The rearranged two subunits can be joined
29 by a peptide linker. A preferred embodiment involves the linking of the N-terminal and
30 C-terminal subunits by a three amino acid linker, although linkers of various sizes can be

used. Additionally, the N-terminal and C-terminal subunits can be joined lacking a linker sequence. Furthermore, a portion of the C-terminal subunit can be deleted and the connection made from the truncated C-terminal subunit to the original N-terminal subunit and vice versa as previously described (Yang and Schachman, *Proc. Natl. Acad. Sci. U.S.A.*, 90: 11980-11984, 1993; Viguera, *et al.*, *Mol. Biol.*, 247: 670-681, 1995; Protasova, *et al.*, *Prot. Eng.*, 7: 1373-1377, 1994).

The novel insecticidal proteins of the present invention can be represented by the formula:



wherein;

a is 0 or 1, and if a is 0, then the permutein does not contain a linker sequence;

X¹ is a polypeptide sequence corresponding to amino acids n+1 through J;

X² is a polypeptide corresponding to amino acids 1 through n;

n is an integer ranging from 1 to J-1;

J is an integer greater than n+1; and

L is a linker.

In the formula above, the constituent amino acid residues of the novel insecticidal protein are numbered sequentially 1 through J from the original amino terminus to the original carboxyl terminus. A pair of adjacent amino acids within this protein can be numbered n and n+1 respectively where n is an integer ranging from 1 to J-1. The residue n+1 becomes the new N-terminus of the novel insecticidal protein and the residue n becomes the new C-terminus of the novel insecticidal protein.

For example, a parent protein sequence consisting of 120 amino acids can be selected as a starting point for designing a permutein (J=120). If the breakpoint is selected as being between position 40 and position 41, then n=40. If a linker is selected to join the two subunits, the resulting permutein will have the formula: (amino acids 41-120)-L-(amino acids 1-40). If a linker was not used, the resulting permutein will have the formula: (amino acids 41-120)-(amino acids 1-40).

The length of the amino acid sequence of the linker can be selected empirically, by using structural information, or by using a combination of the two approaches. When

no structural information is available, a small series of linkers can be made whose length can span a range of 0 to 50 Å and whose sequence is chosen in order to be substantially consistent with surface exposure (Hopp and Woods, *Mol. Immunol.*, 20: 483-489, 1983; Kyte and Doolittle, *J. Mol. Biol.*, 157: 105-132, 1982; Lee and Richards, *J. Mol. Biol.*, 55: 379-400, 1971) and the ability to adopt a conformation which does not significantly affect the overall configuration of the protein (Karplus and Schulz, *Naturwissenschaften*, 72: 212-213, 1985). Assuming an average length of 2.0 to 3.8 Å per residue, this would mean the length to test would be between about 0 to about 30 residues, with 0 to about 15 residues being the preferred range. Accordingly, there are many such sequences that vary in length or composition that can serve as linkers with the primary consideration being that they be neither excessively long nor excessively short (Sandhu, *et al.*, *Critical Rev. Biotech.*, 12: 437-467, 1992). If the linker is too long, entropy effects may destabilize the three-dimensional fold and may affect protein folding. If the linker is too short, it may destabilize the molecule due to torsional or steric strain.

Use of the distance between the chain ends, defined as the distance between the C-alpha carbons, can be used to define the length of the sequence to be used, or at least to limit the number of possibilities that can be tested in an empirical selection of linkers. Using the calculated length as a guide, linkers with a range of number of residues (calculated using 2 to 3.8 Å per residue) can be selected. These linkers can be composed of the original sequence, shortened or lengthened as necessary, and when lengthened the additional residues can be chosen to be flexible and hydrophilic as described above; or optionally the original sequence can be substituted for using a series of linkers, one example being Gly-Pro-Gly (SEQ ID NO:277); or optionally a combination of the original sequence and new sequence having the appropriate total length can be used. An alternative short, flexible linker sequence is Gly-Gly-Gly-Ser-Gly-Gly-Gly (SEQ ID NO:276).

Selection of permutein breakpoints

Sequences of novel patatin analogs capable of folding to biologically active molecules can be prepared by appropriate selection of the beginning (amino terminus) and ending (carboxyl terminus) positions from within the original polypeptide chain

1 while optionally using a linker sequence as described above. Amino and carboxyl
2 termini can be selected from within a common stretch of sequence, referred to as a
3 breakpoint region, using the guidelines described below. A novel amino acid sequence is
4 thus generated by selecting amino and carboxyl termini from within the same breakpoint
5 region. In many cases, the selection of the new termini will be such that the original
6 position of the carboxyl terminus immediately preceded that of the amino terminus.
7 However, selections of termini anywhere within the region may result in a functional
8 protein, and that these will effectively lead to either deletions or additions to the amino or
9 carboxyl portions of the new sequence.

10 The primary amino acid sequence of a protein dictates folding to the three-
11 dimensional structure beneficial for expression of its biological function. It is possible to
12 obtain and interpret three-dimensional structural information using x-ray diffraction of
13 single protein crystals or nuclear magnetic resonance spectroscopy of protein solutions.
14 Examples of structural information that are relevant to the identification of breakpoint
15 regions include the location and type of protein secondary structure (alpha and 3-10
16 helices, parallel and anti-parallel beta sheets, chain reversals and turns, and loops
17 (Kabsch and Sander, *Biopolymers*, 22: 2577-2637, 1983), the degree of solvent exposure
18 of amino acid residues, the extent and type of interactions of residues with one another
19 (Chothia, C., *Ann. Rev. Biochem.*, 53: 537-572, 1984), and the static and dynamic
20 distribution of conformations along the polypeptide chain (Alber and Mathews, *Methods*
21 *Enzymol.*, 154: 511-533, 1987). In some cases additional information is known about
22 solvent exposure of residues, one example is a site of post-translational attachment of
23 carbohydrate which is necessarily on the surface of the protein. When experimental
24 structural information is not available, or when it is not feasible to obtain the information,
25 methods are available to analyze the primary amino acid sequence in order to make
26 predictions of protein secondary and tertiary structure, solvent accessibility and the
27 occurrence of turns and loops (Fasman, G., Ed. Plenum, New York, 1989; Robson, B.
28 and Garnier, J. *Nature* 361: 506, 1993).

29 Biochemical methods can be applicable for empirically determining surface
30 exposure when direct structural methods are not feasible; for example, using the
31 identification of sites of chain scission following limited proteolysis in order to infer

1 surface exposure (Gentile, F. and Salvatore, G., *Eur. J. Biochem.*, 218: 603-621, 1993).
2 Thus, using either the experimentally derived structural information or predictive
3 methods (Srinivasan, R. and Rose, G.D. *Proteins*, 22: 81-99, 1995), the parental amino
4 acid sequence can be analyzed to classify regions according to whether or not they are
5 integral to the maintenance of secondary and tertiary structure. The sequences within
6 regions that are known to be involved in periodic secondary structure (alpha and 3-10
7 helices, parallel and anti-parallel beta sheets) are regions that should be avoided.
8 Similarly, regions of amino acid sequence that are observed or predicted to have a low
9 degree of solvent exposure are more likely to be part of the so-called hydrophobic core of
10 the protein and should also be avoided for selection of amino and carboxyl termini.
11 Regions that are known or predicted to be in surface turns or loops, and especially those
12 regions that are known not to be required for biological activity, can be preferred sites for
13 new amino and carboxyl termini. Stretches of amino acid sequence that are preferred
14 based on the above criteria can be selected as breakpoint regions.

15 An embodiment of the invention is directed towards patatin permutein proteins.
16 The permutein proteins preferably maintain esterase activity and insecticidal properties.
17 The permutein proteins preferably are less allergenic than the wild type patatin protein to
18 individuals or animals allergic to potatoes. This can be assayed by the binding of
19 antibodies to the wild type patatin and patatin permutein proteins.

20 The permutein proteins can optionally contain a linker sequence. The linker can
21 generally be any amino acid sequence, preferably is Gly-Gly-Gly-Ser-Gly-Gly-Gly (SEQ
22 ID NO:276) or Gly-Pro-Gly (SEQ ID NO:277), and more preferably is Gly-Pro-Gly
23 (SEQ ID NO:277). Specific permutein proteins comprise: (amino acids 247-386 of SEQ
24 ID NO:2)-linker-(amino acids 24-246 of SEQ ID NO:2), (amino acids 269-386 of SEQ
25 ID NO:2)-linker-(amino acids 24-268 of SEQ ID NO:2), SEQ ID NO:247, and SEQ ID
26 NO:259.

27 Embodiments of the invention also include isolated nucleic acid molecule
28 segments comprising a structural nucleic acid sequence encoding a patatin permutein
29 protein. The encoded permutein protein can generally be any permutein protein, and
30 preferably comprises (amino acids 247-386 of SEQ ID NO:2)-linker-(amino acids 24-246
31 of SEQ ID NO:2), (amino acids 269-386 of SEQ ID NO:2)-linker-(amino acids 24-268 of

SEQ ID NO:2), SEQ ID NO:247, or SEQ ID NO:259. The linker can generally be any amino acid sequence, preferably is Gly-Gly-Gly-Ser-Gly-Gly-Gly (SEQ ID NO:276) or Gly-Pro-Gly (SEQ ID NO:277), and more preferably is Gly-Pro-Gly (SEQ ID NO:277). Alternatively, the encoded patatin permutein protein can lack a linker sequence. The structural nucleic acid sequence is preferably SEQ ID NO:246 or SEQ ID NO:258.

An embodiment of the invention is directed towards recombinant vectors which encode a patatin permutein protein. The vector can comprise operatively linked in the 5' to 3' orientation: a promoter that directs transcription of a structural nucleic acid sequence; a structural nucleic acid sequence encoding a protein selected from the group consisting of: (amino acids 247-386 of SEQ ID NO:2)-linker-(amino acids 24-246 of SEQ ID NO:2); and (amino acids 269-386 of SEQ ID NO:2)-linker-(amino acids 24-268 of SEQ ID NO:2); and a 3' transcription terminator. The linker can comprise Gly-Pro-Gly (SEQ ID NO:277) or Gly-Gly-Gly-Ser-Gly-Gly-Gly (SEQ ID NO:276). Alternatively, the encoded patatin permutein protein can lack a linker sequence. The structural nucleic acid sequence can preferably be SEQ ID NO:246 or SEQ ID NO:258, and preferably encodes SEQ ID NO:247 or SEQ ID NO:259.

An additional embodiment of the invention is directed towards recombinant host cells useful for the production of a patatin permutein protein. The recombinant host cell preferably produces a patatin permutein protein. More preferably, the recombinant host cell produces a patatin permutein protein in a concentration sufficient to inhibit growth or to kill an insect which ingests the recombinant host cell. The recombinant host cell can comprise a structural nucleic acid sequence encoding a protein selected from the group consisting of: (amino acids 247-386 of SEQ ID NO:2)-linker-(amino acids 24-246 of SEQ ID NO:2); and (amino acids 269-386 of SEQ ID NO:2)-linker-(amino acids 24-268 of SEQ ID NO:2). The linker can generally be any amino acid sequence, and preferably is Gly-Pro-Gly (SEQ ID NO:277) or Gly-Gly-Gly-Ser-Gly-Gly-Gly (SEQ ID NO:276). Alternatively, the encoded patatin permutein protein can lack a linker sequence. The structural nucleic acid sequence is preferably SEQ ID NO:246 or SEQ ID NO:258, and preferably encodes SEQ ID NO:247 or SEQ ID NO:259. The structural nucleic acid sequence can be operatively linked to a promoter sequence that directs transcription of the structural nucleic acid sequence, a 3' transcription terminator, and a 3'

polyadenylation signal sequence. The recombinant host cell can generally be any type of host cell, and preferably is a bacterial, fungal, or plant host cell. The bacterial cell is preferably an *Escherichia coli* bacterial cell. The fungal cell is preferably a *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, or *Pichia pastoris* fungal cell. The plant cell can be a monocot, dicot, or conifer plant cell. The plant cell is preferably an alfalfa, banana, canola, corn, cotton, cucumber, peanut, potato, rice, soybean, sunflower, sweet potato, tobacco, tomato, or wheat plant cell.

An additional embodiment of the invention is directed towards recombinant plants which are useful for the production of patatin permutein proteins. The recombinant plant preferably produces a patatin permutein protein. More preferably, the recombinant plant produces a patatin permutein protein in a concentration sufficient to inhibit growth or to kill an insect which ingests tissue from the recombinant plant. The recombinant plant can comprise a structural nucleic acid sequence encoding a protein selected from the group consisting of: (amino acids 247-386 of SEQ ID NO:2)-linker-(amino acids 24-246 of SEQ ID NO:2); and (amino acids 269-386 of SEQ ID NO:2)-linker-(amino acids 24-268 of SEQ ID NO:2). The linker can comprise Gly-Pro-Gly (SEQ ID NO:277) or Gly-Gly-Gly-Ser-Gly-Gly-Gly (SEQ ID NO:276). Alternatively, the encoded protein can lack a linker sequence. The structural nucleic acid sequence is preferably SEQ ID NO:246 or SEQ ID NO:258, and preferably encodes SEQ ID NO:247 or SEQ ID NO:259. The structural nucleic acid sequence can be operatively linked to a promoter sequence that directs transcription of the structural nucleic acid sequence, a 3' transcription terminator, and a 3' polyadenylation signal sequence. The recombinant plant can generally be any type of plant, and preferably is an alfalfa, banana, canola, corn, cotton, cucumber, peanut, potato, rice, soybean, sunflower, sweet potato, tobacco, tomato, or wheat plant.

Permutein proteins can be prepared by isolating the permutein protein from any one of the above described host cells or plants.

Immunotherapy for potato allergy

Immunotherapy for food allergy has been largely unsuccessful due to the lack of appropriate therapeutic reagents (Sampson, H.A., *J. Allergy Clin. Immunol.*, 90(2): 151-152, 1992). Immunotherapy has typically involved the administration (orally or by

subcutaneous injections) of increasing doses of crude protein extracts of the offending allergenic entities which usually contain variable mixes of many different proteins (Scheiner, O., *Wien Klin Wochenschr.*, 105(22): 653-658, 1993). While there are reports of highly successful clinical applications of immunotherapy for food allergens (Romano, P.C., *et al.*, *Allergol. Immunopathol. (Madr)*, 12(4): 275-281, 1984), those reports are rare and the clinical literature in general recommends avoidance far more strongly than therapy (Gay, G., *Allerg. Immunol. (Paris)*, 29(6): 169-170, 1997). One of the primary reasons for the failure of many clinical attempts to induce tolerance to allergens in general and food allergens in particular relates to anecdotal comments by numerous allergists, that patients don't tolerate the doses of allergen required to achieve tolerance. Animal studies examining the relationship of antigen dose and the induction of tolerance have demonstrated a strong positive correlation (Chen, Y., *et al.*, *Proc. Natl. Acad. Sci., U.S.A.*, 93: 388-391, 1996; Tokai, T., *et al.*, *Nat. Biotechnol.*, 15(8): 754-758, 1997). Due to the very real possibility of inducing an anaphylactic reaction in patients with native allergen, most clinical therapists are quite hesitant to use high doses therapeutically and are therefore compromising the likelihood of successful therapy.

In recent reports, recombinant technology has been used to reduce the allergenic potential of a major allergen without modifying the T cell epitopes, and allowing higher doses of protein to be used in therapy (Tokai, T., *et al.*, *Nat. Biotechnol.*, 15(8): 754-758, 1997). In addition, a lack of understanding about the appropriate route of administration, the uncertainty of mechanisms responsible for induction of allergy and the uncertainty of mechanisms by which immunotherapy suppresses or blocks the T cell-IgE-eosinophil/mast cell cycle have contributed to the large number of equivocal studies and clinical trials. Recent studies in animal models dealing with mechanisms, routes of administration, adjuvants and vaccine formulations have increased the likelihood that immunotherapy for allergies, including food allergies, will become a reproducibly successful clinical treatment when the appropriate therapeutic reagents are available (Sampson, H.A. and Burks, A.W., *Annu. Rev. Nutr.*, 16: 161-177, 1996; Kaminogawa, S., *Biosci. Biotechnol. Biochem.*, 60(11): 1749-1756, 1996; Chapman, M.D., *et al.*, *Allergy*, 52: 374-379, 1997; Barbeau, W.E., *Adv. Exp. Med. Biol.*, 415: 183-193, 1997; Cao, Y., *et al.*, *Immunology*, 90(1): 46-51, 1997; Garside, P. and Mowat, A.M., *Crit. Rev. Immunol.*,

17(2): 119-137, 1997; Rothe, M.J. and Grant-Kels, J.M., *J. Am. Acad. Dermatol.*, 35(1):
 1-13, 1996; Strobel, S., *Allergy*, 50(20): 18-25, 1995; Kruisbeek, A.M. and Amsen, D.,
Curr. Opin. Immunol., 8(2): 233-244, 1996; Herz, U., *et al.*, *Adv. Exp. Med. Biol.*, 409:
 25-32, 1996; Litwin, A., *et al.*, *J. Allergy Clin. Immunol.*, 100: 30-38, 1997;
 Vandewalker, M.L., *Mo. Med.*, 94(7): 311, 1997; Marshall, G.D., Jr. and Davis, F., *Nat.*
Biotechnol., 15(8): 718-719, 1997; Van Deusen, M.A., *et al.*, *Ann. Allergy Asthma*
Immunol., 78: 573-580, 1997; Jacobsen, L., *et al.*, *Allergy*, 52: 914-920, 1997, Scheiner,
 O. and Kraft, D., *Allergy* 50(5): 384-391, 1995).

Relative to immunotherapy, the critical aspects of the modified patatin genes
 described in this patent are that they can be used to synthesize purified, deallergenized-
 protein which can be used for patatin (potato) specific immunotherapy, with reduced
 potential for adverse and potentially fatal anaphylactic reactions in human or veterinary
 patients who have allergies to patatin or potatoes. Various strategies, including fixing or
 cross linking allergens, encapsulation of allergen for oral delivery, the use of small, T-cell
 epitope peptides and most recently, the use of engineered recombinant proteins, or
 modified gene vaccines are being tested in attempts to decrease the potential for
 anaphylactic reactions while inducing tolerance (Cao, Y., *et al.*, *Immunology*, 90(1): 46-
 51, 1997; Chapman, M.D., *et al.*, *Allergy*, 52: 374-379, 1997; Chapman, M.D., *et al.*, *Int.*
Arch. Allergy Immunol., 113(1-3): 102-104, 1997; Collins, S.P., *et al.*, *Clin. Exp. Allergy*,
 26(1): 36-42, 1996; Takai, T., *et al.*, *Mol. Immunol.*, 34(3): 255-261, 1997; Takai, T., *et*
al., *Nat. Biotechnol.*, 15(8) 754-758, 1997; Jirapongsananruek, O. and Leung, D.Y.M.,
Ann. Allergy Asthma Immunol., 79: 5-20, 1997; Litwin, A., *et al.*, *J. Allergy Clin.*
Immunol., 100: 30-38, 1997; Vandewalker, M.L., *Mo. Med.*, 94(7): 311, 1997; Raz, E., *et*
al., *Proc. Natl. Acad. Sci., U.S.A.*, 93: 5141-5145, 1996; Hoyne, G.F., *et al.*, *Clin.*
Immunol. Immunopathol., 80: S23-30, 1996; Hoyne, G.F., *et al.*, *Int. Immunol.*, 9(8):
 1165-1173, 1997; Vrtala, S., *et al.*, *J. Clin. Invest.*, 99(7): 1673-1681, 1997; Sato, Y., *et*
al., *Science*, 273: 352-354, 1996; Lee, D.J., *et al.*, *Int. Arch. Allergy Immunol.*, 113(1-3):
 227-230, 1997; Tsitoura, D.C., *et al.*, *J. Immunol.*, 157(5): 2160-2165, 1996; Hsu, C.H.,
et al., *Int. Immunol.*, 8(9):1405-1411, 1996; Hsu, C.H., *et al.*, *Nat. Med.*, 2(5): 540-544,
 1996).

1 The instant invention uses an engineered patatin protein, as expressed in any
2 living cell, with or without post-synthesis modifications, for immunotherapy by the
3 routes of cutaneous or subcutaneous exposure, injection, or by oral, gastro-intestinal,
4 respiratory or nasal application, either with, or without the use of specific carriers,
5 vehicles and adjuvants. The direct application of nucleic acid encoding recombinant
6 patatin as the *in vivo* (in the patient) expression template (gene) as RNA-, DNA- or gene-
7 vaccines is also the intended use of the engineered genetic materials defined here, coding
8 for patatin, but with modified IgE binding sites. It is also the intent of this patent to cover
9 the use of these modified genes described here including insertion into various DNA
10 vectors including adenovirus, retrovirus, pox virus and replicating or non-replicating
11 eukaryotic expression plasmids (Lee, D.J., *et al.*, *Int. Arch. Allergy Immunol.*, 113(1-3):
12 227-230, 1997) with various promoters and regulatory sequences, which can be inserted
13 into the patient's somatic cells (dendritic cells, epithelial cells, muscle fiber-cells,
14 fibroblasts, etc.) for the purpose of expressing the recombinant gene product to alter the
15 patient's immune response to the patatin proteins (Lee D.J., *et al.*, *Int. Arch. Allergy*
16 *Immunol.*, 113(1-3): 227-230, 1997). Potential routes of administration foreseen in this
17 application include previously described methods of encapsulation, emulsion, receptor or
18 membrane fusion mediated uptake and methods of direct permeabilization or insertion of
19 the DNA or corresponding RNA into the host cells.

20 The following examples are included to demonstrate preferred embodiments of
21 the invention. It should be appreciated by those of skill in the art that the techniques
22 disclosed in the examples which follow represent techniques discovered by the inventors
23 to function well in the practice of the invention, and thus can be considered to constitute
24 preferred modes for its practice. However, those of skill in the art should, in light of the
25 present disclosure, appreciate that many changes can be made in the specific
26 embodiments which are disclosed and still obtain a like or similar result without
27 departing from the spirit and scope of the invention.

EXAMPLES

Example 1: Identification of patatin as an allergen

Since patatin is commonly obtained from an allergenic source (potato), it was hypothesized that patatins in fact encode an important class of offending potato allergens (patatin was reported as allergenic by Seppala, U. et al., *J. Allergy Clin. Immunol.* 103: 165-171, 1999). Assessment of potential allergens preferably include appropriate *in vitro* testing for IgE binding, in this case with potato allergic sera (Fuchs, R.L. and Astwood, J.D., *Food Technology*, 50: 83-88, 1996; Astwood, J.D., et al., *Monographs in allergy Vol. 32: Highlights in food allergy*, pp. 105-120, 1996, Metcalfe, D.D., et al., *Critical Reviews in Food Science and Nutrition*, 36S: 165-186, 1996). It is the recommendation of a working group organized by the IFBC and the ILSI Allergy and Immunology Institute that proteins encoded by nucleic acid sequences from allergenic sources such as potato (a “less-commonly” allergenic source) should be examined for their ability to react with IgEs of potato-allergic patients using a minimum of five individual patient sera (Metcalfe, D.D., et al., *Critical Reviews in Food Science and Nutrition*, 36S: 165-186, 1996). Patatin-17 protein was tested for IgE binding using standard *in vitro* testing with serum taken from patients with bona fide well defined clinically displayed potato allergy as described below.

Clinical Characterization of Potato Allergic Subjects (Serum donors)

Patients who suffer from potato allergy were identified at Johns Hopkins Clinic (Baltimore, MD) and were evaluated for potato allergy using clinical criteria outlined in Table 2.

Serum was obtained from patients with convincing clinical history of potato allergy. The convincing history was defined as being one or more of the following: a) positive potato allergic as evaluated by double-blind placebo-control food challenge b) anaphylaxis and/or hospitalization due to the consumption of potatoes or c) dramatic skin test results.

Table 2: Clinical patient data

Patient	Clinical History	Flare/Wheal (Skin prick test)	DBPCFC (potato)
HS01	Most recent hospitalization: 10/19/93 AD, A, AR, FH, MFS, IgE =1397 KIAUa/L	7/19, 4/14, 7/17	Not performed
HS02	Most recent hospitalization: 6/94 AD, FH, Latex (+) RAST, MFS, IgE=7544K/L	20/26	Not performed
HS03	Most recent hospitalization: 7/27/95 AD, A, FH, MFS, IgE = N/A	5/13	Yes
HS05	Most recent hospitalization 5/30/95 AD, A, FH, MFS, IgE = 12341 ng/ml	4/9	Yes
HS06	Most recent hospitalization 6/13/95 AD, A, FH, MFS IgE = N/A	5/20, 4/13, 5/12	Yes
HS07	Not potato allergic, allergic to egg, milk, peanuts, seafood. AD, A, AR, FH, MFS	High IgE control serum, not allergic to potato.	
HS08	Non-atopic (normal)	Low IgE control serum	

AD= Atopic dermatitis; FH= Food hypersensitivity; AR = Allergic rhinitis; A= Asthma; MFS= Multiple food sensitivity; N/A = not available.

Example 2: Western blotting of patatin proteins

Western blotting experiments were performed using patatin protein purified to near homogeneity from corn plants genetically engineered to produce patatin, patatin producing crude genetically engineered corn leaf extracts, crude potato tuber extracts, and non-transgenic corn leaf samples.

Protein samples were electrophoresed by SDS-PAGE (Laemmli, U.K., *Nature* 227: 680-685, 1970) and were electroblotted onto nitrocellulose. Protein blots were processed by standard Western blotting (immunoblotting) techniques and were incubated in potato allergic serum diluted 1:5 in PBS buffer for 1 hour. After washing the blots 3 times with PBS, the blots were incubated in biotinylated anti-IgE (Johns Hopkins Hospital, Baltimore, MD) for 1 hour, followed by a 30 minute incubation in HRP-linked avidin(Promega, New York, NY). IgE-reactive protein bands were visualized by DAB staining (3,3 diaminobenzidine). The blots were dried and photographed. Individual

blots are labeled according to patient serum used. As a control, one blot was incubated in anti-IgE only.

Patatins were shown to be an allergen of potato by examining the reactivity of purified patatin to sera obtained from patients allergic to potato. Sera from five potato allergic subjects were tested by Western blotting techniques. All five sera reacted with purified patatin protein.

Patatin isozymes (SEQ ID NOS:278-282, Figure 1) were tested for IgE binding by Western blotting. Isozymes of patatin were cloned into a yeast expression system and purified prior to analysis. The isozymes were subjected to IgE western blotting as described above with the exception that all five patient sera were pooled. The resulting Western blot of the yeast-expressed isozymes showed that all five isozymes bound IgE in a manner similar to patatin 17, and that all isozymes of patatin tested are also allergens.

Example 3: Western blotting of patatin proteins

Eighty-nine 10-mer peptides were synthesized using the Genosys SPOTs system, each consecutive 10-mer overlapping by 6 amino acids based on the amino acid sequence of patatin 17 (SEQ ID NO:2). The peptides were evaluated for IgE binding with five different potato allergic patient sera using the same incubation procedures as described above. The results are summarized graphically in Figure 2, showing major and minor allergenic epitopes. Interestingly, many of the immunogenic epitopes contain tyrosine. The peptide numbers, sequences, and immunoreactivity is detailed in Table 3.

Table 3: Peptide scan of patatin 17

Peptide # (SEQ ID NO)	Peptide Sequence	HS01	HS02	HS03	HS05	HS06	Cumulative Total
1 (16)	QLGEMVTVLS	0.47	0.33	0.02	0.05	0.06	0.93
2 (17)	MVTVLSIDGG	0.53	0.33	0.02	0.07	0.05	1
3 (18)	LSIDGGGIRG	0.52	0.38	0.07	0.08	0.09	1.14
4 (19)	GGIRGIIPA	0.53	0.19	0.06	0.19	0.23	1.2
5 (20)	RGIIPATILE	0.46	0.28	0.04	0.09	0.05	0.92
6 (21)	PATILEFLEG	0.49	0.31	0.05	0.09	0.07	1.01
7 (22)	LEFLEGQLQE	0.36	0.24	0.04	0.1	0.06	0.8
8 (23)	EGQLQEMDNN	0.29	0.19	0.02	0.09	0.05	0.64
9 (24)	QEMDNNADAR	0.22	0.13	0.01	0.05	0.04	0.45
10 (25)	NNADARLADY	0.21	0.17	0.03	0.05	0.07	0.53

11 (26)	ARLADYFDV1	0.54	0.31	0.16	0.15	0.25	1.41
12 (27)	DYFDVIGGTS	0.61	0.34	0.46	0.06	0.15	1.62
13 (28)	VIGGTSTGGL	0.63	0.72	0.05	0.15	0.09	1.64
14 (29)	TSTGGLLTAM	0.3	0.17	0.03	0.06	0.09	0.65
15 (30)	GLLTAMISTP	0.63	0.41	0.05	0.24	0.12	1.45
16 (31)	AMISTPNENN	0.34	0.18	0.02	0.07	0.02	0.63
17 (32)	TPNENNRFFA	0.46	0.22	0.03	0.19	0.07	0.97
18 (33)	NNRPFAAAKE	0.37	0.21	0.05	0.07	0.06	0.76
19 (34)	FAAAKEIVPF	0.52	0.29	0.08	0.11	0.08	1.08
20 (35)	KEIVPFYFEH	0.29	0.14	0.28	0.29	0.23	1.23
21 (36)	PFYFEHGPQI	0.65	0.06	1.08	0.51	0.17	2.47
22 (37)	EHGPQIFNPS	0.34	0.15	0.03	0.05	0.06	0.63
23 (38)	QIFNPSGQIL	0.33	0.29	0.02	0.07	0.07	0.78
24 (39)	PSGQILGPKY	0	0	0.02	0	0.05	0.07
25 (40)	ILGPKYDGKY	0	0	0.07	0	0.02	0.09
26 (41)	KYDGKYLMOV	0.02	0	0.11	0.01	0.04	0.18
27 (42)	KYLMQVLQEK	0.12	0.04	1.08	0.07	0.79	2.1
28 (43)	QVLQEKLGTE	0.46	0.16	0.01	0.07	0.02	0.72
29 (44)	EKLGETRVHQ	0.5	0.12	0.01	0.07	0.04	0.74
30 (45)	ETRVHQALTE	0.42	0.16	0.03	0.05	0.03	0.69
31 (46)	HQALTEVVIS	0.43	0.21	0.04	0.1	0.05	0.83
32 (47)	TEVVISSFDI	0.44	0.25	0.05	0.08	0.04	0.86
33 (48)	ISSFDIKTNK	0.1	0.02	0.04	0.06	0.13	0.35
34 (49)	DIKTNKPVIF	0.57	0.22	0.04	0.18	0.28	1.29
35 (50)	NKPVIFTKSN	0	0.01	0.02	0.07	0.24	0.34
36 (51)	IFTKSNLANS	0	0	0.03	0.06	0.17	0.26
37 (52)	SNLANSPELD	0.43	0.96	0.01	0.09	0.02	1.51
38 (53)	NSPELDKAMY	0.18	0.12	0.01	0.05	0.05	0.41
39 (54)	LDKAMYDISY	0.54	0.26	0.19	0.15	0.23	1.37
40 (55)	MYDISYSTAA	0.92	0.08	0.52	0.04	0.22	1.78
41 (56)	SYSTAAPTY	1.15	0.25	1.04	0.33	0.55	3.32
42 (57)	AAPTYFPPH	1.02	0.52	1.12	0.81	0.86	4.33
43 (58)	TYFPFHYFVT	0.02	0.01	0.54	0.03	0.24	0.84
44 (59)	PHYFVTNTSN	0.03	0.01	1.17	0.13	0.44	1.78
45 (60)	VTNTSNGDEY	0.23	0.15	0.04	0.03	0.03	0.48
46 (61)	SNGDEYFNL	0.33	0.25	0.08	0.1	0.11	0.87
47 (62)	EYFNLVDGA	0.34	0.25	0.07	0.1	0.2	0.96
48 (63)	NLVDGAVATV	0.3	0.18	0.02	0.06	0.05	0.61
49 (64)	GAVATVADPA	0.45	0.54	0.01	0.07	0.02	1.09
50 (65)	TVADPALLSI	0.48	0.29	0.01	0.07	0.03	0.88
51 (66)	PALLSISVAT	0.65	0.33	0.02	0.1	0.01	1.11
52 (67)	SISVATRLAQ	0.61	0.23	0.14	0.53	0.53	2.04
53 (68)	ATRLAQKDPA	0.87	0.34	0.05	0.29	0.22	1.77
54 (69)	AQKDPAFASI	0.86	0.32	0.04	0.12	0.03	1.37
55 (70)	PAFASIRSLN	0.81	0.15	0.05	0.51	0.59	2.11
56 (71)	SIRSLNYKKM	0.07	0.01	0.17	0.07	0.11	0.43
57 (72)	LNYKKMLLS	0.05	0.01	0.35	0.08	0.39	0.88
58 (73)	KMLLSLGTG	1.15	0.15	0.04	0.38	0.71	2.43
59 (74)	LSLGTGTSE	0.34	0.23	0.02	0.04	0.03	0.66
60 (75)	TGTTSEFDKT	0.92	0.39	0.6	0.1	0.09	2.1
61 (76)	SEFDKTYTAK	1.33	1.35	1.41	0.12	0.28	4.49
62 (77)	KTYTAKEAAT	1.36	0.94	1.11	0.76	0.4	4.57

63 (78)	AKEAATWTAV	0.45	0.15	0.01	0.2	0.03	0.88
64 (79)	ATWTAVHWMML	0.1	0.02	0.01	0.08	0.03	0.27
65 (80)	AVHWMMLVIQK	0.69	0.05	0.03	0.43	0.62	1.82
66 (81)	MLVIQKMTDA	0.32	0.15	0.02	0.15	0.03	0.67
67 (82)	QKMTDYYLST	0.26	0.125	0.03	0.21	0.03	0.675
68 (83)	DAASSYMTDY	0.2	0.14	0.08	0.08	0.1	0.6
69 (84)	SYMTDYYLST	0.5	0.03	0.32	0.05	0.11	1.02
70 (85)	DYYLSTAFQA	0.44	0	0.22	0.43	0.03	0.52
71 (86)	STAFQALDSK	0.1	0.3	0.08	0.05	0.03	0.88
72 (87)	QALDSKNNYL	0.44	0.46	0.28	0.26	0.03	1.87
73 (88)	SKNNYLRLVQE	0.44	0.05	1.31	0.07	0.21	2.08
74 (89)	YLRVQENALT	1.38	0.03	1.31	0.11	0.2	3.03
75 (90)	QENALTGTTT	0.47	0.25	0	0.06	0	0.78
76 (91)	LTGTTTTEMDD	0.41	0.24	0	0.06	0	0.71
77 (92)	TTEMDDASEA	0.38	0.3	0	0.05	0	0.73
78 (93)	DDASEANMEL	0.44	0.24	0	0.06	0	0.74
79 (94)	EANMELLVQV	0.42	0.27	0	0.04	0	0.73
80 (95)	ELLVQVGENL	0.4	0.25	0	0.05	0	0.7
81 (96)	QVGENLLKKP	0.44	0.14	0	0.07	0	0.65
82 (97)	NLLKKPVSED	0.47	0.2	0	0.03	0	0.7
83 (98)	KPVSEDNPET	0.27	0.21	0	0.03	0	0.51
84 (99)	EDNPETYEEA	0.13	0.11	0	0.01	0	0.25
85 (100)	ETYEEALKRF	1.26	1.2	1.36	0.53	0.71	5.06
86 (101)	EALKRFAKLL	1.38	0.04	0	1.06	1.12	3.6
87 (102)	RFAKLLSDRK	0.98	0.05	0	0.84	0.94	2.81
88 (103)	LLSDRKKLRA	0.2	0.01	0	0.37	0.51	1.09
89 (104)	RKKLRANKAS	0.28	0	0	0.31	0.64	1.23
	Patient Cumulative Totals	41.84	20.565	18.1	14.17	16.55	

Example 4: Identification of result effective substitutions

For each major and several minor allergenic epitopes of patatin, result effective substitutions were identified by synthesizing peptides that were altered by individually substituting an alanine residue at each non-alanine position in the epitope. Similarly, the reported nucleic acid sequence encoding corn patatin (U.S. Patent No. 5,882,668; clone 5c9) was evaluated for IgE binding by producing peptides at corresponding positions to the potato patatin protein.

For example, Epitope 41 was analyzed by alanine scanning and rational substitution as follows.

Epitope 41	SEFDKTYTAK (SEQ ID NO:76)
Alanine scan	AEFDKTYTAK (SEQ ID NO:165)

1 SAFDKITYTAK (SEQ ID NO:166)
2 SEADKITYTAK (SEQ ID NO:167)
3 SEFAKITYTAK (SEQ ID NO:168)
4 SEFDATYTAK (SEQ ID NO:169)
5 SEFDKAYTAK (SEQ ID NO:170)
6 SEFDKTATAK (SEQ ID NO:171)
7 SEFDKTYAAK (SEQ ID NO:172)
8 SEFDKITYTAA (SEQ ID NO:173)
9 Rational substitution AFFDKITYTAK (SEQ ID NO:283)
10 SEFDKTFTAK (SEQ ID NO:176)
11 Corn homolog CIFDSTYTAK (SEQ ID NO:284)

12
13 Selected epitopes were analyzed by alanine scanning and rational substitution.
14 Immunoassay with potato-allergic serum was used as described above. Table 4
15 summarizes the results of these experiments to identify result effective substitutions for
16 patatin. Blank spaces in the table indicate that binding of the peptide to patient IgE was
17 not detectable.

18 Table 4: Scanning of patatin for result effective substitutions

Sequence	SEQ ID NO	Binding of modified peptides by patient IgE as measured by OD			
		HS03	HS06	HS01	HS02
DYFDVIGGTS	105		0.12	0.16	0.36
DYFDVIAGTS	106		0.14	0.17	0.4
VIGGTSTGGL	107				0.04
VIAGTSTGAL	108				
AFYFEHGPQI	109		0.96	0.5	0.78
PAYFEHGPQI	110		0.75	0.41	0.69
PFAFEHGPQI	111				
PFYAEHGPQI	112		0.7	0.43	0.79
PFYFAHGPQI	113	0.93	1.07	0.59	1.44
PFYFEAGPQI	114	0.08	0.93	0.65	1.34
PFYFEHAPQI	115		0.75	0.54	1.11
PFYFEHGAQI	116		0.63	0.29	0.6
PFYFEHGPPI	117		0.63	0.25	0.56
PFYFEHGPQA	118		0.27	0.16	0.33
TFYLENGPKI	119	0.05	0.48	0.68	1.07
PFFFEHGPQI	120				
AYLMQVLQEK	121		0.26	0.11	0.53
KALMQVLQEK	122				

KYAMQVLQEK	123	0.11	0.43	0.1	1.25
KYLAQVLQEK	124	0.22	0.48	0.11	1.34
KYLMAVLQEK	125	0.22	0.83	0.16	1.33
KYLMQALQEK	126	0.11	0.6	0.15	0.95
KYLMQVQAEK	127		0.53	0.15	0.81
KYLMQVLAEK	128	0.06	0.69	0.11	1.34
KYLMQVLQAK	129	0.74	0.79	0.05	0.58
KYLMQVLQEA	130		0.28	0.27	0.37
VFLHDKIKSL	131	0.06	0.26		0.41
AYSTAAAPTY	132		0.1	0.12	0.12
SASTAAAPTY	133				
SYATAAAPTY	134		0.16	0.13	0.37
SYSAAAPTY	135		0.13	0.12	0.32
SYSTAAAPTY	136		0.15	0.13	0.34
SYSTAAAPAY	137		0.15	0.14	0.29
SYSTAAAPTA	138		0.55	0.54	1.13
CISTAAAPTY	139	0.4			
SYSTAAAPAF	140	0.39	1.02	0.65	1.42
AFAAAAAPTY	141				0.07
SYSTAAAPTF	142	0.15	0.97	0.48	1.09
STSAAPTYFP	143		0.21	0.23	0.39
STSAAPTFPP	144				0.23
STSAAPTAFP	145				0.08
STAAAPTFPP	146			0.12	0.28
AAAAATYFPPH	147		0.13	0.1	0.05
AAAPAYFPPH	148			0.07	0.04
AAAPTAFFPPH	149				
AAAPTYAPPH	150		0.23	0.14	0.21
AAAPTYFAPH	151		0.45	0.18	0.44
AAAPTYFPAH	152		0.15	0.07	0.18
AAAPTYFPPA	153		0.1	0.06	0.31
SAAPTYFPAH	154		0.77	0.73	0.96
AAAPAFFPPH	155				
AAAPPFFPPH	156				
AAAPTFPPPH	157				
SISVATRLAQ	158			0.26	0.26
AMSMKTKEVH	159				
PAFASIRSLN	160				
PNFNAGSPTE	161				
KMLLLSLGTG	162				
NYLIISVGTG	163	0.49	1.08	0.64	1.48
KMLLLSLGAG	164		0.13		
AEFDKTYTAK	165	0.09	0.22		1.34
SAFDKTYTAK	166	0.66	0.71	0.06	1.42
SEADKTYTAK	167				0.99
SEFAKTYTAK	168	0.5	0.57		0.91
SEFDATYTAK	169				0.17
SEFDKAYTAK	170	0.1	0.24		1.38
SEFDKTATAK	171				0.81
SEFDKTYAAK	172	0.2	0.35		1.39
SEFDKTYTAA	173			0.1	1.18
KQAEKYTAEO	174			0.08	0.24

SEFDAAFAAA	175				
SEFDKTTAK	176	0.09	0.16	0.07	1.45
AEKYTAEQCA	177				
ATYTAKEAAT	178		0.24		0.18
KAYTAKEAAT	179		0.28		0.33
KTATAKEAAT	180				
KTYAAKEAAT	181	0.1	0.32		0.73
KTYTAAEAAT	182				0.35
KTYTAKAAAT	183	0.4	0.59		0.82
KTYTAKEAAA	184				0.36
EKYTAEQCAK	185				
AAFAAAEAAT	186				
KTFTAKEAAT	187				
QALHCEKKYL	188				
QALDSKAAYL	189				
QALDSKNNFL	190				
QALHCENNFL	191				
CEKKYLRIQD	192	1.01	0.16		
SKNNFLRVQE	193				
SENNYLRVQE	194	0.31	0.96	0.42	1
ALRVQENALT	195				
YARVQENALT	196	1.06	1.02	0.05	0.54
YLAVQENALT	197	0.37	1.04	0.11	1.06
YLRAQENALT	198	1.1	1	0.06	1.26
YLRVAENALT	199	1.03	0.92	0.08	1.26
YLRVQANALT	200	1.05	0.92	0.06	1.24
YLRVQEAALT	201	0.93	0.92	0.07	1.11
YLRVQENALT	202	0.94	0.93	0.04	1.24
YLRVQENALA	203	1.05	0.96	0.43	1.16
YLRIQDDTLT	204	1.07	0.85	0.39	1.12
YLTVAALALT	205	1.05	0.86	0.28	1.33
FLRVQENALT	206				
NNYLRVQENA	207	0.23	0.88	0.5	1.17
KKYLRIQDDT	208		0.26	0.09	0.37
NNFLRVQENA	209				
NAYLRVQENA	210	0.17	1.02	0.53	1.06
ATYEEAKLRF	211	0.26	1.03		0.65
EAYEEALKRF	212	0.06	0.43		0.33
ETAEEALKRF	213		1.04		
ETYAEALKRF	214	0.62	1.02		1.15
ETYEALKRF	215	1.06	0.38		0.89
ETYEAAKRF	216	0.08	0.1		0.9
ETYEALARF	217				0.11
ETYEALKAF	218				0.1
ETYEALKRA	219				0.1
GTNAQSLADF	220				
ETYEALAAAF	221	0.07	0.78	0.33	0.77
ETFEALKRF	222				
YEEALKTFK	223	1.08	0.85	0.14	1.46
FEEALKRFK	224	0.46	0.72		0.67
AALKRFKALL	225	0.15	0.17		
EAAKRFKALL	226	0.08	0.33		0.05

EALARFAKLL	227		0.09		
EALKAFKLL	228				
EALKRAAKLL	229	0.08	0.07		
EALKRFAALL	230				
EALKRFAKAL	231	0.06	0.09		0.1
EALKRFAKLA	232	0.06			0.1
QSLADFAKQL	233				
AALAAFAKLL	234				
LADFAKQLSD	235				
DFAKQLSDER	236				0.17
AFAALLSDRK	237				

Result effective substitutions were identified by a reduction in IgE binding ability with respect to the non-substituted peptide sequence. Table 5 shows the identified result effective substitutions. Blank spaces in the table indicate that binding of the peptide to patient IgE was not detectable. Many substitutions of alanine or phenylalanine for the original tyrosine resulted in reduced or eliminated antibody binding.

Table 5: Result effective substitutions of patatin

Location (SEQ ID NO)	Peptide (SEQ ID NO)	HS03	HS06	HS01	HS02
Minor Epitope 21	PFYFEHGPQI (36) ::A::: (111) ::F::: (r) (120) :::A (118)	1.08	0.17 0.27	0.65 0.16	0.06 0.33
Minor Epitope 27	KYLMQVLQEK (42) :A::: (122) :::A (130) VFLHDKIKSL (c) (131)	1.08 0.06	0.79 0.28 0.26	0.12 0.27	0.04 0.37 0.41
Major Epitope 41	SYSTAAPTY (56) A::: (132) :A::: (133) AFAA::: (r) (141) CI::S::: (c) (139)	1.04 0.04	0.55 0.1	1.15 0.12	0.25 0.12 0.007
Overlap Epitope 41/42	STAAPTYFP (238) ::S:::A:: (r) (145)				0.08
Major Epitope 42 (57)	AAAPTYFPPH (57) :::A::: (148) :::A::: (149) :::AF::: (r) (155) :::PF::: (r) (156) :::F::: (r) (157)	1.12	0.86	1.02 0.07	0.52 0.04
Major Epitope 61	SEFDKTYTAK (76) :::A::: (169) KQAE:YTAEQ (c) (174) :::AAFA:A (r) (175)	0.12	0.28	1.33 0.08	1.35 0.17 0.24
Major	KTYTAKEAAT (77)	1.11	0.04	1.36	0.94

Epitope 62	A::: (178) ::A::: (180) :::A::: (182) AAFA:A::: (r) (186) ::F::: (r) (187) EK::EQC:K (c) (185)		0.24		0.18 0.35
Minor Epitope 72	QALDSKNNYL (87) ::HCEKK:: (c) (188) :::AA:: (r) (189) :::F:: (r) (190) :::E::F:: (r) (240)	0.28	0.43	0.44	0.46
Minor epitope 73	SKNNYLRVQE (88) :::F::: (r) (193)	1.31	0.21	0.44	0.05
Minor epitope 74	YLRVQENALT (87) A::: (195) F::: (r) (206)	1.31	0.2	1.38	0.03
Overlap epitope 73/74	NNYLRVQENA (207) ::F::: (r) (209)	0.23	0.88	0.5	1.17
Major epitope 85	ETYEALKRF (100) :::A:: (217) :::A:: (218) :::A:: (219) ::F::: (r) (222) G:NAQS:AD: (c) (220)	1.36	0.71	1.26	1.2 0.11 0.1 0.1
Major Epitope 86	EALKRFKLL (101) ::A::: (227) :::A::: (228) :::A::: (229) :::A:: (230) :::A:: (231) :::A:: (232) SD:AD::Q: (c) (241) A::AA::: (r) (234)	0 0.08 0.06 0.06	1.12 0.09 0.07 0.09	1.38	0.04
Epitope overlap 86/87	LKRFAKLLSD (239) (NO BINDING)				
Major Epitope 87	RFAKLLSDRK (102) D:::Q:::ER (c) (236) A::A::: (r) (237)	0	0.94	0.98	0.05 0.17

(r) = rational; (c) = corn.

Example 5: Site directed mutagenesis

To introduce site specific mutations, the cloned DNA sequence of patatin (SEQ ID NO:1 encoding patatin protein SEQ ID NO:2; pMON 26820) was subjected to PCR with primers SEQ ID NO:3 and SEQ ID NO:4 to incorporate part of the α -factor signal sequence (*Pichia* expression manual, Invitrogen, Carlsbad, CA), and EcoRI and XhoI restriction sites to facilitate cloning into the *Pichia pastoris* yeast secretion vector pPIC9 (GenBank accession number Z46233; Invitrogen, Carlsbad, CA). Typical PCR

1 conditions are 25 cycles 94°C denaturation for 1 minute, 45°C annealing for one minute
2 and 72°C extension for 2 minutes; plus one cycle 72°C extension for 10 minutes. A 50
3 µL reaction contains 30 pmol of each primer and 1 µg of template DNA; and 1 X PCR
4 buffer with MgCl₂, 200 µM dGTP, 200 µM dATP, 200 µM dTTP, 200 µM dCTP, 2.5
5 units of *Pwo* DNA polymerase. PCR reactions are performed in RoboCycler Gradient 96
6 Temperature Cycler (Stratagene, La Jolla, CA).

7 The amplified fragment SEQ ID NO:5 was digested with restriction enzymes
8 XhoI and EcoRI and cloned into the pBluescript vector (Stratagene, La Jolla, CA),
9 digested with the same two restriction enzymes. The resulting plasmid (pMON 26869)
10 was used for oligonucleotide-directed mutagenesis using the Bio-Rad mutagenesis kit
11 based on the method of Kunkel (*Proc. Natl. Acad. Sci. U.S.A.*, 82: 477-492, 1985).
12 Briefly, single-stranded pMON26869 was used as template for mutagenesis and was
13 prepared by superinfection of plasmid containing cells with M13K07 (Gorman, *et al.*,
14 *DNA Prot. Eng. Techniques*, 2: 3-10, 1990). The mutagenic oligonucleotides are SEQ ID
15 NOS:8-15 (reverse complement). DNA purified from transformed DH5α *E. coli* colonies
16 was used for sequence determination. Sequencing was performed using the ABI PRISM
17 sequencing kit (Perkin Elmer Biosystems, Foster City, CA). The resulting plasmid
18 containing the mutation in the patatin gene was digested with restriction enzymes XhoI
19 and EcoRI.

20 The patatin nucleic acid fragment was then ligated into the pPIC9 vector
21 (Invitrogen, Carlsbad, CA), digested with the same two restriction enzymes to afford
22 plasmid pMON37401. *Pichia pastoris* KM71 cells were electroporated with
23 pMON37401 containing the appropriate mutation. The resulting transformed cells were
24 used to produce protein in *Pichia pastoris* using the procedure supplied by the
25 manufacturer (Invitrogen, Carlsbad, CA). The encoded protein contains an alpha factor
26 signal cleavage site. Plasmid pMON37401 encodes SEQ ID NO:6 which is cleaved to
27 afford SEQ ID NO:7, having four amino acids added at the N-terminus of amino acids
28 24-386 of SEQ ID NO:2. Position four of SEQ ID NO:7 therefore corresponds to
29 position 23 of SEQ ID NO:2.

30 The concentration of patatin in the culture was determined using a patatin ELISA
31 assay and the enzyme activity was measured using the method of Hofgen and Willmitzer

(*Plant Science*, 66: 221-230, 1990). The variants containing multiple mutations were further purified using Mono Q and hydrophobic interaction chromatography (HIC). Each culture was purified by first sizing on Amicon YM10 membranes (Millipore, Bedford, MA) to a >10 kDa fraction, followed by chromatography on the Mono Q HR 10/10 column (Pharmacia, Piscataway, NJ). For chromatography on the Mono Q column, the samples were loaded on the column in 25 mM Tris pH 7.5 and eluted with a gradient of 1.0 M KCl in 25 mM Tris pH 7.5. Fractions containing patatin protein were determined using SDS-PAGE. For chromatography on the HIC column, the appropriate fractions were pooled and dialyzed into 1 M ammonium sulfate in 25 mM Tris pH 7.5. The dialyzed sample was then loaded on 16/10 phenyl Sepharose column (Pharmacia, Piscataway, NJ) and eluted with a gradient of 25 mM Tris pH 7.5.

The protein concentration was determined using the Bradford method, using BSA as a standard. SDS-PAGE analysis showed that these proteins were essentially pure. The esterase activity of the newly formed variants are shown in Table 6. The activity was determined using p-nitrophenyl caprate substrate as described by Hofgen and Willmitzer (*Plant Science*, 66: 221-230, 1990).

Table 6: Esterase activity of patatin mutants

Variant	Activity (mOD.min ⁻¹ .μg ⁻¹)
Wild type	93.2
Y106F	51.1
Y129F	74.7
Y185F	85.6
Y193F	82.2
Y185F/Y193F	99.4
Y270F	163.4
Y316F	94.88
Y362F	130.7
Y106F/Y129F/Y185F/Y193F/Y270F/Y316F/Y362F	57.1
Y185F/Y193F/Y270F/Y316F/Y362F	161.5

Patatin proteins having a phenylalanine substitution at each of the amino acid positions 106, 129, 185, 193, 270, 316 and 362 (numbers correspond to positions in SEQ

1 ID NO:2) of expressed SEQ ID NO:7 exhibit full enzyme activity. Proteins having
2 multiple substitutions also displayed full enzyme activity.

3 In addition to nucleotide sequences encoding conservative amino acid changes
4 within the fundamental polypeptide sequence, biologically functional equivalent
5 nucleotide sequences include nucleotide sequences containing other base substitutions,
6 additions, or deletions. These include nucleic acids containing the same inherent genetic
7 information as that contained in the cDNA which encode peptides, polypeptides, or
8 proteins conferring pathogen resistance the same as or similar to that of pathogen upon
9 host cells and plants. Such nucleotide sequences can be referred to as “genetically
10 equivalent modified forms” of the cDNA, and can be identified by the methods
11 described herein.

12 Mutations made in the cDNA, plasmid DNA, genomic DNA, synthetic DNA, or
13 other nucleic acid encoding the deallergenized gene preferably preserve the reading
14 frame of the coding sequence. Furthermore, these mutations preferably do not create
15 complementary regions that could hybridize to produce secondary mRNA structures,
16 such as loops or hairpins, that would adversely affect mRNA translation.

17 Although mutation sites can be predetermined, it is not necessary that the nature
18 of the mutations *per se* be predetermined. For example, in order to select for optimum
19 characteristics of mutants at a given site, random mutagenesis can be conducted at the
20 target codon.

21 Alternatively, mutations can be introduced at particular loci by synthesizing
22 oligonucleotides containing a mutant sequence, flanked by restriction sites enabling
23 ligation to fragments of the native cDNA sequence. Following ligation, the resulting
24 reconstructed nucleotide sequence encodes a derivative form having the desired amino
25 acid insertion, substitution, or deletion.

26 Example 6: Construction of permutein sequences

27 Nucleic acid sequences encoding permutein proteins having rearranged N-
28 terminus/C-terminus protein sequences can be made by following the general method
29 described by Mullins et al. (*J. Am. Chem. Soc.* 116: 5529-5533, 1994). The steps are
30 shown in Figure 3. The Figure and the following Examples involve the design and use of

1 a linker region separating the original C-terminus and N-terminus, but the use of a linker
2 is not a critical or required element of permutein design.

3 Two sets of oligonucleotide primers are used in the construction of a nucleic acid
4 sequence encoding a permutein protein. In the first step, oligonucleotide primers “new
5 N-termini” and “linker start” are used in a PCR reaction to create amplified nucleic acid
6 molecule “new N-termini fragment” that contains the nucleic acid sequence encoding the
7 new N-terminal portion of the permutein protein, followed by the polypeptide linker that
8 connects the C-terminal and N-terminal ends of the original protein. In the second step,
9 oligonucleotide primers “new C-termini” and “linker end” are used in a PCR reaction to
10 create amplified nucleic acid molecule “new C-termini fragment” that contains the
11 nucleic acid sequence encoding the same linker as used above, followed by the new C-
12 termini portion of the permutein protein. The “new N-termini” and “new C-termini”
13 oligonucleotide primers are designed to include appropriate restriction enzyme
14 recognition sites which assist in the cloning of the nucleic acid sequence encoding the
15 permutein protein into plasmids.

16 Any suitable PCR conditions and polymerase can be used. It is desirable to use a
17 thermostable DNA polymerase with high fidelity to reduce or eliminate the introduction
18 of sequence errors. Typical PCR conditions are 25 cycles 94°C denaturation for 1
19 minute, 45°C annealing for one minute and 72°C extension for 2 minutes; plus one cycle
20 72°C extension for 10 minutes. A 50 µL reaction contains 30 pmol of each primer and 1
21 µg of template DNA; and 1 X PCR buffer with MgCl₂, 200 µM dGTP, 200 µM dATP,
22 200 µM dTTP, 200 µM dCTP, 2.5 units of *Pwo* DNA polymerase. PCR reactions are
23 performed in RoboCycler Gradient 96 Temperature Cycler (Stratagene, La Jolla, CA).

24 The amplified “new N-termini fragment” and “new C-termini fragment” are
25 annealed to form a template in a third PCR reaction to amplify the full-length nucleic
26 acid sequence encoding the permutein protein. The DNA fragments “new N-termini
27 fragment” and “new C-termini fragment” are resolved on a 1% TAE gel, stained with
28 ethidium bromide, and isolated using the QIAquick Gel Extraction Kit (Qiagen, Valencia,
29 CA). These fragments are combined in equimolar quantities with oligonucleotide
30 primers “new N-termini” and “new C-termini” in the third PCR reaction. The conditions

1 for the PCR are the same as used previously. PCR reaction products can be purified
2 using the QIAquick PCR purification kit (Qiagen, Valencia, CA).

3 Alternatively, a linker sequence can be designed containing a restriction site,
4 allowing direct ligation of the two amplified PCR products.

5 Example 7: Construction of plasmid pMON 37402

6 The patatin protein contains a trypsin protease sensitive site at the arginine amino
7 acid at position 246, as determined by electrophoresis of a trypsin digest reaction. In
8 order to determine if the exposed protease site is an antigenic epitope, a permutein was
9 constructed using positions 246-247 as a breakpoint.

10 The nucleic acid sequence encoding the permutein protein in plasmid pMON
11 37402 was created using the method illustrated in Figure 3 and described in Example 6.
12 Nucleic acid molecule "new N-termini fragment" was created and amplified from the
13 sequence encoding patatin in plasmid pMON26820 using oligonucleotide primers 27
14 (SEQ ID NO:242) and 48 (SEQ ID NO:243). Nucleic acid molecule "new C-termini
15 fragment" was created and amplified from the sequence encoding patatin in plasmid
16 pMON26820 using oligonucleotide primers 47 (SEQ ID NO:244) and 36 (SEQ ID
17 NO:245). The full-length nucleic acid molecule encoding the permutein protein was
18 created and amplified from annealed fragments "new N-termini fragment" and "new C-
19 termini fragment" using oligonucleotide primers 27 (SEQ ID NO:242) and 36 (SEQ ID
20 NO:245).

21 The resulting amplified nucleic acid molecule was digested with restriction
22 endonucleases XhoI and EcoRI, and purified using the QIAquick PCR purification kit
23 (Qiagen, Valencia, CA). Plasmid pMON 26869 (derivative of pPIC9, Invitrogen,
24 Carlsbad, CA) was digested with restriction endonucleases XhoI and EcoRI, and gel
25 purified, resulting in an approximately 2900 base pair vector fragment. The purified
26 restriction fragments were combined and ligated using T4 DNA ligase.

27 The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells
28 (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on
29 ampicillin-containing plates. Plasmid DNA was isolated and sequenced to confirm the

1 presence of the correct insert. The resulting plasmid was designated pMON 37402
2 (containing SEQ ID NO:246, encoding protein sequence SEQ ID NO:247).

3 Example 8: Construction of plasmid pMON 37405

4 Amino acids 201-202, near tyrosine 193, were chosen as a breakpoint for the
5 construction of a permutein protein.

6 The nucleic acid sequence encoding the permutein protein in plasmid pMON
7 37405 was created using the method illustrated in Figure 3 and described in Example 6.
8 Nucleic acid molecule "New N-termini fragment" was created and amplified from the
9 sequence encoding patatin in plasmid pMON26820 using oligonucleotide primers 48
10 (SEQ ID NO:243) and 58 (SEQ ID NO:249). Nucleic acid molecule "New C-termini
11 fragment" was created and amplified from the sequence encoding patatin in plasmid
12 pMON26820 using oligonucleotide primers 47 (SEQ ID NO:244) and 59 (SEQ ID
13 NO:249). The full-length nucleic acid molecule encoding the permutein protein was
14 created and amplified from annealed fragments "New N-termini fragment" and "New C-
15 termini fragment" using oligonucleotide primers 58 (SEQ ID NO:248) and 59 (SEQ ID
16 NO:249).

17 The resulting amplified nucleic acid molecule was digested with restriction
18 endonucleases XhoI and EcoRI, and purified using the QIAquick PCR purification kit
19 (Qiagen, Valencia, CA). Plasmid pMON 26869 (derivative of pPIC9, Invitrogen,
20 Carlsbad, CA) was digested with restriction endonucleases XhoI and EcoRI, and gel
21 purified, resulting in an approximately 2900 base pair vector fragment. The purified
22 restriction fragments were combined and ligated using T4 DNA ligase.

23 The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells
24 (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on
25 ampicillin-containing plates. Plasmid DNA was isolated and sequenced to confirm the
26 presence of the correct insert. The resulting plasmid was designated pMON 37405
27 (containing SEQ ID NO:250, encoding protein sequence SEQ ID NO:251).

Example 9: Construction of plasmid pMON 37406

Amino acids 183-184, adjacent to tyrosine 185, were chosen as a breakpoint for the construction of a permutein protein.

The nucleic acid sequence encoding the permutein protein in plasmid pMON 37406 was created using the method illustrated in Figure 3 and described in Example 6. Nucleic acid molecule "New N-termini fragment" was created and amplified from the sequence encoding patatin in plasmid pMON26820 using oligonucleotide primers 48 (SEQ ID NO:243) and 60 (SEQ ID NO:252). Nucleic acid molecule "New C-termini fragment" was created and amplified from the sequence encoding patatin in plasmid pMON26820 using oligonucleotide primers 47 (SEQ ID NO:244) and 61 (SEQ ID NO:253). The full-length nucleic acid molecule encoding the permutein protein was created and amplified from annealed fragments "New N-termini fragment" and "New C-termini fragment" using oligonucleotide primers 60 (SEQ ID NO:252) and 61 (SEQ ID NO:253).

The resulting amplified nucleic acid molecule was digested with restriction endonucleases XhoI and EcoRI, and purified using the QIAquick PCR purification kit (Qiagen, Valencia, CA). Plasmid pMON 26869 (derivative of pPIC9, Invitrogen, Carlsbad, CA) was digested with restriction endonucleases XhoI and EcoRI, and gel purified, resulting in an approximately 2900 base pair vector fragment. The purified restriction fragments were combined and ligated using T4 DNA ligase.

The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on ampicillin-containing plates. Plasmid DNA was isolated and sequenced to confirm the presence of the correct insert. The resulting plasmid was designated pMON37406 (containing SEQ ID NO:254, encoding protein sequence SEQ ID NO:255).

Example 10: Construction of plasmid pMON 37407

Amino acids 268-269, adjacent to tyrosine 270, were chosen as a breakpoint for the construction of a permutein protein.

1 The nucleic acid sequence encoding the permutein protein in plasmid pMON
2 37407 was created using the method illustrated in Figure 3 and described in Example 6.
3 Nucleic acid molecule "New N-termini fragment" was created and amplified from the
4 sequence encoding patatin in plasmid pMON26820 using oligonucleotide primers 48
5 (SEQ ID NO:243) and 62 (SEQ ID NO:256). Nucleic acid molecule "New C-termini
6 fragment" was created and amplified from the sequence encoding patatin in plasmid
7 pMON26820 using oligonucleotide primers 47 (SEQ ID NO:244) and 63 (SEQ ID
8 NO:257). The full-length nucleic acid molecule encoding the permutein protein was
9 created and amplified from annealed fragments "New N-termini fragment" and "New C-
10 termini fragment" using oligonucleotide primers 62 (SEQ ID NO:256) and 63 (SEQ ID
11 NO:257).

12 The resulting amplified nucleic acid molecule was digested with restriction
13 endonucleases XhoI and EcoRI, and purified using the QIAquick PCR purification kit
14 (Qiagen, Valencia, CA). Plasmid pMON 26869 (derivative of pPIC9, Invitrogen,
15 Carlsbad, CA) was digested with restriction endonucleases XhoI and EcoRI, and gel
16 purified, resulting in an approximately 2900 base pair vector fragment. The purified
17 restriction fragments were combined and ligated using T4 DNA ligase.

18 The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells
19 (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on
20 ampicillin-containing plates. Plasmid DNA was isolated and sequenced to confirm the
21 presence of the correct insert. The resulting plasmid was designated pMON37407
22 (containing SEQ ID NO:258, encoding protein sequence SEQ ID NO:259).

23 Example 11: Construction of plasmid pMON 37408

24 Amino acids 321-322, near tyrosine 216, were chosen as a breakpoint for the
25 construction of a permutein protein.

26 The nucleic acid sequence encoding the permutein protein in plasmid pMON
27 37408 was created using the method illustrated in Figure 3 and described in Example 6.
28 Nucleic acid molecule "New N-termini fragment" was created and amplified from the
29 sequence encoding patatin in plasmid pMON26820 using oligonucleotide primers 48
30 (SEQ ID NO:243) and 64 (SEQ ID NO:260). Nucleic acid molecule "New C-termini

1 fragment” was created and amplified from the sequence encoding patatin in plasmid
2 pMON26820 using oligonucleotide primers 47 (SEQ ID NO:244) and 65 (SEQ ID
3 NO:261). The full-length nucleic acid molecule encoding the permutein protein was
4 created and amplified from annealed fragments “New N-termini fragment” and “New C-
5 termini fragment” using oligonucleotide primers 64 (SEQ ID NO:260) and 65 (SEQ ID
6 NO:261).

7 The resulting amplified nucleic acid molecule was digested with restriction
8 endonucleases XhoI and EcoRI, and purified using the QIAquick PCR purification kit
9 (Qiagen, Valencia, CA). Plasmid pMON 26869 (derivative of pPIC9, Invitrogen,
10 Carlsbad, CA) was digested with restriction endonucleases XhoI and EcoRI, and gel
11 purified, resulting in an approximately 2900 base pair vector fragment. The purified
12 restriction fragments were combined and ligated using T4 DNA ligase.

13 The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells
14 (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on
15 ampicillin-containing plates. Plasmid DNA was isolated and sequenced to confirm the
16 presence of the correct insert. The resulting plasmid was designated pMON37408
17 (containing SEQ ID NO:262, encoding protein sequence SEQ ID NO:263).

18 Example 12: Production of permutein proteins in *Pichia pastoris*

19 Plasmids pMON37402, pMON37405, pMON37406, pMON37407, and
20 pMON37408 were individually used to electroporate KM71 cells from *Pichia pastoris*
21 according to the procedure supplied by the manufacturer (Invitrogen, Carlsbad, CA). The
22 resulting transformed cells were used to produce protein in *Pichia pastoris* following the
23 procedure supplied by the manufacturer (Invitrogen, Carlsbad, CA).

24 The concentration of patatin in the culture was determined using a patatin ELISA
25 assay and the enzyme activity was measured using the method of Hofgen and Willmitzer
26 (*Plant Science*, 66: 221-230, 1990). The variants containing multiple mutations were
27 further purified using Mono Q and hydrophobic interaction chromatography (HIC). Each
28 culture was purified by first sizing on YM10 membranes (Amicon, MA) to a [>10 kDa]
29 fraction, followed by chromatography on the Mono Q HR 10/10 column (Pharmacia, NJ).
30 For chromatography on the Mono Q column, the samples were loaded on the column in

25 mM Tris pH 7.5 and eluted with a gradient of 1.0 M KCl in 25 mM Tris pH 7.5. Fractions containing patatin protein were determined using SDS-PAGE. For chromatography on the HIC column, the appropriate fractions were pooled and dialyzed into 1 M ammonium sulfate in 25 mM Tris pH 7.5. The dialyzed sample was then loaded on 16/10 phenyl Sepharose column (Pharmacia, NJ) and eluted with a gradient of 25 mM Tris pH 7.5.

The protein concentration was determined using the Bradford method, using BSA as a standard. SDS-PAGE analysis showed that these proteins were essentially pure. The esterase activity of the variants are shown in Table 7.

Table 7: Activity of permuteins

pMON	Breakpoint	Activity ($\Delta OD \text{ min}^{-1} \mu\text{g}^{-1}$)
Native enzyme		83.21
pMON37402	246/247	66.7
pMON37405	201/202	No expression
pMON37406	183/184	No expression
pMON37407	268/269	12.1
pMON37408	321/322	No expression

The activity was determined using *p*-nitrophenyl caprate substrate as described by Hofgen and Willmitzer (*Plant Science*, 66: 221-230, 1990).

Example 13: Insect bioefficacy assays

Assays for activity against larvae of SCRW are carried out by overlaying the test sample on an agar diet similar to that described by Marrone (*J. Econ. Entom.* 78: 290-293, 1985). Test samples were prepared in 25 mM Tris, pH 7.5 buffer. Neonate larvae are allowed to feed on the treated diet at 26°C, and mortality and growth stunting were evaluated after 5 or 6 days. The results of this assay are shown in Table 8.

Table 8: Insect bioefficacy assay

Protein (200 ppm)	Mean Survival Weight	% Weight Reduction
Tris buffer (control)	1.26 \pm 0.3	-
Wild Type	0.21 \pm 0.02	83

pMON37402	0.21 ± 0.03	83
pMON37407	0.32 ± 0.04	75

These data demonstrate that the growth of the SCRW larvae is similarly reduced upon ingestion of the proteins encoded by pMON37402 and pMON37407 as compared to the wild type patatin protein.

Example 14: Permutein sequences improved for monocot expression

Modification of coding sequences has been demonstrated above to improve expression of insecticidal proteins. A modified coding sequence was thus designed to improve expression in plants, especially corn (SEQ ID NO:264).

Example 15: Construction of pMON40701 for monocot expression

Plasmid pMON19767 was digested with restriction endonucleases NcoI and EcoRI and the 1100 bp gene fragment was purified using the QIAquick PCR purification kit (Qiagen, Valencia, CA). Plasmid pMON33719 was digested with restriction endonucleases NcoI and EcoRI, and gel purified, resulting in an approximately 3900 base pair vector fragment. The two purified restriction fragments were combined and ligated using T4 DNA ligase.

The ligation reaction mixture was used to transform *E. coli* strain DH5α cells (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on ampicillin-containing plates. Plasmid DNA was isolated and sequenced to confirm the presence of the correct insert. The resulting plasmid was designated pMON40700. Plasmid pMON40700 was digested with restriction endonuclease NotI and the resulting 2200 bp DNA fragment was purified using the QIAquick PCR purification kit (Qiagen, Valencia, CA). Plasmid pMON30460 was digested with restriction endonuclease NotI, and gel purified, resulting in an approximately 4200 base pair vector fragment. The two purified restriction fragments were combined and ligated using T4 DNA ligase.

The ligation reaction mixture was used to transform *E. coli* strain DH5α cells (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on

1 kanamycin-containing plates. The resulting plasmid was designated pMON40701
2 (containing SEQ ID NO:264, encoding protein sequence SEQ ID NO:265).

3 Example 16: Construction of pMON40703 for monocot expression

4 The nucleic acid sequence encoding the permutein protein in plasmid
5 pMON40703 was created using the method illustrated in Figure 3 and described in
6 Example 6. Nucleic acid molecule “New N-termini fragment” was created and amplified
7 from the sequence encoding patatin in plasmid pMON19767 using oligonucleotide
8 primers Syn1 (SEQ ID NO:266) and Syn2 (SEQ ID NO:267). Nucleic acid molecule
9 “New C-termini fragment” was created and amplified from the sequence encoding patatin
10 in plasmid pMON19767 using oligonucleotide primers Syn3 (SEQ ID NO:268) and Syn4
11 (SEQ ID NO:269). The full-length nucleic acid molecule encoding the permutein protein
12 was created and amplified from annealed fragments “New N-termini fragment” and
13 “New C-termini fragment” using oligonucleotide primers Syn1 (SEQ ID NO:266) and
14 Syn4 (SEQ ID NO:269).

15 The resulting amplified nucleic acid molecule was digested with restriction
16 endonucleases NcoI and EcoRI, and purified using the QIAquick PCR purification kit
17 (Qiagen, Valencia, CA). Plasmid pMON33719 was digested with restriction
18 endonucleases NcoI and EcoRI, and gel purified, resulting in an approximately 3900 base
19 pair vector fragment. The purified restriction fragments were combined and ligated using
20 T4 DNA ligase.

21 The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells
22 (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on
23 ampicillin-containing plates. Plasmid DNA was isolated and sequenced to confirm the
24 presence of the correct insert. The resulting plasmid was designated pMON40702.
25 Plasmid pMON40702 was digested with NotI, and the resulting 2200 bp DNA fragment
26 was purified using the QIAquick PCR purification kit (Qiagen, Valencia, CA). Plasmid
27 pMON30460 was digested with restriction endonuclease NotI, and gel purified, resulting
28 in an approximately 4200 base pair vector fragment. The purified restriction fragments
29 were combined and ligated using T4 DNA ligase.

1 The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells
2 (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on
3 kanamycin-containing plates. The resulting plasmid was designated pMON40703
4 (containing SEQ ID NO:270, encoding protein sequence SEQ ID NO:271). Plasmid
5 pMON40703 encodes a permutein protein with a “breakpoint” at positions 246/247 of the
6 wild type patatin protein sequence (SEQ ID NO:2). The first 23 amino acids of SEQ ID
7 NO:2 are a signal peptide sequence which is cleaved in the mature protein.

8 Example 17: Construction of pMON40705 for monocot expression

9 The nucleic acid sequence encoding the permutein protein in plasmid
10 pMON40705 was created using the method illustrated in Figure 3 and described in
11 Example 6. Nucleic acid molecule “New N-termini fragment” was created and amplified
12 from the sequence encoding patatin in plasmid pMON19767 using oligonucleotide
13 primers Syn10 (SEQ ID NO:272) and Syn2 (SEQ ID NO:267). Nucleic acid molecule
14 “New C-termini fragment” was created and amplified from the sequence encoding patatin
15 in plasmid pMON19767 using oligonucleotide primers Syn3 (SEQ ID NO:268) and
16 Syn11 (SEQ ID NO:273). The full-length nucleic acid molecule encoding the permutein
17 protein was created and amplified from annealed fragments “New N-termini fragment”
18 and “New C-termini fragment” using oligonucleotide primers Syn10 (SEQ ID NO:272)
19 and Syn11 (SEQ ID NO:273).

20 The resulting amplified nucleic acid molecule was digested with restriction
21 endonucleases NcoI and EcoRI, and purified using the QIAquick PCR purification kit
22 (Qiagen, Valencia, CA). Plasmid pMON33719 was digested with restriction
23 endonucleases NcoI and EcoRI, and gel purified, resulting in an approximately 3900 base
24 pair vector fragment. The purified restriction fragments were combined and ligated using
25 T4 DNA ligase.

26 The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells
27 (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on
28 ampicillin-containing plates. Plasmid DNA was isolated and sequenced to confirm the
29 presence of the correct insert. The resulting plasmid was designated pMON40704.
30 Plasmid pMON40704 was digested with restriction endonuclease NotI, and the resulting

1 2200 bp DNA fragment was purified using the QIAquick PCR purification kit (Qiagen,
2 Valencia, CA). Plasmid pMON30460 was digested with restriction endonuclease NotI,
3 and gel purified, resulting in an approximately 4200 base pair vector fragment. The
4 purified restriction fragments were combined and ligated using T4 DNA ligase.

5 The ligation reaction mixture was used to transform *E. coli* strain DH5 α cells
6 (Life Technologies, Gaithersburg, MD). Transformant bacteria were selected on plates
7 containing kanamycin. The resulting plasmid was designated pMON40705 (containing
8 SEQ ID NO:274, encoding protein sequence SEQ ID NO:275). Plasmid pMON40705
9 encodes a permutein protein with a “breakpoint” at positions 268/269 of the wild type
10 patatin protein sequence (SEQ ID NO:2). The first 23 amino acids of SEQ ID NO:2 are a
11 signal peptide sequence which is cleaved in the mature protein.

12 Example 18: Transient expression of protein in corn leaf protoplasts

13 Plasmids pMON40701, pMON40703, and pMON40705 (all containing the native
14 signal sequence for vacuolar targeting) were separately electroporated into corn leaf
15 protoplasts as described by Sheen (*Plant Cell* 3: 225-245, 1991). Protein was extracted
16 with glass beads and the supernatant was assayed for protein expression using ELISA for
17 patatin and NPTII. Expression of protein by the transformed corn protoplasts was
18 confirmed by Western blot analysis. Expression results are shown in Table 9.

19 Table 9: ELISA data

Sample	Patatin ELISA ($\mu\text{g/mL}$)	NPTII ELISA ($\mu\text{g/mL}$)	Normalized Expression (Patatin ELISA/NPTII ELISA)
pMON40701	1.1	0.6	1.8
pMON40703	2.1	0.3	7.0
pMON40705	1.3	0.6	2.2

20 The results indicate that the permutein encoded by plasmid pMON40703
21 surprisingly shows approximately 4-fold higher expression compared to the wild type
22 enzyme.

Example 19: Deglycosylation of protein sequences

This example provides evidence that glycosylation of can contribute to the allergenicity of a protein. Accordingly, rational substitution of amino acid residues likely to be the targets of glycosylation within a subject allergen protein may reduce or eliminate the allergenic properties of the protein without adversely affecting the enzymatic, insecticidal, antifungal or other functional properties of the protein.

Glycosylation commonly occurs as either N-linked or O-linked forms. N-linked glycosylation usually occurs at the motif Asn-Xaa-Ser/Thr, where Xaa is any amino acid except Pro (Kasturi, L. et al., *Biochem J.* 323: 415-519, 1997; Melquist, J.L. et al., *Biochemistry* 37: 6833-6837, 1998). O-linked glycosylation occurs between the hydroxyl group of serine or threonine and an amino sugar.

Site directed mutagenesis of selected asparagine, serine, or threonine may be used to reduce or eliminate the glycosylation of patatin proteins. A search of SEQ ID NO:2 for the Asn-Xaa-Ser/Thr motif reveals one occurrence at amino acid positions 202-204. Mutagenization of the nucleic acid sequence encoding this region results in a reduced allergenicity of the encoded protein.

In order to test this approach to reducing allergenicity of patatin proteins, two sets of experiments were performed: a) production of patatin proteins in *Escherichia coli*, which do not glycosylate proteins; and b) production of patatin proteins with an N202Q site directed mutation.

Antibodies obtained from patients HS-07 and G15-MON (not potato allergic) did not show specific binding to wild type patatin, patatin produced in *E. coli*, or the N202Q variant. Antibodies obtained from patient HS-01 (potato allergic) bound to wild type patatin, but not to patatin produced in *E. coli* or the N202Q variant. Antibodies obtained from patient HS-02 (potato allergic) bound strongly to wild type patatin, but extremely weakly to patatin produced in *E. coli*, and binding to the N202Q variant resembled vector controls. Antibodies obtained from patient HS-03 (potato allergic) bound to wild type patatin, but not to patatin produced in *E. coli* or the N202Q variant. Antibodies obtained from patient HS-05 (potato allergic) bound to wild type patatin, but very weakly to patatin produced in *E. coli* and the N202Q variant. Antibodies obtained from patient HS-06 (potato allergic) strongly bound wild type patatin, the N202Q variant, and to patatin

1 produced in *E. coli*. These results strongly suggest that glycosylation is at least partially
2 responsible for the antigenic properties of patatin proteins, and that site directed
3 mutagenesis may be used to reduce or eliminate specific antibody binding. Mutagenesis
4 at position 202 of SEQ ID NO:2 may be useful for reducing or eliminating specific
5 antibody binding.

6 The deglycosylation approach was also tested using a patatin homolog, Pat17. As
7 demonstrated above, patatin epitopes exhibiting IgE binding were identified, and each
8 contained a Tyr residue. Substitution of these Tyr residues within each epitope led to loss
9 of IgE binding. Site-directed mutagenesis was used to produce variants with individual
10 and multiple Tyr substitutions in the protein, which was expressed in *Pichia pastoris* and
11 assessed for enzyme activity. All the variants were found to have enzymatic activity no
12 less than the wild type protein. A single variant with all 5 tyrosine residues substituted
13 with phenylalanine was found to have insecticidal activity no less than the unsubstituted
14 protein and was expressed in *E.coli* to produce the non-glycosylated version. The *E.coli*
15 5-"Tyr to Phe" variant was assessed for IgE binding. An isozyme of patatin, designated
16 Pat17, was also expressed in corn to produce a plant glycoprotein and in *E.coli* to
17 produce a nonglycosylated protein. Sera of seven patients (five exhibiting potato allergy
18 and one exhibiting other allergies but no allergy to potatoes) were used to assay
19 Pat17 or Pat17 variant binding by immunoblot assay. Four of the five sera from patients
20 exhibiting potato allergy showed either very weak or no binding to wild type patatin
21 expressed in *E.coli* but did bind to the 5-Tyr variant. Serum from one patient exhibiting
22 potato allergy showed strong binding to recombinant wild type patatin protein expressed
23 in *E.coli* but weak binding to the 5-Tyr variant. Sera from all five patients exhibiting
24 potato allergy bound strongly to patatin expressed in corn and native patatin present in
25 potatoes. Serum from a control patient allergic to eggs, milk, peanuts and seafood, but
26 exhibiting no allergy to potatoes showed no binding to patatin expressed in *E.coli* but did
27 bind to patatin expressed in corn. Immunoblot results suggested that the sugar moiety in
28 patatin is a non-specific IgE binding epitope and the polypeptide portion of patatin also
29 contains immunogenic IgE epitopes.

1 Patients who suffer from potato allergy were identified at Johns Hopkins Clinic
2 (Baltimore, MD) and were evaluated for potato allergy using clinical criteria outlined in
3 Table 2.

4 Serum was obtained from patients with convincing clinical history of potato
5 allergy. The convincing history was defined as being one or more of the following: a)
6 positive potato allergic reaction as evaluated by double-blind placebo-control food
7 challenge b) anaphylaxis and/or hospitalization due to the consumption of potatoes or c)
8 dramatic skin test results.

9 Peptide Synthesis

10 Peptides were synthesized on cellulose membranes using the SPOTS system
11 (Genosys Biotechnologies, TX). Membranes were stored at -20°C until use.

12 Site directed Mutagenesis

13 Site specific mutations were introduced into patatin by first incorporating part of
14 the a-factor signal sequence (*Pichia* expression manual, Invitrogen, Carlsbad, CA) to the
15 patatin gene using PCR. Primers used for the PCR were
16 GGAGCTCGAGAAAAGAGAGGCTGAAGCTCAGTTGGGAGAAATGGTGACTGT
17 TCT (*XhoI* site in italics) and GGTCTAGAG GAATTCTCATTAATAAGAAG (*EcoRI*
18 site in italics). The primers contained restriction sites to facilitate cloning into *Pichia*
19 *pastoris* yeast secretion vector pPIC9 (GenBank accession number Z46233; Invitrogen,
20 Carlsbad, CA). Typical PCR conditions are 25 cycles 94°C denaturation for 1 minute,
21 45°C annealing for one minute and 72°C extension for 2 minutes; plus one cycle 72°C
22 extension for 10 minutes. A 50 mL reaction contained 30 pmol of each primer and 1 mg
23 of template DNA; and 1 X PCR buffer with MgCl₂, 200 mM dGTP, 200 mM dATP, 200
24 mM dTTP, 200 mM dCTP, 2.5 units of *Pwo* DNA polymerase. PCR reactions are
25 performed in RoboCycler Gradient 96 Temperature Cycler (Stratagene, La Jolla, CA).

26 The amplified patatin gene was digested with restriction enzymes *XhoI* and *EcoRI*
27 and cloned into the pBluescript vector (Stratagene, La Jolla, CA), digested with the same
28 two restriction enzymes. The template plasmid DNA used for the PCR was
29 pMON26820. The resulting plasmid (pMON 26869) was used for oligonucleotide-
30 directed mutagenesis using the Bio-Rad mutagenesis kit based on the method of Kunkel

1 et al., *Proc Natl Acad Sci USA* 82, 477-92 (1985). Briefly, single-stranded pMON26869
2 was used as template for mutagenesis and was prepared by superinfection of plasmid
3 containing cells with M13K07 (Gorman et al., *DNA and Protein Engineering techniques*
4 2, 3-10 (1990)). DNA purified from transformed DH5a *E. coli* colonies was used for
5 sequence determination. Sequencing was performed using the ABI PRISM sequencing
6 kit (Perkin Elmer Biosystems, Foster City, CA).

7 Protein Expression in *Pichia pastoris*

8 Plasmids containing the mutations in the patatin gene were digested with restriction
9 enzymes *Xho*I and *Eco*RI. The patatin nucleic acid fragment was then ligated into the
10 pPIC9 vector (Invitrogen, Carlsbad, CA), digested with the same two restriction enzymes
11 to afford plasmid pMON37401. *Pichia pastoris* KM71 cells were electroporated with
12 pMON37401 containing the appropriate mutation. The resulting transformed cells were
13 used to produce protein in *Pichia pastoris* using the procedure supplied by the
14 manufacturer (Invitrogen, Carlsbad, CA). The proteins were purified in the same way as
15 the proteins expressed in *E. coli* (see below).

16 Western Blotting of Proteins

17 Protein samples were electrophoresed by SDS-PAGE and electroblotted onto PVDF
18 membrane (Millipore, Bedford MA). Protein blots were processed by standard Western
19 blotting (immunoblotting) techniques and were incubated in potato allergic serum diluted
20 1:5 in PBS buffer for 1 hour. After washing the blots 3 times with PBS, the blots were
21 incubated in biotinylated anti-IgE (Johns Hopkins Hospital, Baltimore MD) for 1 hour,
22 followed by a 30 minute incubation in HRP-linked avidin (Promega, New York, NY).
23 IgE-reactive protein bands were visualized by using the ECL system (Amersham
24 Pharmacia Biotech, NJ). As a control, one blot was incubated in anti-IgE only. His-
25 tagged glyphosate oxidase and potato extracts was prepared and provided for this study
26 by Regulatory Sciences, Monsanto Company. The peptides were evaluated using the
27 same incubation procedures as described above.

28 Expression and purification of patatin in corn

29 An isozyme of patatin, Pat17, was generated for expression in corn using a modified
30 plant optimized gene sequence as described by Brown *et al* (US Patent 5,689,052). All

the constructs contained the native 23 amino acid signal peptide for vacuolar targeting. Corn was transformed by microprojectile bombardment (Morrish et al., in *Transgenic plants. Fundamentals and Applications* (ed. Hiatt, A.) 133-171 (Marcel Dekker, New York, 1993); Songstad et al., *In Vitro Cell Dev Biol - Plant* 32, 179-183 (1996)). Protein from the transformed corn plants was purified by first grinding the leaves in liquid nitrogen and extracting the protein using 25 mM Tris/HCl. The plant extract was further dialyzed against 25 mM Tris/HCl pH 7.5. The plant extract was then loaded onto Mono Q HR 10/10 anion-exchange column (Amersham Pharmacia, NJ) equilibrated with 25 mM Tris/HCl pH 7.5 (buffer A). The protein was eluted with 25 mM Tris/HCl pH 7.5, 1 M KCl (buffer B) using a linear gradient of 0-100% buffer B using an HPLC system (Shimadzu). Fractions containing protein were assayed for esterase activity and dialyzed against 25 mM Tris/HCl pH 7.5, 1 M Ammonium Sulfate (buffer C). The protein was purified to homogeneity by loading onto a phenyl-Sepharose 16/10 column (Amersham Pharmacia, NJ) equilibrated with buffer C. Esterase active fractions were pooled and dialyzed against 25 mM Tris/HCl pH 7.5.

Expression and purification of patatin in *E.coli*

Pat17 was expressed in *E.coli* using the pET expression system (Novagen, WI). The coding region of the mature Pat17 gene (without its signal peptide) was amplified by PCR using the primers 5'-GGGCCATGGCGCAGTTGGGAGAAATGGTG-3' (*NcoI* site in italics) and 5'-AACAAAGCTTCTTATTGAGGTGCGGCCGCTTGCATGC-3' (*NotI* site in italics) using standard PCR reaction conditions as described in the Gene Amp kit (Perkin-Elmer Cetus, CT) and an annealing temperature of 40°C. The template was plasmid pMON26820. The resulting DNA was digested with *NcoI* and *NotI* and cloned into a modified pET24d plasmid, designed to add an N-terminal hexa-histidine tag to the protein. The correct sequence of the PCR product was verified by sequencing, and the plasmid was transformed into *E.coli* BL21 (DE3), and transformants selected on LB containing 25 mg/mL kanamycin. The expression strain was grown in LB containing 25 mg/mL kanamycin and induced for 8 hrs at 28°C with 1 mM IPTG. Cells were harvested and washed in 50 mM Tris/HCl pH 8.5, 150 mM NaCl, and lysed by French Press at 20,000 psi. His-tagged protein was recovered in the soluble fraction of lysed cells and subsequently purified using Ni-NTA resin as described in the QIAexpressionist manual

(Qiagen CA). The partially purified protein was then dialyzed against 25 mM Tris/HCl pH 7.5 (buffer A) and loaded onto Mono Q HR 10/10 anion-exchange column (Amersham Pharmacia, NJ) equilibrated with buffer A. The protein was eluted with 25 mM Tris/HCl pH 7.5, 1 M KCl (buffer B) using a linear gradient of 0-100% buffer B run over 30 min at a flow rate of 4 mL/min using an HPLC system (Shimadzu). Fractions containing protein were assayed for esterase activity. Esterase active fractions were pooled, concentrated and dialyzed against 25 mM Tris/HCl pH 7.5 and stored at 4°C.

Enzyme Activity Assays

Enzyme activity was measured as described previously using *p*-nitrophenyl caprate (Sigma, MO) as a substrate, dissolved in dimethylsulfoxide (5 mM stock solution) and diluted in 4% Triton X-100, 1% SDS to a final concentration of 1 mM. For the assay, 20 mL of protein solution was added to a mixture of 25 mL of the 1 mM substrate solution and 80 mL of 50 mM Tris pH 8.5. The enzyme activity was monitored at 405 nm in 6 sec interval for a period of 10 min. Esterase activity was expressed as DOD min⁻¹mg⁻¹ protein.

Insect Bioassay

The protein was also assayed for activity against larvae of *Diabrotica virgifera* (Western corn rootworm) by overlaying the test sample on an agar diet similar to that described previously (Marrone et al., *J. Econ. Entom.* 78, 290-3 (1985)). Proteins to be tested were diluted in 25 mM Tris/HCl pH 7.5 and overlayed on the diet surface. Neonate larvae were allowed to feed on the diet and mortality and growth stunting were evaluated after 6 days.

IgE Binding Epitopes on Patatin

A panel of eighty-nine overlapping peptides representing the amino acid sequence of patatin were synthesized to determine the regions responsible for IgE binding. Each peptide was 10 amino acids long and consisted of 6 amino acid overlap between the consecutive peptides. The peptides were evaluated for IgE binding with five different potato allergic patient sera. Patatin has 3 major epitopes. These major IgE binding regions represent amino acids 184-193, 188-197, 269-278 and 360-369. Other minor IgE binding regions represent amino acids 104-113, 138-147 and 316-325. The amino acids

essential for IgE binding in each major and minor epitopes were determined by synthesizing peptides with single amino acid changes at each position by individually substituting an alanine residue at each non-alanine position in the epitopes. The resulting alanine substituted peptides were evaluated for IgE binding. Result effective substitutions were identified by a reduction in IgE binding with respect to the non-substituted peptide sequence. It was very interesting to note that all the epitopes contained a Tyr residue and substitution of this Tyr for Ala or Phe eliminated IgE binding.

Enzyme and Bioactivity

The Tyr residues identified to be critical for IgE binding in each of the epitopes were substituted with Phe either individually or in concert using site-directed mutagenesis. All the variants were expressed in *Pichia pastoris* and assessed for enzyme activity and insecticidal activity. The variants included Y106F, Y129F, Y185F, Y193F, Y270F, Y316F, Y362F, Y185F/Y193F, Y185F/Y193F/Y270F/Y316F/Y362F (5-Tyr) and Y106F/Y129F/Y185F/Y193F/Y270F/Y316F/ Y362F (7-Tyr). All the variants maintained enzyme activity. The 5-Tyr and 7-Tyr variants were then assessed for insecticidal activity by overlaying protein (200 ppm final concentration). The proteins caused significant stunting of the larval growth as measured by the weight of the larvae after 6 days with the 5-Tyr variant showing higher insecticidal activity compared to the 7-Tyr and wild type proteins. The 7-Tyr variant was unstable upon long term storage at 4°C and thus was not pursued further.

Immunoblotting

In order to test if the glycan moiety on patatin was important for binding of IgE, Pat17 was expressed in *E.coli* to produce a nonglycosylated protein and in corn to produce a plant glycosylated protein. The 5-Tyr variant was also expressed in *E.coli* to assess the individual contribution of the linear epitopes without the glycan moiety on the protein. The proteins were tested for binding to IgE using sera from five patients with allergy to potatoes and sera from one patient with allergies to many things but no allergy to potatoes. Proteins from both corn and *E.coli* were purified to homogeneity. These proteins were transferred to PVDF membrane (Millipore, MA) and subsequently probed

1 with sera from patients with and without allergy to potatoes. A His-tagged glyphosate
2 oxidase control was included in all the studies to verify that the His-tag did not affect the
3 binding of IgE. Serum obtained from patient HS-07 (no allergy to potatoes) did not bind
4 Pat17 expressed in *E.coli* but showed good binding to Pat17 from corn and also a protein
5 at the same molecular weight in potato extract. It is interesting to note that this sera also
6 showed strong binding to another protein (> 46kDa) in the potato. Sera from patients
7 HS-01, HS-02, HS-03, HS-05 (allergy to potatoes) shows strong binding to Pat17
8 expressed in corn, but very weak to no binding to Pat17 produced in *E.coli*. Also, the
9 sera from patients HS-01, HS-2, HS-03 and HS-05 bound to a protein of similar
10 molecular weight in the potato extract. Sera from patients HS-01, HS-02 and HS-03 also
11 showed binding to another protein in potato extract of a lower molecular weight (<
12 30kDa). Serum obtained from patient HS-06 (allergic to potatoes) showed very strong
13 binding to wild type patatin expressed in both corn and *E.coli* but weaker binding to the
14 5-Tyr variant expressed in *E.coli*. Sera from HS-06 also showed very strong binding to a
15 protein in potato extract with similar molecular weight as patatin. The sera from all the
16 patients showed no binding to His-tagged glyphosate oxidase indicating that the His-tag
17 does not bind IgE. These results strongly suggest that the glycan moiety on Pat17 is
18 responsible for IgE binding in some potato allergic patients and linear epitopes also
19 contribute to the antigenicity of patatin.

20 Example 20: Alternative nucleic acid and protein sequences

21 For future variations of the patatin protein, sequences showing high similarity to
22 the sequences disclosed herein could be used in producing deallergenized patatin proteins
23 and permuteins. For example, a BLAST search (Altschul, S.F. et al., *J. Mol. Biol.* 215:
24 403-410, 1990) can be performed to identify additional patatin sequences. Sources other
25 than those disclosed herein can be used to obtain a patatin nucleic acid sequence, and the
26 encoded patatin protein. Furthermore, subunit sequences from different organisms can be
27 combined to create a novel patatin sequence incorporating structural, regulatory, and
28 enzymatic properties from different sources.

Example 21: Nucleic acid mutation and hybridization

Variations in the nucleic acid sequence encoding a patatin protein may lead to mutant patatin protein sequences that display equivalent or superior enzymatic characteristics when compared to the sequences disclosed herein. This invention accordingly encompasses nucleic acid sequences which are similar to the sequences disclosed herein, protein sequences which are similar to the sequences disclosed herein, and the nucleic acid sequences that encode them. Mutations can include deletions, insertions, truncations, substitutions, fusions, shuffling of subunit sequences, and the like.

Mutations to a nucleic acid sequence can be introduced in either a specific or random manner, both of which are well known to those of skill in the art of molecular biology. A myriad of site-directed mutagenesis techniques exist, typically using oligonucleotides to introduce mutations at specific locations in a nucleic acid sequence. Examples include single strand rescue (Kunkel, T. *Proc. Natl. Acad. Sci. U.S.A.*, 82: 488-492, 1985), unique site elimination (Deng and Nickloff, *Anal. Biochem.* 200: 81, 1992), nick protection (Vandeyar, et al. *Gene* 65: 129-133, 1988), and PCR (Costa, et al. *Methods Mol. Biol.* 57: 31-44, 1996). Random or non-specific mutations can be generated by chemical agents (for a general review, see Singer and Kusmierek, *Ann. Rev. Biochem.* 52: 655-693, 1982) such as nitrosoguanidine (Cerdeña-Olmedo et al., *J. Mol. Biol.* 33: 705-719, 1968; Guerola, et al. *Nature New Biol.* 230: 122-125, 1971) and 2-aminopurine (Rogan and Bessman, *J. Bacteriol.* 103: 622-633, 1970), or by biological methods such as passage through mutator strains (Greener et al. *Mol. Biotechnol.* 7: 189-195, 1997).

Nucleic acid hybridization is a technique well known to those of skill in the art of DNA manipulation. The hybridization properties of a given pair of nucleic acids is an indication of their similarity or identity. Mutated nucleic acid sequences can be selected for their similarity to the disclosed patatin nucleic acid sequences on the basis of their hybridization to the disclosed sequences. Low stringency conditions can be used to select sequences with multiple mutations. One may wish to employ conditions such as about 0.15 M to about 0.9 M sodium chloride, at temperatures ranging from about 20°C to about 55°C. High stringency conditions can be used to select for nucleic acid sequences with higher degrees of identity to the disclosed sequences. Conditions employed may

include about 0.02 M to about 0.15 M sodium chloride, about 0.5% to about 5% casein, about 0.02% SDS and/or about 0.1% N-laurylsarcosine, about 0.001 M to about 0.03 M sodium citrate, at temperatures between about 50°C and about 70°C. More preferably, high stringency conditions are 0.02 M sodium chloride, 0.5% casein, 0.02% SDS, 0.001 M sodium citrate, at a temperature of 50°C.

Example 22: Determination of homologous and degenerate nucleic acid sequences

Modification and changes can be made in the sequence of the proteins of the present invention and the nucleic acid segments which encode them and still obtain a functional molecule that encodes a protein with desirable properties. The following is a discussion based upon changing the amino acid sequence of a protein to create an equivalent, or possibly an improved, second-generation molecule. The amino acid changes can be achieved by changing the codons of the nucleic acid sequence, according to the codons given in Table 10.

Table 10: Codon degeneracies of amino acids

Amino acid	One letter	Three letter	Codons
Alanine	A	Ala	GCA GCC GCG GCT
Cysteine	C	Cys	TGC TGT
Aspartic acid	D	Asp	GAC GAT
Glutamic acid	F	Glu	GAA GAG
Phenylalanine	F	Phe	TTC TTT
Glycine	G	Gly	GGA GGC GGG GGT
Histidine	H	His	CAC CAT
Isoleucine	I	Ile	ATA ATC ATT
Lysine	K	Lys	AAA AAG
Leucine	L	Leu	TTA TTG CTA CTC CTG CTT
Methionine	M	Met	ATG
Asparagine	N	Asn	AAC AAT
Proline	P	Pro	CCA CCC CCG CCT
Glutamine	Q	Gln	CAA CAG
Arginine	R	Arg	AGA AGG CGA CGC CGG CGT
Serine	S	Ser	AGC AGT TCA TCC TCG TCT
Threonine	T	Thr	ACA ACC ACG ACT
Valine	V	Val	GTA GTC GTG GTT
Tryptophan	W	Trp	TGG
Tyrosine	Y	Tyr	TAC TAT

1 Certain amino acids can be substituted for other amino acids in a protein sequence
2 without appreciable loss of enzymatic activity. It is thus contemplated that various
3 changes can be made in the peptide sequences of the disclosed protein sequences, or their
4 corresponding nucleic acid sequences without appreciable loss of the biological activity.

5 In making such changes, the hydropathic index of amino acids can be considered.
6 The importance of the hydropathic amino acid index in conferring interactive biological
7 function on a protein is generally understood in the art (Kyte and Doolittle, *J. Mol. Biol.*,
8 157: 105-132, 1982). It is accepted that the relative hydropathic character of the amino
9 acid contributes to the secondary structure of the resultant protein, which in turn defines
10 the interaction of the protein with other molecules, for example, enzymes, substrates,
11 receptors, DNA, antibodies, antigens, and the like.

12 Each amino acid has been assigned a hydropathic index on the basis of their
13 hydrophobicity and charge characteristics. These are: isoleucine (+4.5); valine (+4.2);
14 leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine
15 (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3);
16 proline (-1.6); histidine (-3.2); glutamate/glutamine/aspartate/asparagine (-3.5); lysine (-
17 3.9); and arginine (-4.5).

18 It is known in the art that certain amino acids can be substituted by other amino
19 acids having a similar hydropathic index or score and still result in a protein with similar
20 biological activity, i.e., still obtain a biologically functional protein. In making such
21 changes, the substitution of amino acids whose hydropathic indices are within ± 2 is
22 preferred, those within ± 1 are more preferred, and those within ± 0.5 are most preferred.

23 It is also understood in the art that the substitution of like amino acids can be
24 made effectively on the basis of hydrophilicity. U.S. Patent No. 4,554,101 (Hopp, T.P.,
25 issued November 19, 1985) states that the greatest local average hydrophilicity of a
26 protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a
27 biological property of the protein. The following hydrophilicity values have been
28 assigned to amino acids: arginine/lysine (+3.0); aspartate/glutamate (+3.0 ± 1); serine
29 (+0.3); asparagine/glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 ± 1);
30 alanine/histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5);
31 leucine/isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); and tryptophan (-3.4).

1 It is understood that an amino acid can be substituted by another amino acid
2 having a similar hydrophilicity score and still result in a protein with similar biological
3 activity, i.e., still obtain a biologically functional protein. In making such changes, the
4 substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those
5 within ± 1 are more preferred, and those within ± 0.5 are most preferred.

6 As outlined above, amino acid substitutions are therefore based on the relative
7 similarity of the amino acid side-chain substituents, for example, their hydrophobicity,
8 hydrophilicity, charge, size, and the like. Exemplary substitutions which take various of
9 the foregoing characteristics into consideration are well known to those of skill in the art
10 and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine
11 and asparagine; and valine, leucine, and isoleucine. Changes which are not expected to
12 be advantageous may also be used if these resulted in functional patatin proteins.

13 Example 23: Production of patatin proteins and permuteins in plants

14 Plant Vectors

15 In plants, transformation vectors capable of introducing nucleic acid sequences
16 encoding patatin proteins and permuteins are easily designed, and generally contain one
17 or more nucleic acid coding sequences of interest under the transcriptional control of 5'
18 and 3' regulatory sequences. Such vectors generally comprise, operatively linked in
19 sequence in the 5' to 3' direction, a promoter sequence that directs the transcription of a
20 downstream heterologous structural nucleic acid sequence in a plant; optionally, a 5' non-
21 translated leader sequence; a nucleic acid sequence that encodes a protein of interest; and
22 a 3' non-translated region that encodes a polyadenylation signal which functions in plant
23 cells to cause the termination of transcription and the addition of polyadenylate
24 nucleotides to the 3' end of the mRNA encoding the protein. Plant transformation
25 vectors also generally contain a selectable marker. Typical 5'-3' regulatory sequences
26 include a transcription initiation start site, a ribosome binding site, an RNA processing
27 signal, a transcription termination site, and/or a polyadenylation signal. Vectors for plant
28 transformation have been reviewed in Rodriguez et al. (Vectors: A Survey of Molecular
29 Cloning Vectors and Their Uses, Butterworths, Boston., 1988), Glick et al. (Methods in
30 Plant Molecular Biology and Biotechnology, CRC Press, Boca Raton, Fla., 1993), and

1 Croy (Plant Molecular Biology Labfax, Hames and Rickwood (Eds.), BIOS Scientific
2 Publishers Limited, Oxford, UK., 1993).

3 Plant Promoters

4 Plant promoter sequences can be constitutive or inducible, environmentally- or
5 developmentally-regulated, or cell- or tissue-specific. Often-used constitutive promoters
6 include the CaMV 35S promoter (Odell, J.T. et al., *Nature* 313: 810-812, 1985), the
7 enhanced CaMV 35S promoter, the Figwort Mosaic Virus (FMV) promoter (Richins et
8 al., *Nucleic Acids Res.* 20: 8451-8466, 1987), the mannopine synthase (*mas*) promoter,
9 the nopaline synthase (*nos*) promoter, and the octopine synthase (*ocs*) promoter. Useful
10 inducible promoters include promoters induced by salicylic acid or polyacrylic acids (PR-
11 1, Williams, S. W. et al, *Biotechnology* 10: 540-543, 1992), induced by application of
12 safeners (substituted benzenesulfonamide herbicides, Hershey, H.P. and Stoner, T.D.,
13 *Plant Mol. Biol.* 17: 679-690, 1991), heat-shock promoters (Ou-Lee et al., *Proc. Natl.*
14 *Acad. Sci. U.S.A.* 83: 6815-6819, 1986; Ainley et al., *Plant Mol. Biol.* 14: 949-967,
15 1990), a nitrate-inducible promoter derived from the spinach nitrite reductase gene (Back
16 et al., *Plant Mol. Biol.* 17: 9-18, 1991), hormone-inducible promoters (Yamaguchi-
17 Shinozaki, K. et al., *Plant Mol. Biol.* 15: 905-912, 1990; Kares et al., *Plant Mol. Biol.* 15:
18 225-236, 1990), and light-inducible promoters associated with the small subunit of RuBP
19 carboxylase and LHCP gene families (Kuhlemeier et al., *Plant Cell* 1: 471, 1989;
20 Feinbaum, R.L. et al., *Mol. Gen. Genet.* 226: 449-456, 1991; Weisshaar, B. et al., *EMBO*
21 *J.* 10: 1777-1786, 1991; Lam, E. and Chua, N.H., *J. Biol. Chem.* 266: 17131-17135,
22 1990; Castresana, C. et al., *EMBO J.* 7: 1929-1936, 1988; Schulze-Lefert et al., *EMBO J.*
23 8: 651, 1989). Examples of useful tissue-specific, developmentally-regulated promoters
24 include the β -conglycinin 7S promoter (Doyle, J.J. et al., *J. Biol. Chem.* 261: 9228-9238,
25 1986; Slighton and Beachy, *Planta* 172: 356-363, 1987), and seed-specific promoters
26 (Knutzon, D.S. et al., *Proc. Natl. Acad. Sci. U.S.A.* 89: 2624-2628, 1992; Bustos, M.M. et
27 al., *EMBO J.* 10: 1469-1479, 1991; Lam and Chua, *Science* 248: 471, 1991; Stayton et
28 al., *Aust. J. Plant. Physiol.* 18: 507, 1991). Plant functional promoters useful for
29 preferential expression in seed plastids include those from plant storage protein genes and
30 from genes involved in fatty acid biosynthesis in oilseeds. Examples of such promoters

1 include the 5' regulatory regions from such genes as napin (Kridl et al., *Seed Sci. Res.* 1:
2 209-219, 1991), phaseolin, zein, soybean trypsin inhibitor, ACP, stearyl-ACP
3 desaturase, and oleosin. Seed-specific gene regulation is discussed in EP 0 255 378.
4 Promoter hybrids can also be constructed to enhance transcriptional activity (Comai, L.
5 and Moran, P.M., U.S. Patent No. 5,106,739, issued April 21, 1992), or to combine
6 desired transcriptional activity and tissue specificity.

7 Plant transformation and regeneration

8 A variety of different methods can be employed to introduce such vectors into
9 plant protoplasts, cells, callus tissue, leaf discs, meristems, etcetera, to generate
10 transgenic plants, including *Agrobacterium*-mediated transformation, particle gun
11 delivery, microinjection, electroporation, polyethylene glycol mediated protoplast
12 transformation, liposome-mediated transformation, etcetera (reviewed in Potrykus, I.
13 *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 42: 205-225, 1991). In general, transgenic
14 plants comprising cells containing and expressing DNAs encoding patatin proteins and
15 permuteins can be produced by transforming plant cells with a DNA construct as
16 described above via any of the foregoing methods; selecting plant cells that have been
17 transformed on a selective medium; regenerating plant cells that have been transformed
18 to produce differentiated plants; and selecting a transformed plant which expresses the
19 protein-encoding nucleotide sequence.

20 Specific methods for transforming a wide variety of dicots and obtaining
21 transgenic plants are well documented in the literature (Gasser and Fraley, *Science* 244:
22 1293-1299, 1989; Fisk and Dandekar, *Scientia Horticulturae* 55: 5-36, 1993; Christou,
23 *Agro Food Industry Hi Tech*, p.17, 1994; and the references cited therein).

24 Successful transformation and plant regeneration have been reported in the
25 monocots as follows: asparagus (*Asparagus officinalis*; Bytebier et al., *Proc. Natl. Acad.*
26 *Sci. U.S.A.* 84: 5345-5349, 1987); barley (*Hordeum vulgare*; Wan and Lemaux, *Plant*
27 *Physiol.* 104: 37-48, 1994); maize (*Zea mays*; Rhodes, C.A. et al., *Science* 240: 204-207,
28 1988; Gordon-Kamm et al., *Plant Cell* 2: 603-618, 1990; Fromm, M.E. et al.,
29 *Bio/Technology* 8: 833-839, 1990; Koziel et al., *Bio/Technology* 11: 194-200, 1993); oats
30 (*Avena sativa*; Somers et al., *Bio/Technology* 10: 1589-1594, 1992); orchardgrass

1 (*Dactylis glomerata*; Horn et al., *Plant Cell Rep.* 7: 469-472, 1988); rice (*Oryza sativa*,
2 including indica and japonica varieties; Toriyama et al., *Bio/Technology* 6: 10, 1988;
3 Zhang et al., *Plant Cell Rep.* 7: 379-384, 1988; Luo and Wu, *Plant Mol. Biol. Rep.* 6:
4 165-174, 1988; Zhang and Wu, *Theor. Appl. Genet.* 76: 835-840, 1988; Christou et al.,
5 *Bio/Technology* 9: 957-962, 1991); rye (*Secale cereale*; De la Pena et al., *Nature* 325:
6 274-276, 1987); sorghum (*Sorghum bicolor*; Casas, A.M. et al., *Proc. Natl. Acad. Sci.*
7 *U.S.A.* 90: 11212-11216, 1993); sugar cane (*Saccharum* spp.; Bower and Birch, *Plant J.*
8 2: 409-416, 1992); tall fescue (*Festuca arundinacea*; Wang, Z.Y. et al., *Bio/Technology*
9 10: 691-696, 1992); turfgrass (*Agrostis palustris*; Zhong et al., *Plant Cell Rep.* 13: 1-6,
10 1993); wheat (*Triticum aestivum*; Vasil et al., *Bio/Technology* 10: 667-674, 1992; Weeks,
11 T. et al., *Plant Physiol.* 102: 1077-1084, 1993; Becker et al., *Plant J.* 5: 299-307, 1994),
12 and alfalfa (Masoud, S.A. et al., *Transgen. Res.* 5: 313, 1996); *Brassica* (canola/oilseed
13 rape) (Fry, J. *Plant Cell Rep.* 6: 321-325, 1987); and soybean (Hinchee, M. *Bio/Technol.*
14 6: 915-922, 1988).

15 All of the compositions and/or methods disclosed and claimed herein can be made
16 and executed without undue experimentation in light of the present disclosure. While the
17 compositions and methods of this invention have been described in terms of preferred
18 embodiments, it will be apparent to those of skill in the art that variations can be applied
19 to the compositions and/or methods and in the steps or in the sequence of steps of the
20 methods described herein without departing from the concept, spirit and scope of the
21 invention. More specifically, it will be apparent that certain agents which are both
22 chemically and physiologically related can be substituted for the agents described herein
23 while the same or similar results would be achieved. All such similar substitutes and
24 modifications apparent to those skilled in the art are deemed to be within the spirit, scope
25 and concept of the invention.
26
27

Dup.

SEQUENCE LISTING

1
2
3 <110> ALIBHAI, MURTAZA F.
4 ASTWOOD, JAMES D.
5 SAMPSON, HUGH A.
6 McWHERTER, CHARLES F.
7
8 <120> PREPARATION OF DEALLERGENIZED PATATIN PROTEINS AND PERMUTEINS
9
10 <130> MOBT217
11
12 <160> 293
13
14 <170> PatentIn version 3.0
15
16 <210> 1
17 <211> 1158
18 <212> DNA
19 <213> Solanum Tuberosum
20
21 <400> 1
22 atggcaacta ctaaatcttt tttaatTTta atatttatga tattagcaac tactagtTca 60
23
24 acatttgctc agttgggaga aatggTgact gttcttagta ttgatggagg tggaattaga 120
25
26 gggatcatc cggctaccat tctcgaattt ctTgaaggac aacttcagga aatggacaat 180
27
28 aatgcagatg caagactTgc agattacttt gatgtaattg gaggaacaag tacaggaggT 240
29
30 ttattgactg ctatgataag tactccaaat gaaaacaatc gaccctTtgc Tgctgccaaa 300
31
32 gaaattgtac ctttttactt cgaacatggc cctcagattt ttaatcctag Tggtcaaatt 360
33
34 ttaggcccaa aatatgatgg aaaatatctt atgcaagTtc tTcaagaaaa actTggagaa 420
35
36 actcgtgtgc atcaagcttt gacagaagtt gTcatctcaa gctTtgacat caaaacaaat 480
37
38 aagccagtaa tattcactaa gtcaaattta gcaaactctc cagaattgga Tgctaagatg 540
39
40 tatgacataa gttattccac agcagcagct ccaacatatt tTcctccgca tTactttgtt 600
41
42 actaatacta gtaatggaga tgaatatgag tTcaatcttg ttgatggTgc Tgttgctact 660
43
44 gttgctgato cggcgTtatt atccattagc gTtgcaacga gactTgcaca aaaggatcca 720
45
46 gcatttgctt caattaggTc attgaattac aaaaaaatgc Tgttgctctc attaggcact 780
47
48 ggcactactt cagagTttga taaaacatat acagcaaaag aggcagctac ctggactgct 840
49
50 gtacattgga Tgttagttat acagaaaatg actgatgcag caagtTctta catgactgat 900
51
52 tattaccctt ctactgcttt tcaagctctt gattcaaaaa acaattacct cagggtTcaa 960
53
54 gaaaatgcat taacaggcac aactactgaa atggatgatg cttctgaggc taatatggaa 1020
55
56 ttattagtac aagTtggtga aaacttattg aagaaccag tTccgaaga caatcctgaa 1080

1 acctatgagg aagctctaaa gaggtttgca aaattgctct ctgataggaa gaaactccga 1140
 2
 3
 4 gcaaacaag cttcttat 1158
 5
 6 <210> 2
 7 <211> 386
 8 <212> PRT
 9 <213> Solanum Tuberosum
 10
 11 <400> 2
 12
 13 Met Ala Thr Thr Lys Ser Phe Leu Ile Leu Ile Phe Met Ile Leu Ala
 14 1 5 10 15
 15
 16 Thr Thr Ser Ser Thr Phe Ala Gln Leu Gly Glu Met Val Thr Val Leu
 17 20 25 30
 18
 19 Ser Ile Asp Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala Thr Ile Leu
 20 35 40 45
 21
 22 Glu Phe Leu Glu Gly Gln Leu Gln Glu Met Asp Asn Ala Asp Ala
 23 50 55 60
 24
 25 Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly
 26 65 70 75 80
 27
 28 Leu Leu Thr Ala Met Ile Ser Thr Pro Asn Glu Asn Asn Arg Pro Phe
 29 85 90 95
 30
 31 Ala Ala Ala Lys Glu Ile Val Pro Phe Tyr Phe Glu His Gly Pro Gln
 32 100 105 110
 33
 34 Ile Phe Asn Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr Asp Gly Lys
 35 115 120 125
 36
 37 Tyr Leu Met Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His
 38 130 135 140
 39
 40 Gln Ala Leu Thr Glu Val Val Ile Ser Ser Phe Asp Ile Lys Thr Asn
 41 145 150 155 160
 42
 43 Lys Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Asn Ser Pro Glu Leu
 44 165 170 175
 45
 46 Asp Ala Lys Met Tyr Asp Ile Ser Tyr Ser Thr Ala Ala Ala Pro Thr
 47 180 185 190
 48
 49 Tyr Phe Pro Pro His Tyr Phe Val Thr Asn Thr Ser Asn Gly Asp Glu
 50 195 200 205
 51
 52 Tyr Glu Phe Asn Leu Val Asp Gly Ala Val Ala Thr Val Ala Asp Pro
 53 210 215 220
 54
 55 Ala Leu Leu Ser Ile Ser Val Ala Thr Arg Leu Ala Gln Lys Asp Pro
 56 225 230 235 240
 57

1 Ala Phe Ala Ser Ile Arg Ser Leu Asn Tyr Lys Lys Met Leu Leu Leu
 2 245 250 255
 3
 4 Ser Leu Gly Thr Gly Thr Thr Ser Glu Phe Asp Lys Thr Tyr Thr Ala
 5 260 265 270
 6
 7 Lys Glu Ala Ala Thr Trp Thr Ala Val His Trp Met Leu Val Ile Gln
 8 275 280 285
 9
 10 Lys Met Thr Asp Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Leu Ser
 11 290 295 300
 12
 13 Thr Ala Phe Gln Ala Leu Asp Ser Lys Asn Asn Tyr Leu Arg Val Gln
 14 305 310 315 320
 15
 16 Glu Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu
 17 325 330 335
 18
 19 Ala Asn Met Glu Leu Leu Val Gln Val Gly Glu Asn Leu Leu Lys Lys
 20 340 345 350
 21
 22 Pro Val Ser Glu Asp Asn Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg
 23 355 360 365
 24
 25 Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala
 26 370 375 380
 27
 28 Ser Tyr
 29 385
 30
 31 <210> 3
 32 <211> 54
 33 <212> DNA
 34 <213> Synthetic
 35
 36 <400> 3
 37 ggagctcgag aaaagagagg ctgaagctca gttgggagaa atggtgactg ttct 54
 38
 39 <210> 4
 40 <211> 29
 41 <212> DNA
 42 <213> Synthetic
 43
 44 <400> 4
 45 ggtctagagg aattctcatt_aataagaag 29
 46
 47 <210> 5
 48 <211> 1138
 49 <212> DNA
 50 <213> Synthetic
 51
 52 <400> 5
 53 ggagctcgag aaaagagagg ctgaagctca gttgggagaa atggtgactg ttcttagtat 60
 54
 55 tgatggagggt ggaattagag ggatcattcc ggctaccatt ctgaatttc ttgaaggaca 120
 56
 57 acttcaggaa atggacaata atgcagatgc aagacttgca gattactttg atgtaattgg 180

1
 2 aggaacaagt acaggagggtt tattgactgc tatgataagt actccaaatg aaaacaatcg 240
 3
 4 acccttttgc gctgccaaag aaattgtacc tttttacttc gaacatggcc ctccagatttt 300
 5
 6 taatcctagt ggtcaaattt taggccccaa atatgatgga aaatatctta tgcaagttct 360
 7
 8 tcaagaaaaa cttggagaaa ctggtgtgca tcaagctttg acagaagttg tcatctcaag 420
 9
 10 ctttgacatc aaaacaaata agccagtaat attcactaag tcaaatttag caaactctcc 480
 11
 12 agaattggat gctaaagatg atgacataag ttattccaca gcagcagctc caacatatct 540
 13
 14 tctctccgat tactttgtta ctaatactag taatggagat gaatatgagt tcaatcttgt 600
 15
 16 tgatggtgct gttgctactg ttgctgatcc ggcgttatta tccattagcg ttgcaacgag 660
 17
 18 acttgcacaa aaggatccag catttgcttc aattagggtc ttgaattaca aaaaaatgct 720
 19
 20 gttgctctca ttaggcactg gcactacttc agagtttgat aaaacatata cagcaaaaga 780
 21
 22 ggcagctacc tggactgctg tacattggat gttagttata cagaaaatga ctgatgcagc 840
 23
 24 aagtctctac atgactgatt attacctttc tactgctttt caagctcttg attcaaaaaa 900
 25
 26 caattacctc aggggttcaag aaaatgcatt aacaggcaca actactgaaa tggatgatgc 960
 27
 28 ttctgaggct aatatggaat tattagtaca agttggtgaa aacttattga agaaaccagt 1020
 29
 30 ttccgaagac aatcctgaaa cctatgagga agctctaag aggtttgcaa aattgctctc 1080
 31
 32 tgataggaag aaactccgag caaacaaga cttctattaa tgagaattcc tctagacc 1138
 33
 34 <210> 6
 35 <211> 452
 36 <212> PRT
 37 <213> Synthetic
 38
 39 <400> 6
 40
 41 Met Arg Phe Pro Ser Ile Phe Thr Ala Val Leu Phe Ala Ala Ser Ser
 42 1 5 10 15
 43
 44 Ala Leu Ala Ala Pro Val Asn Thr Thr Thr Glu Asp Glu Thr Ala Gln
 45 20 25 30
 46
 47 Ile Pro Ala Glu Ala Val Ile Gly Tyr Ser Asp Leu Glu Gly Asp Phe
 48 35 40 45
 49
 50 Asp Val Ala Val Leu Pro Phe Ser Asn Ser Thr Asn Asn Gly Leu Leu
 51 50 55 60
 52
 53 Phe Ile Asn Thr Thr Ile Ala Ser Ile Ala Ala Lys Glu Glu Gly Val
 54 65 70 75 80
 55
 56 Ser Leu Glu Lys Arg Glu Ala Glu Ala Gln Leu Gly Glu Met Val Thr
 57 85 90 95

1
 2 Val Leu Ser Ile Asp Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala Thr
 3 100 105 110
 4
 5 Ile Leu Glu Phe Leu Glu Gly Gln Leu Gln Glu Met Asp Asn Asn Ala
 6 115 120 125
 7
 8 Asp Ala Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr
 9 130 135 140
 10
 11 Gly Gly Leu Leu Thr Ala Met Ile Ser Thr Pro Asn Glu Asn Asn Arg
 12 145 150 155 160
 13
 14 Pro Phe Ala Ala Ala Lys Glu Ile Val Pro Phe Tyr Phe Glu His Gly
 15 165 170 175
 16
 17 Pro Gln Ile Phe Asn Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr Asp
 18 180 185 190
 19
 20 Gly Lys Tyr Leu Met Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg
 21 195 200 205
 22
 23 Val His Gln Ala Leu Thr Glu Val Val Ile Ser Ser Phe Asp Ile Lys
 24 210 215 220
 25
 26 Thr Asn Lys Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Asn Ser Pro
 27 225 230 235 240
 28
 29 Glu Leu Asp Ala Lys Met Tyr Asp Ile Ser Tyr Ser Thr Ala Ala Ala
 30 245 250 255
 31
 32 Pro Thr Tyr Phe Pro Pro His Tyr Phe Val Thr Asn Thr Ser Asn Gly
 33 260 265 270
 34
 35 Asp Glu Tyr Glu Phe Asn Leu Val Asp Gly Ala Val Ala Thr Val Ala
 36 275 280 285
 37
 38 Asp Pro Ala Leu Leu Ser Ile Ser Val Ala Thr Arg Leu Ala Gln Lys
 39 290 295 300
 40
 41 Asp Pro Ala Phe Ala Ser Ile Arg Ser Leu Asn Tyr Lys Lys Met Leu
 42 305 310 315 320
 43
 44 Leu Leu Ser Leu Gly Thr Gly Thr Thr Ser Glu Phe Asp Lys Thr Tyr
 45 325 330 335
 46
 47 Thr Ala Lys Glu Ala Ala Thr Trp Thr Ala Val His Trp Met Leu Val
 48 340 345 350
 49
 50 Ile Gln Lys Met Thr Asp Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr
 51 355 360 365
 52
 53 Leu Ser Thr Ala Phe Gln Ala Leu Asp Ser Lys Asn Asn Tyr Leu Arg
 54 370 375 380
 55
 56 Val Gln Glu Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala
 57 385 390 395 400

1
 2 Ser Glu Ala Asn Met Glu Leu Leu Val Gln Val Gly Glu Asn Leu Leu
 3 405 410 415
 4
 5 Lys Lys Pro Val Ser Glu Asp Asn Pro Glu Thr Tyr Glu Glu Ala Leu
 6 420 425 430
 7
 8 Lys Arg Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn
 9 435 440 445
 10
 11 Lys Ala Ser Tyr
 12 450
 13
 14 <210> 7
 15 <211> 367
 16 <212> PRT
 17 <213> Synthetic
 18
 19 <400> 7
 20
 21 Glu Ala Glu Ala Gln Leu Gly Glu Met Val Thr Val Leu Ser Ile Asp
 22 1 5 10 15
 23
 24 Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala Thr Ile Leu Glu Phe Leu
 25 20 25 30
 26
 27 Glu Gly Gln Leu Gln Glu Met Asp Asn Asn Ala Asp Ala Arg Leu Ala
 28 35 40 45
 29
 30 Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly Leu Leu Thr
 31 50 55 60
 32
 33 Ala Met Ile Ser Thr Pro Asn Glu Asn Asn Arg Pro Phe Ala Ala Ala
 34 65 70 75 80
 35
 36 Lys Glu Ile Val Pro Phe Tyr Phe Glu His Gly Pro Gln Ile Phe Asn
 37 85 90 95
 38
 39 Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr Asp Gly Lys Tyr Leu Met
 40 100 105 110
 41
 42 Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His Gln Ala Leu
 43 115 120 125
 44
 45 Thr Glu Val Val Ile Ser Ser Phe Asp Ile Lys Thr Asn Lys Pro Val
 46 130 135 140
 47
 48 Ile Phe Thr Lys Ser Asn Leu Ala Asn Ser Pro Glu Leu Asp Ala Lys
 49 145 150 155 160
 50
 51 Met Tyr Asp Ile Ser Tyr Ser Thr Ala Ala Ala Pro Thr Tyr Phe Pro
 52 165 170 175
 53
 54 Pro His Tyr Phe Val Thr Asn Thr Ser Asn Gly Asp Glu Tyr Glu Phe
 55 180 185 190
 56
 57 Asn Leu Val Asp Gly Ala Val Ala Thr Val Ala Asp Pro Ala Leu Leu

1 195 200 205
2
3 Ser Ile Ser Val Ala Thr Arg Leu Ala Gln Lys Asp Pro Ala Phe Ala
4 210 215 220
5
6 Ser Ile Arg Ser Leu Asn Tyr Lys Lys Met Leu Leu Ser Leu Gly
7 225 230 235 240
8
9 Thr Gly Thr Thr Ser Glu Phe Asp Lys Thr Tyr Thr Ala Lys Glu Ala
10 245 250 255
11
12 Ala Thr Trp Thr Ala Val His Trp Met Leu Val Ile Gln Lys Met Thr
13 260 265 270
14
15 Asp Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Leu Ser Thr Ala Phe
16 275 280 285
17
18 Gln Ala Leu Asp Ser Lys Asn Asn Tyr Leu Arg Val Gln Glu Asn Ala
19 290 295 300
20
21 Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu Ala Asn Met
22 305 310 315 320
23
24 Glu Leu Leu Val Gln Val Gly Glu Asn Leu Leu Lys Lys Pro Val Ser
25 325 330 335
26
27 Glu Asp Asn Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg Phe Ala Lys
28 340 345 350
29
30 Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala Ser Tyr
31 355 360 365
32
33 <210> 8
34 <211> 24
35 <212> DNA
36 <213> Synthetic
37
38 <400> 8
39 atgttcgaag aaaaaaggta caat 24
40
41 <210> 9
42 <211> 24
43 <212> DNA
44 <213> Synthetic
45
46 <400> 9
47 ttgcataaga aattttccat cata 24
48
49 <210> 10
50 <211> 24
51 <212> DNA
52 <213> Synthetic
53
54 <400> 10
55 tgctgtggaa aaacttatgt cata 24
56
57 <210> 11

```

1  <211> 24
2  <212> DNA
3  <213> Synthetic
4
5  <400> 11
6  cggaggaaaa aatgttgagg ctgc 24
7
8  <210> 12
9  <211> 51
10 <212> DNA
11 <213> Synthetic
12
13 <400> 12
14 atgcggagga aaaaatgttg gagctgctgc tgtggaaaaa cttatgtcat a 51
15
16 <210> 13
17 <211> 24
18 <212> DNA
19 <213> Synthetic
20
21 <400> 13
22 ttttgctgta aatgttttat caaa 24
23
24 <210> 14
25 <211> 24
26 <212> DNA
27 <213> Synthetic
28
29 <400> 14
30 aaccctgagg aaattgtttt ttga 24
31
32 <210> 15
33 <211> 24
34 <212> DNA
35 <213> Synthetic
36
37 <400> 15
38 agcttcctca aaggtttcag gatt 24
39
40 <210> 16
41 <211> 10
42 <212> PRT
43 <213> Synthetic
44
45 <400> 16
46
47 Gln Leu Gly Glu Met Val Thr Val Leu Ser
48 1 5 10
49
50 <210> 17
51 <211> 10
52 <212> PRT
53 <213> Synthetic
54
55 <400> 17
56
57 Met Val Thr Val Leu Ser Ile Asp Gly Gly

```

```

1      1              5              10
2      <210> 18
3      <211> 10
4      <212> PRT
5      <213> Synthetic
6
7      <400> 18
8
9      Leu Ser Ile Asp Gly Gly Gly Ile Arg Gly
10     1              5              10
11
12     <210> 19
13     <211> 10
14     <212> PRT
15     <213> Synthetic
16
17     <400> 19
18
19     Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala
20     1              5              10
21
22     <210> 20
23     <211> 10
24     <212> PRT
25     <213> Synthetic
26
27     <400> 20
28
29     Arg Gly Ile Ile Pro Ala Thr Ile Leu Glu
30     1              5              10
31
32     <210> 21
33     <211> 10
34     <212> PRT
35     <213> Synthetic
36
37     <400> 21
38
39     Pro Ala Thr Ile Leu Glu Phe Leu Glu Gly
40     1              5              10
41
42     <210> 22
43     <211> 10
44     <212> PRT
45     <213> Synthetic
46
47     <400> 22
48
49     Leu Glu Phe Leu Glu Gly Gln Leu Gln Glu
50     1              5              10
51
52     <210> 23
53     <211> 10
54     <212> PRT
55     <213> Synthetic
56
57     <400> 23

```

```

1
2   Glu Gly Gln Leu Gln Glu Met Asp Asn Asn
3   1                      5                      10
4
5   <210> 24
6   <211> 10
7   <212> PRT
8   <213> Synthetic
9
10  <400> 24
11
12  Gln Glu Met Asp Asn Asn Ala Asp Ala Arg
13  1                      5                      10
14
15  <210> 25
16  <211> 10
17  <212> PRT
18  <213> Synthetic
19
20  <400> 25
21
22  Asn Asn Ala Asp Ala Arg Leu Ala Asp Tyr
23  1                      5                      10
24
25  <210> 26
26  <211> 10
27  <212> PRT
28  <213> Synthetic
29
30  <400> 26
31
32  Ala Arg Leu Ala Asp Tyr Phe Asp Val Ile
33  1                      5                      10
34
35  <210> 27
36  <211> 10
37  <212> PRT
38  <213> Synthetic
39
40  <400> 27
41
42  Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser
43  1                      5                      10
44
45  <210> 28
46  <211> 10
47  <212> PRT
48  <213> Synthetic
49
50  <400> 28
51
52  Val Ile Gly Gly Thr Ser Thr Gly Gly Leu
53  1                      5                      10
54
55  <210> 29
56  <211> 10
57  <212> PRT

```



```

1  <213>  Synthetic
2
3  <400>  29
4
5  Thr Ser Thr Gly Gly Leu Leu Thr Ala Met
6  1          5          10
7
8  <210>  30
9  <211>  10
10 <212>  PRT
11 <213>  Synthetic
12
13 <400>  30
14
15 Gly Leu Leu Thr Ala Met Ile Ser Thr Pro
16 1          5          10
17
18 <210>  31
19 <211>  10
20 <212>  PRT
21 <213>  Synthetic
22
23 <400>  31
24
25 Ala Met Ile Ser Thr Pro Asn Glu Asn Asn
26 1          5          10
27
28 <210>  32
29 <211>  10
30 <212>  PRT
31 <213>  Synthetic
32
33 <400>  32
34
35 Thr Pro Asn Glu Asn Asn Arg Pro Phe Ala
36 1          5          10
37
38 <210>  33
39 <211>  10
40 <212>  PRT
41 <213>  Synthetic
42
43 <400>  33
44
45 Asn Asn Arg Pro Phe Ala Ala Ala Lys Glu
46 1          5          10
47
48 <210>  34
49 <211>  10
50 <212>  PRT
51 <213>  Synthetic
52
53 <400>  34
54
55 Phe Ala Ala Ala Lys Glu Ile Val Pro Phe
56 1          5          10
57

```

```

1  <210> 35
2  <211> 10
3  <212> PRT
4  <213> Synthetic
5
6  <400> 35
7
8  Lys Glu Ile Val Pro Phe Tyr Phe Glu His
9  1                      5                      10
10
11 <210> 36
12 <211> 10
13 <212> PRT
14 <213> Synthetic
15
16 <400> 36
17
18 Pro Phe Tyr Phe Glu His Gly Pro Gln Ile
19 1                      5                      10
20
21 <210> 37
22 <211> 10
23 <212> PRT
24 <213> Synthetic
25
26 <400> 37
27
28 Glu His Gly Pro Gln Ile Phe Asn Pro Ser
29 1                      5                      10
30
31 <210> 38
32 <211> 10
33 <212> PRT
34 <213> Synthetic
35
36 <400> 38
37
38 Gln Ile Phe Asn Pro Ser Gly Gln Ile Leu
39 1                      5                      10
40
41 <210> 39
42 <211> 10
43 <212> PRT
44 <213> Synthetic
45
46 <400> 39
47
48 Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr
49 1                      5                      10
50
51 <210> 40
52 <211> 10
53 <212> PRT
54 <213> Synthetic
55
56 <400> 40
57

```

```

1 Ile Leu Gly Pro Lys Tyr Asp Gly Lys Tyr
2 1 5 10
3
4 <210> 41
5 <211> 10
6 <212> PRT
7 <213> Synthetic
8
9 <400> 41
10
11 Lys Tyr Asp Gly Lys Tyr Leu Met Gln Val
12 1 5 10
13
14 <210> 42
15 <211> 10
16 <212> PRT
17 <213> Synthetic
18
19 <400> 42
20
21 Lys Tyr Leu Met Gln Val Leu Gln Glu Lys
22 1 5 10
23
24 <210> 43
25 <211> 10
26 <212> PRT
27 <213> Synthetic
28
29 <400> 43
30
31 Gln Val Leu Gln Glu Lys Leu Gly Glu Thr
32 1 5 10
33
34 <210> 44
35 <211> 10
36 <212> PRT
37 <213> Synthetic
38
39 <400> 44
40
41 Glu Lys Leu Gly Glu Thr Arg Val His Gln
42 1 5 10
43
44 <210> 45
45 <211> 10
46 <212> PRT
47 <213> Synthetic
48
49 <400> 45
50
51 Glu Thr Arg Val His Gln Ala Leu Thr Glu
52 1 5 10
53
54 <210> 46
55 <211> 10
56 <212> PRT
57 <213> Synthetic

```

```

1
2 <400> 46
3
4 His Gln Ala Leu Thr Glu Val Val Ile Ser
5 1 5 10
6
7 <210> 47
8 <211> 10
9 <212> PRT
10 <213> Synthetic
11
12 <400> 47
13
14 Thr Glu Val Val Ile Ser Ser Phe Asp Ile
15 1 5 10
16
17 <210> 48
18 <211> 10
19 <212> PRT
20 <213> Synthetic
21
22 <400> 48
23
24 Ile Ser Ser Phe Asp Ile Lys Thr Asn Lys
25 1 5 10
26
27 <210> 49
28 <211> 10
29 <212> PRT
30 <213> Synthetic
31
32 <400> 49
33
34 Asp Ile Lys Thr Asn Lys Pro Val Ile Phe
35 1 5 10
36
37 <210> 50
38 <211> 10
39 <212> PRT
40 <213> Synthetic
41
42 <400> 50
43
44 Asn Lys Pro Val Ile Phe Thr Lys Ser Asn
45 1 5 10
46
47 <210> 51
48 <211> 10
49 <212> PRT
50 <213> Synthetic
51
52 <400> 51
53
54 Ile Phe Thr Lys Ser Asn Leu Ala Asn Ser
55 1 5 10
56
57 <210> 52

```

```

1 <211> 10
2 <212> PRT
3 <213> Synthetic
4
5 <400> 52
6
7 Ser Asn Leu Ala Asn Ser Pro Glu Leu Asp
8 1 5 10
9
10 <210> 53
11 <211> 10
12 <212> PRT
13 <213> Synthetic
14
15 <400> 53
16
17 Asn Ser Pro Glu Leu Asp Ala Lys Met Tyr
18 1 5 10
19
20 <210> 54
21 <211> 10
22 <212> PRT
23 <213> Synthetic
24
25 <400> 54
26
27 Leu Asp Ala Lys Met Tyr Asp Ile Ser Tyr
28 1 5 10
29
30 <210> 55
31 <211> 10
32 <212> PRT
33 <213> Synthetic
34
35 <400> 55
36
37 Met Tyr Asp Ile Ser Tyr Ser Thr Ala Ala
38 1 5 10
39
40 <210> 56
41 <211> 10
42 <212> PRT
43 <213> Synthetic
44
45 <400> 56
46
47 Ser Tyr Ser Thr Ala Ala Pro Thr Tyr
48 1 5 10
49
50 <210> 57
51 <211> 10
52 <212> PRT
53 <213> Synthetic
54
55 <400> 57
56
57 Ala Ala Ala Pro Thr Tyr Phe Pro Pro His

```

```

1      1          5          10
2
3      <210>  58
4      <211>  10
5      <212>  PRT
6      <213>  Synthetic
7
8      <400>  58
9
10     Thr Tyr Phe Pro Pro His Tyr Phe Val Thr
11     1          5          10
12
13     <210>  59
14     <211>  10
15     <212>  PRT
16     <213>  Synthetic
17
18     <400>  59
19
20     Pro His Tyr Phe Val Thr Asn Thr Ser Asn
21     1          5          10
22
23     <210>  60
24     <211>  10
25     <212>  PRT
26     <213>  Synthetic
27
28     <400>  60
29
30     Val Thr Asn Thr Ser Asn Gly Asp Glu Tyr
31     1          5          10
32
33     <210>  61
34     <211>  10
35     <212>  PRT
36     <213>  Synthetic
37
38     <400>  61
39
40     Ser Asn Gly Asp Glu Tyr Glu Phe Asn Leu
41     1          5          10
42
43     <210>  62
44     <211>  10
45     <212>  PRT
46     <213>  Synthetic
47
48     <400>  62
49
50     Glu Tyr Glu Phe Asn Leu Val Asp Gly Ala
51     1          5          10
52
53     <210>  63
54     <211>  10
55     <212>  PRT
56     <213>  Synthetic
57

```

```

1  <400> 63
2
3  Asn Leu Val Asp Gly Ala Val Ala Thr Val
4  1          5          10
5
6  <210> 64
7  <211> 10
8  <212> PRT
9  <213> Synthetic
10
11 <400> 64
12
13 Gly Ala Val Ala Thr Val Ala Asp Pro Ala
14 1          5          10
15
16 <210> 65
17 <211> 10
18 <212> PRT
19 <213> Synthetic
20
21 <400> 65
22
23 Thr Val Ala Asp Pro Ala Leu Leu Ser Ile
24 1          5          10
25
26 <210> 66
27 <211> 10
28 <212> PRT
29 <213> Synthetic
30
31 <400> 66
32
33 Pro Ala Leu Leu Ser Ile Ser Val Ala Thr
34 1          5          10
35
36 <210> 67
37 <211> 10
38 <212> PRT
39 <213> Synthetic
40
41 <400> 67
42
43 Ser Ile Ser Val Ala Thr Arg Leu Ala Gln
44 1          5          10
45
46 <210> 68
47 <211> 10
48 <212> PRT
49 <213> Synthetic
50
51 <400> 68
52
53 Ala Thr Arg Leu Ala Gln Lys Asp Pro Ala
54 1          5          10
55
56 <210> 69
57 <211> 10

```

```

1  <212>  PRT
2  <213>  Synthetic
3
4  <400>  69
5
6  Ala Gln Lys Asp Pro Ala Phe Ala Ser Ile
7  1          5          10
8
9  <210>  70
10 <211>  10
11 <212>  PRT
12 <213>  Synthetic
13
14 <400>  70
15
16 Pro Ala Phe Ala Ser Ile Arg Ser Leu Asn
17 1          5          10
18
19 <210>  71
20 <211>  10
21 <212>  PRT
22 <213>  Synthetic
23
24 <400>  71
25
26 Ser Ile Arg Ser Leu Asn Tyr Lys Lys Met
27 1          5          10
28
29 <210>  72
30 <211>  10
31 <212>  PRT
32 <213>  Synthetic
33
34 <400>  72
35
36 Leu Asn Tyr Lys Lys Met Leu Leu Leu Ser
37 1          5          10
38
39 <210>  73
40 <211>  10
41 <212>  PRT
42 <213>  Synthetic
43
44 <400>  73
45
46 Lys Met Leu Leu Leu Ser Leu Gly Thr Gly
47 1          5          10
48
49 <210>  74
50 <211>  10
51 <212>  PRT
52 <213>  Synthetic
53
54 <400>  74
55
56 Leu Ser Leu Gly Thr Gly Thr Thr Ser Glu
57 1          5          10

```



```

1
2 <210> 75
3 <211> 10
4 <212> PRT
5 <213> Synthetic
6
7 <400> 75
8
9 Thr Gly Thr Thr Ser Glu Phe Asp Lys Thr
10 1 5 10
11
12 <210> 76
13 <211> 10
14 <212> PRT
15 <213> Synthetic
16
17 <400> 76
18
19 Ser Glu Phe Asp Lys Thr Tyr Thr Ala Lys
20 1 5 10
21
22 <210> 77
23 <211> 10
24 <212> PRT
25 <213> Synthetic
26
27 <400> 77
28
29 Lys Thr Tyr Thr Ala Lys Glu Ala Ala Thr
30 1 5 10
31
32 <210> 78
33 <211> 10
34 <212> PRT
35 <213> Synthetic
36
37 <400> 78
38
39 Ala Lys Glu Ala Ala Thr Trp Thr Ala Val
40 1 5 10
41
42 <210> 79
43 <211> 10
44 <212> PRT
45 <213> Synthetic
46
47 <400> 79
48
49 Ala Thr Trp Thr Ala Val His Trp Met Leu
50 1 5 10
51
52 <210> 80
53 <211> 10
54 <212> PRT
55 <213> Synthetic
56
57 <400> 80

```

```

1
2   Ala Val His Trp Met Leu Val Ile Gln Lys
3   1                      5                      10
4
5   <210> 81
6   <211> 10
7   <212> PRT
8   <213> Synthetic
9
10  <400> 81
11
12  Met Leu Val Ile Gln Lys Met Thr Asp Ala
13  1                      5                      10
14
15  <210> 82
16  <211> 10
17  <212> PRT
18  <213> Synthetic
19
20  <400> 82
21
22  Gln Lys Met Thr Asp Tyr Tyr Leu Ser Thr
23  1                      5                      10
24
25  <210> 83
26  <211> 10
27  <212> PRT
28  <213> Synthetic
29
30  <400> 83
31
32  Asp Ala Ala Ser Ser Tyr Met Thr Asp Tyr
33  1                      5                      10
34
35  <210> 84
36  <211> 10
37  <212> PRT
38  <213> Synthetic
39
40  <400> 84
41
42  Ser Tyr Met Thr Asp Tyr Tyr Leu Ser Thr
43  1                      5                      10
44
45  <210> 85
46  <211> 10
47  <212> PRT
48  <213> Synthetic
49
50  <400> 85
51
52  Asp Tyr Tyr Leu Ser Thr Ala Phe Gln Ala
53  1                      5                      10
54
55  <210> 86
56  <211> 10
57  <212> PRT

```

```

1  <213> Synthetic
2
3  <400> 86
4
5  Ser Thr Ala Phe Gln Ala Leu Asp Ser Lys
6  1             5             10
7
8  <210> 87
9  <211> 10
10 <212> PRT
11 <213> Synthetic
12
13 <400> 87
14
15 Gln Ala Leu Asp Ser Lys Asn Asn Tyr Leu
16 1             5             10
17
18 <210> 88
19 <211> 10
20 <212> PRT
21 <213> Synthetic
22
23 <400> 88
24
25 Ser Lys Asn Asn Tyr Leu Arg Val Gln Glu
26 1             5             10
27
28 <210> 89
29 <211> 10
30 <212> PRT
31 <213> Synthetic
32
33 <400> 89
34
35 Tyr Leu Arg Val Gln Glu Asn Ala Leu Thr
36 1             5             10
37
38 <210> 90
39 <211> 10
40 <212> PRT
41 <213> Synthetic
42
43 <400> 90
44
45 Gln Glu Asn Ala Leu Thr Gly Thr Thr Thr
46 1             5             10
47
48 <210> 91
49 <211> 10
50 <212> PRT
51 <213> Synthetic
52
53 <400> 91
54
55 Leu Thr Gly Thr Thr Thr Glu Met Asp Asp
56 1             5             10
57

```

```

1  <210> 92
2  <211> 10
3  <212> PRT
4  <213> Synthetic
5
6  <400> 92
7
8  Thr Thr Glu Met Asp Asp Ala Ser Glu Ala
9  1          5          10
10
11 <210> 93
12 <211> 10
13 <212> PRT
14 <213> Synthetic
15
16 <400> 93
17
18 Asp Asp Ala Ser Glu Ala Asn Met Glu Leu
19 1          5          10
20
21 <210> 94
22 <211> 10
23 <212> PRT
24 <213> Synthetic
25
26 <400> 94
27
28 Glu Ala Asn Met Glu Leu Leu Val Gln Val
29 1          5          10
30
31 <210> 95
32 <211> 10
33 <212> PRT
34 <213> Synthetic
35
36 <400> 95
37
38 Glu Leu Leu Val Gln Val Gly Glu Asn Leu
39 1          5          10
40
41 <210> 96
42 <211> 10
43 <212> PRT
44 <213> Synthetic
45
46 <400> 96
47
48 Gln Val Gly Glu Asn Leu Leu Lys Lys Pro
49 1          5          10
50
51 <210> 97
52 <211> 10
53 <212> PRT
54 <213> Synthetic
55
56 <400> 97
57

```

```

1  Asn Leu Leu Lys Lys Pro Val Ser Glu Asp
2  1                      5                      10
3
4  <210> 98
5  <211> 10
6  <212> PRT
7  <213> Synthetic
8
9  <400> 98
10
11 Lys Pro Val Ser Glu Asp Asn Pro Glu Thr
12 1                      5                      10
13
14 <210> 99
15 <211> 10
16 <212> PRT
17 <213> Synthetic
18
19 <400> 99
20
21 Glu Asp Asn Pro Glu Thr Tyr Glu Glu Ala
22 1                      5                      10
23
24 <210> 100
25 <211> 10
26 <212> PRT
27 <213> Synthetic
28
29 <400> 100
30
31 Glu Thr Tyr Glu Glu Ala Leu Lys Arg Phe
32 1                      5                      10
33
34 <210> 101
35 <211> 10
36 <212> PRT
37 <213> Synthetic
38
39 <400> 101
40
41 Glu Ala Leu Lys Arg Phe Ala Lys Leu Leu
42 1                      5                      10
43
44 <210> 102
45 <211> 10
46 <212> PRT
47 <213> Synthetic
48
49 <400> 102
50
51 Arg Phe Ala Lys Leu Leu Ser Asp Arg Lys
52 1                      5                      10
53
54 <210> 103
55 <211> 10
56 <212> PRT
57 <213> Synthetic

```

```

1
2 <400> 103
3
4 Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala
5 1 5 10
6
7 <210> 104
8 <211> 10
9 <212> PRT
10 <213> Synthetic
11
12 <400> 104
13
14 Arg Lys Lys Leu Arg Ala Asn Lys Ala Ser
15 1 5 10
16
17 <210> 105
18 <211> 10
19 <212> PRT
20 <213> Synthetic
21
22 <400> 105
23
24 Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser
25 1 5 10
26
27 <210> 106
28 <211> 10
29 <212> PRT
30 <213> Synthetic
31
32 <400> 106
33
34 Asp Tyr Phe Asp Val Ile Ala Gly Thr Ser
35 1 5 10
36
37 <210> 107
38 <211> 10
39 <212> PRT
40 <213> Synthetic
41
42 <400> 107
43
44 Val Ile Gly Gly Thr Ser Thr Gly Gly Leu
45 1 5 10
46
47 <210> 108
48 <211> 10
49 <212> PRT
50 <213> Synthetic
51
52 <400> 108
53
54 Val Ile Ala Gly Thr Ser Thr Gly Ala Leu
55 1 5 10
56
57 <210> 109

```

```

1  <211> 10
2  <212> PRT
3  <213> Synthetic
4
5  <400> 109
6
7  Ala Phe Tyr Phe Glu His Gly Pro Gln Ile
8  1             5             10
9
10 <210> 110
11 <211> 10
12 <212> PRT
13 <213> Synthetic
14
15 <400> 110
16
17 Pro Ala Tyr Phe Glu His Gly Pro Gln Ile
18 1             5             10
19
20 <210> 111
21 <211> 10
22 <212> PRT
23 <213> Synthetic
24
25 <400> 111
26
27 Pro Phe Ala Phe Glu His Gly Pro Gln Ile
28 1             5             10
29
30 <210> 112
31 <211> 10
32 <212> PRT
33 <213> Synthetic
34
35 <400> 112
36
37 Pro Phe Tyr Ala Glu His Gly Pro Gln Ile
38 1             5             10
39
40 <210> 113
41 <211> 10
42 <212> PRT
43 <213> Synthetic
44
45 <400> 113
46
47 Pro Phe Tyr Phe Ala His Gly Pro Gln Ile
48 1             5             10
49
50 <210> 114
51 <211> 10
52 <212> PRT
53 <213> Synthetic
54
55 <400> 114
56
57 Pro Phe Tyr Phe Glu Ala Gly Pro Gln Ile

```

```

1      1              5              10
2
3      <210> 115
4      <211> 10
5      <212> PRT
6      <213> Synthetic
7
8      <400> 115
9
10     Pro Phe Tyr Phe Glu His Ala Pro Gln Ile
11     1              5              10
12
13     <210> 116
14     <211> 10
15     <212> PRT
16     <213> Synthetic
17
18     <400> 116
19
20     Pro Phe Tyr Phe Glu His Gly Ala Gln Ile
21     1              5              10
22
23     <210> 117
24     <211> 10
25     <212> PRT
26     <213> Synthetic
27
28     <400> 117
29
30     Pro Phe Tyr Phe Glu His Gly Pro Ala Ile
31     1              5              10
32
33     <210> 118
34     <211> 10
35     <212> PRT
36     <213> Synthetic
37
38     <400> 118
39
40     Pro Phe Tyr Phe Glu His Gly Pro Gln Ala
41     1              5              10
42
43     <210> 119
44     <211> 10
45     <212> PRT
46     <213> Synthetic
47
48     <400> 119
49
50     Thr Phe Tyr Leu Glu Asn Gly Pro Lys Ile
51     1              5              10
52
53     <210> 120
54     <211> 10
55     <212> PRT
56     <213> Synthetic
57

```



```

1  <400> 120
2
3  Pro Phe Phe Phe Glu His Gly Pro Gln Ile
4  1          5          10
5
6  <210> 121
7  <211> 10
8  <212> PRT
9  <213> Synthetic
10
11 <400> 121
12
13 Ala Tyr Leu Met Gln Val Leu Gln Glu Lys
14 1          5          10
15
16 <210> 122
17 <211> 10
18 <212> PRT
19 <213> Synthetic
20
21 <400> 122
22
23 Lys Ala Leu Met Gln Val Leu Gln Glu Lys
24 1          5          10
25
26 <210> 123
27 <211> 10
28 <212> PRT
29 <213> Synthetic
30
31 <400> 123
32
33 Lys Tyr Ala Met Gln Val Leu Gln Glu Lys
34 1          5          10
35
36 <210> 124
37 <211> 10
38 <212> PRT
39 <213> Synthetic
40
41 <400> 124
42
43 Lys Tyr Leu Ala Gln Val Leu Gln Glu Lys
44 1          5          10
45
46 <210> 125
47 <211> 10
48 <212> PRT
49 <213> Synthetic
50
51 <400> 125
52
53 Lys Tyr Leu Met Ala Val Leu Gln Glu Lys
54 1          5          10
55
56 <210> 126
57 <211> 10

```

```

1  <212>  PRT
2  <213>  Synthetic
3
4  <400>  126
5
6  Lys Tyr Leu Met Gln Ala Leu Gln Glu Lys
7  1                      5                      10
8
9  <210>  127
10 <211>  10
11 <212>  PRT
12 <213>  Synthetic
13
14 <400>  127
15
16 Lys Tyr Leu Met Gln Val Ala Gln Glu Lys
17 1                      5                      10
18
19 <210>  128
20 <211>  10
21 <212>  PRT
22 <213>  Synthetic
23
24 <400>  128
25
26 Lys Tyr Leu Met Gln Val Leu Ala Glu Lys
27 1                      5                      10
28
29 <210>  129
30 <211>  10
31 <212>  PRT
32 <213>  Synthetic
33
34 <400>  129
35
36 Lys Tyr Leu Met Gln Val Leu Gln Ala Lys
37 1                      5                      10
38
39 <210>  130
40 <211>  10
41 <212>  PRT
42 <213>  Synthetic
43
44 <400>  130
45
46 Lys Tyr Leu Met Gln Val Leu Gln Glu Ala
47 1                      5                      10
48
49 <210>  131
50 <211>  10
51 <212>  PRT
52 <213>  Synthetic
53
54 <400>  131
55
56 Val Phe Leu His Asp Lys Ile Lys Ser Leu
57 1                      5                      10

```

```

1
2 <210> 132
3 <211> 10
4 <212> PRT
5 <213> Synthetic
6
7 <400> 132
8
9 Ala Tyr Ser Thr Ala Ala Ala Pro Thr Tyr
10 1 5 10
11
12 <210> 133
13 <211> 10
14 <212> PRT
15 <213> Synthetic
16
17 <400> 133
18
19 Ser Ala Ser Thr Ala Ala Ala Pro Thr Tyr
20 1 5 10
21
22 <210> 134
23 <211> 10
24 <212> PRT
25 <213> Synthetic
26
27 <400> 134
28
29 Ser Tyr Ala Thr Ala Ala Ala Pro Thr Tyr
30 1 5 10
31
32 <210> 135
33 <211> 10
34 <212> PRT
35 <213> Synthetic
36
37 <400> 135
38
39 Ser Tyr Ser Ala Ala Ala Ala Pro Thr Tyr
40 1 5 10
41
42 <210> 136
43 <211> 10
44 <212> PRT
45 <213> Synthetic
46
47 <400> 136
48
49 Ser Tyr Ser Thr Ala Ala Ala Ala Thr Tyr
50 1 5 10
51
52 <210> 137
53 <211> 10
54 <212> PRT
55 <213> Synthetic
56
57 <400> 137

```

```

1
2   Ser Tyr Ser Thr Ala Ala Ala Pro Ala Tyr
3   1           5                      10
4
5   <210> 138
6   <211> 10
7   <212> PRT
8   <213> Synthetic
9
10  <400> 138
11
12  Ser Tyr Ser Thr Ala Ala Ala Pro Thr Ala
13  1           5                      10
14
15  <210> 139
16  <211> 10
17  <212> PRT
18  <213> Synthetic
19
20  <400> 139
21
22  Cys Ile Ser Thr Ser Ala Ala Pro Thr Tyr
23  1           5                      10
24
25  <210> 140
26  <211> 10
27  <212> PRT
28  <213> Synthetic
29
30  <400> 140
31
32  Ser Tyr Ser Thr Ala Ala Ala Pro Ala Phe
33  1           5                      10
34
35  <210> 141
36  <211> 10
37  <212> PRT
38  <213> Synthetic
39
40  <400> 141
41
42  Ala Phe Ala Ala Ala Ala Ala Pro Thr Tyr
43  1           5                      10
44
45  <210> 142
46  <211> 10
47  <212> PRT
48  <213> Synthetic
49
50  <400> 142
51
52  Ser Tyr Ser Thr Ala Ala Ala Pro Thr Phe
53  1           5                      10
54
55  <210> 143
56  <211> 10
57  <212> PRT

```

```

1  <213> Synthetic
2
3  <400> 143
4
5  Ser Thr Ser Ala Ala Pro Thr Tyr Phe Pro
6  1          5          10
7
8  <210> 144
9  <211> 10
10 <212> PRT
11 <213> Synthetic
12
13 <400> 144
14
15 Ser Thr Ser Ala Ala Pro Thr Phe Phe Pro
16 1          5          10
17
18 <210> 145
19 <211> 10
20 <212> PRT
21 <213> Synthetic
22
23 <400> 145
24
25 Ser Thr Ser Ala Ala Pro Thr Ala Phe Pro
26 1          5          10
27
28 <210> 146
29 <211> 10
30 <212> PRT
31 <213> Synthetic
32
33 <400> 146
34
35 Ser Thr Ala Ala Ala Pro Thr Phe Phe Pro
36 1          5          10
37
38 <210> 147
39 <211> 10
40 <212> PRT
41 <213> Synthetic
42
43 <400> 147
44
45 Ala Ala Ala Ala Thr Tyr Phe Pro Pro His
46 1          5          10
47
48 <210> 148
49 <211> 10
50 <212> PRT
51 <213> Synthetic
52
53 <400> 148
54
55 Ala Ala Ala Pro Ala Tyr Phe Pro Pro His
56 1          5          10
57

```

```

1  <210> 149
2  <211> 10
3  <212> PRT
4  <213> Synthetic
5
6  <400> 149
7
8  Ala Ala Ala Pro Thr Ala Phe Pro Pro His
9  1                               5               10
10
11 <210> 150
12 <211> 10
13 <212> PRT
14 <213> Synthetic
15
16 <400> 150
17
18 Ala Ala Ala Pro Thr Tyr Ala Pro Pro His
19 1                               5               10
20
21 <210> 151
22 <211> 10
23 <212> PRT
24 <213> Synthetic
25
26 <400> 151
27
28 Ala Ala Ala Pro Thr Tyr Phe Ala Pro His
29 1                               5               10
30
31 <210> 152
32 <211> 10
33 <212> PRT
34 <213> Synthetic
35
36 <400> 152
37
38 Ala Ala Ala Pro Thr Tyr Phe Pro Ala His
39 1                               5               10
40
41 <210> 153
42 <211> 10
43 <212> PRT
44 <213> Synthetic
45
46 <400> 153
47
48 Ala Ala Ala Pro Thr Tyr Phe Pro Pro Ala
49 1                               5               10
50
51 <210> 154
52 <211> 10
53 <212> PRT
54 <213> Synthetic
55
56 <400> 154
57

```

```

1  Ser Ala Ala Pro Thr Tyr Phe Pro Ala His
2  1                      5                      10
3
4  <210> 155
5  <211> 10
6  <212> PRT
7  <213> Synthetic
8
9  <400> 155
10
11 Ala Ala Ala Pro Ala Phe Phe Pro Pro His
12 1                      5                      10
13
14 <210> 156
15 <211> 10
16 <212> PRT
17 <213> Synthetic
18
19 <400> 156
20
21 Ala Ala Ala Pro Pro Phe Phe Pro Pro His
22 1                      5                      10
23
24 <210> 157
25 <211> 10
26 <212> PRT
27 <213> Synthetic
28
29 <400> 157
30
31 Ala Ala Ala Pro Thr Phe Phe Pro Pro His
32 1                      5                      10
33
34 <210> 158
35 <211> 10
36 <212> PRT
37 <213> Synthetic
38
39 <400> 158
40
41 Ser Ile Ser Val Ala Thr Arg Leu Ala Gln
42 1                      5                      10
43
44 <210> 159
45 <211> 10
46 <212> PRT
47 <213> Synthetic
48
49 <400> 159
50
51 Ala Met Ser Met Leu Thr Lys Glu Val His
52 1                      5                      10
53
54 <210> 160
55 <211> 10
56 <212> PRT
57 <213> Synthetic

```

```

1
2 <400> 160
3
4 Pro Ala Phe Ala Ser Ile Arg Ser Leu Asn
5 1 5 10
6
7 <210> 161
8 <211> 10
9 <212> PRT
10 <213> Synthetic
11
12 <400> 161
13
14 Pro Asn Phe Asn Ala Gly Ser Pro Thr Glu
15 1 5 10
16
17 <210> 162
18 <211> 10
19 <212> PRT
20 <213> Synthetic
21
22 <400> 162
23
24 Lys Met Leu Leu Leu Ser Leu Gly Thr Gly
25 1 5 10
26
27 <210> 163
28 <211> 10
29 <212> PRT
30 <213> Synthetic
31
32 <400> 163
33
34 Asn Tyr Leu Ile Ile Ser Val Gly Thr Gly
35 1 5 10
36
37 <210> 164
38 <211> 10
39 <212> PRT
40 <213> Synthetic
41
42 <400> 164
43
44 Lys Met Leu Leu Leu Ser Leu Gly Ala Gly
45 1 5 10
46
47 <210> 165
48 <211> 10
49 <212> PRT
50 <213> Synthetic
51
52 <400> 165
53
54 Ala Glu Phe Asp Lys Thr Tyr Thr Ala Lys
55 1 5 10
56
57 <210> 166

```



```

1  <211> 10
2  <212> PRT
3  <213> Synthetic
4
5  <400> 166
6
7  Ser Ala Phe Asp Lys Thr Tyr Thr Ala Lys
8  1          5          10
9
10 <210> 167
11 <211> 10
12 <212> PRT
13 <213> Synthetic
14
15 <400> 167
16
17 Ser Glu Ala Asp Lys Thr Tyr Thr Ala Lys
18 1          5          10
19
20 <210> 168
21 <211> 10
22 <212> PRT
23 <213> Synthetic
24
25 <400> 168
26
27 Ser Glu Phe Ala Lys Thr Tyr Thr Ala Lys
28 1          5          10
29
30 <210> 169
31 <211> 10
32 <212> PRT
33 <213> Synthetic
34
35 <400> 169
36
37 Ser Glu Phe Asp Ala Thr Tyr Thr Ala Lys
38 1          5          10
39
40 <210> 170
41 <211> 10
42 <212> PRT
43 <213> Synthetic
44
45 <400> 170
46
47 Ser Glu Phe Asp Lys Ala Tyr Thr Ala Lys
48 1          5          10
49
50 <210> 171
51 <211> 10
52 <212> PRT
53 <213> Synthetic
54
55 <400> 171
56
57 Ser Glu Phe Asp Lys Thr Ala Thr Ala Lys

```

```

1      1              5              10
2
3      <210>  172
4      <211>  10
5      <212>  PRT
6      <213>  Synthetic
7
8      <400>  172
9
10     Ser Glu Phe Asp Lys Thr Tyr Ala Ala Lys
11     1              5              10
12
13     <210>  173
14     <211>  10
15     <212>  PRT
16     <213>  Synthetic
17
18     <400>  173
19
20     Ser Glu Phe Asp Lys Thr Tyr Thr Ala Ala
21     1              5              10
22
23     <210>  174
24     <211>  10
25     <212>  PRT
26     <213>  Synthetic
27
28     <400>  174
29
30     Lys Gln Ala Glu Lys Tyr Thr Ala Glu Gln
31     1              5              10
32
33     <210>  175
34     <211>  10
35     <212>  PRT
36     <213>  Synthetic
37
38     <400>  175
39
40     Ser Glu Phe Asp Ala Ala Phe Ala Ala Ala
41     1              5              10
42
43     <210>  176
44     <211>  10
45     <212>  PRT
46     <213>  Synthetic
47
48     <400>  176
49
50     Ser Glu Phe Asp Lys Thr Phe Thr Ala Lys
51     1              5              10
52
53     <210>  177
54     <211>  10
55     <212>  PRT
56     <213>  Synthetic
57

```

```

1  <400> 177
2
3  Ala Glu Lys Tyr Thr Ala Glu Gln Cys Ala
4  1                      5                      10
5
6  <210> 178
7  <211> 10
8  <212> PRT
9  <213> Synthetic
10
11 <400> 178
12
13 Ala Thr Tyr Thr Ala Lys Glu Ala Ala Thr
14 1                      5                      10
15
16 <210> 179
17 <211> 10
18 <212> PRT
19 <213> Synthetic
20
21 <400> 179
22
23 Lys Ala Tyr Thr Ala Lys Glu Ala Ala Thr
24 1                      5                      10
25
26 <210> 180
27 <211> 10
28 <212> PRT
29 <213> Synthetic
30
31 <400> 180
32
33 Lys Thr Ala Thr Ala Lys Glu Ala Ala Thr
34 1                      5                      10
35
36 <210> 181
37 <211> 10
38 <212> PRT
39 <213> Synthetic
40
41 <400> 181
42
43 Lys Thr Tyr Ala Ala Lys Glu Ala Ala Thr
44 1                      5                      10
45
46 <210> 182
47 <211> 10
48 <212> PRT
49 <213> Synthetic
50
51 <400> 182
52
53 Lys Thr Tyr Thr Ala Ala Glu Ala Ala Thr
54 1                      5                      10
55
56 <210> 183
57 <211> 10

```

```

1  <212>  PRT
2  <213>  Synthetic
3
4  <400>  183
5
6  Lys Thr Tyr Thr Ala Lys Ala Ala Ala Thr
7  1                      5                      10
8
9  <210>  184
10 <211>  10
11 <212>  PRT
12 <213>  Synthetic
13
14 <400>  184
15
16 Lys Thr Tyr Thr Ala Lys Glu Ala Ala Ala
17 1                      5                      10
18
19 <210>  185
20 <211>  10
21 <212>  PRT
22 <213>  Synthetic
23
24 <400>  185
25
26 Glu Lys Tyr Thr Ala Glu Gln Cys Ala Lys
27 1                      5                      10
28
29 <210>  186
30 <211>  10
31 <212>  PRT
32 <213>  Synthetic
33
34 <400>  186
35
36 Ala Ala Phe Ala Ala Ala Glu Ala Ala Thr
37 1                      5                      10
38
39 <210>  187
40 <211>  10
41 <212>  PRT
42 <213>  Synthetic
43
44 <400>  187
45
46 Lys Thr Phe Thr Ala Lys Glu Ala Ala Thr
47 1                      5                      10
48
49 <210>  188
50 <211>  10
51 <212>  PRT
52 <213>  Synthetic
53
54 <400>  188
55
56 Gln Ala Leu His Cys Glu Lys Lys Tyr Leu
57 1                      5                      10

```

```

1
2 <210> 189
3 <211> 10
4 <212> PRT
5 <213> Synthetic
6
7 <400> 189
8
9 Gln Ala Leu Asp Ser Lys Ala Ala Tyr Leu
10 1 5 10
11
12 <210> 190
13 <211> 10
14 <212> PRT
15 <213> Synthetic
16
17 <400> 190
18
19 Gln Ala Leu Asp Ser Lys Asn Asn Phe Leu
20 1 5 10
21
22 <210> 191
23 <211> 10
24 <212> PRT
25 <213> Synthetic
26
27 <400> 191
28
29 Gln Ala Leu His Cys Glu Asn Asn Phe Leu
30 1 5 10
31
32 <210> 192
33 <211> 10
34 <212> PRT
35 <213> Synthetic
36
37 <400> 192
38
39 Cys Glu Lys Lys Tyr Leu Arg Ile Gln Asp
40 1 5 10
41
42 <210> 193
43 <211> 10
44 <212> PRT
45 <213> Synthetic
46
47 <400> 193
48
49 Ser Lys Asn Asn Phe Leu Arg Val Gln Glu
50 1 5 10
51
52 <210> 194
53 <211> 10
54 <212> PRT
55 <213> Synthetic
56
57 <400> 194

```

```

1
2 Ser Glu Asn Asn Tyr Leu Arg Val Gln Glu
3 1 5 10
4
5 <210> 195
6 <211> 10
7 <212> PRT
8 <213> Synthetic
9
10 <400> 195
11
12 Ala Leu Arg Val Gln Glu Asn Ala Leu Thr
13 1 5 10
14
15 <210> 196
16 <211> 10
17 <212> PRT
18 <213> Synthetic
19
20 <400> 196
21
22 Tyr Ala Arg Val Gln Glu Asn Ala Leu Thr
23 1 5 10
24
25 <210> 197
26 <211> 10
27 <212> PRT
28 <213> Synthetic
29
30 <400> 197
31
32 Tyr Leu Ala Val Gln Glu Asn Ala Leu Thr
33 1 5 10
34
35 <210> 198
36 <211> 10
37 <212> PRT
38 <213> Synthetic
39
40 <400> 198
41
42 Tyr Leu Arg Ala Gln Glu Asn Ala Leu Thr
43 1 5 10
44
45 <210> 199
46 <211> 10
47 <212> PRT
48 <213> Synthetic
49
50 <400> 199
51
52 Tyr Leu Arg Val Ala Glu Asn Ala Leu Thr
53 1 5 10
54
55 <210> 200
56 <211> 10
57 <212> PRT

```

1 <213> Synthetic
 2
 3 <400> 200
 4
 5 Tyr Leu Arg Val Gln Ala Asn Ala Leu Thr
 6 1 5 10
 7
 8 <210> 201
 9 <211> 10
 10 <212> PRT
 11 <213> Synthetic
 12
 13 <400> 201
 14
 15 Tyr Leu Arg Val Gln Glu Ala Ala Leu Thr
 16 1 5 10
 17
 18 <210> 202
 19 <211> 10
 20 <212> PRT
 21 <213> Synthetic
 22
 23 <400> 202
 24
 25 Tyr Leu Arg Val Gln Glu Asn Ala Ala Thr
 26 1 5 10
 27
 28 <210> 203
 29 <211> 10
 30 <212> PRT
 31 <213> Synthetic
 32
 33 <400> 203
 34
 35 Tyr Leu Arg Val Gln Glu Asn Ala Leu Ala
 36 1 5 10
 37
 38 <210> 204
 39 <211> 10
 40 <212> PRT
 41 <213> Synthetic
 42
 43 <400> 204
 44
 45 Tyr Leu Arg Ile Gln Asp Asp Thr Leu Thr
 46 1 5 10
 47
 48 <210> 205
 49 <211> 10
 50 <212> PRT
 51 <213> Synthetic
 52
 53 <400> 205
 54
 55 Tyr Leu Thr Val Ala Ala Ala Ala Leu Thr
 56 1 5 10
 57

```

1  <210> 206
2  <211> 10
3  <212> PRT
4  <213> Synthetic
5
6  <400> 206
7
8  Phe Leu Arg Val Gln Glu Asn Ala Leu Thr
9  1                      5                      10
10
11 <210> 207
12 <211> 10
13 <212> PRT
14 <213> Synthetic
15
16 <400> 207
17
18 Asn Asn Tyr Leu Arg Val Gln Glu Asn Ala
19 1                      5                      10
20
21 <210> 208
22 <211> 10
23 <212> PRT
24 <213> Synthetic
25
26 <400> 208
27
28 Lys Lys Tyr Leu Arg Ile Gln Asp Asp Thr
29 1                      5                      10
30
31 <210> 209
32 <211> 10
33 <212> PRT
34 <213> Synthetic
35
36 <400> 209
37
38 Asn Asn Phe Leu Arg Val Gln Glu Asn Ala
39 1                      5                      10
40
41 <210> 210
42 <211> 10
43 <212> PRT
44 <213> Synthetic
45
46 <400> 210
47
48 Asn Ala Tyr Leu Arg Val Gln Glu Asn Ala
49 1                      5                      10
50
51 <210> 211
52 <211> 10
53 <212> PRT
54 <213> Synthetic
55
56 <400> 211
57

```



```

1  Ala Thr Tyr Glu Glu Ala Lys Leu Arg Phe
2  1                      5                      10
3
4  <210> 212
5  <211> 10
6  <212> PRT
7  <213> Synthetic
8
9  <400> 212
10
11 Glu Ala Tyr Glu Glu Ala Leu Lys Arg Phe
12 1                      5                      10
13
14 <210> 213
15 <211> 10
16 <212> PRT
17 <213> Synthetic
18
19 <400> 213
20
21 Glu Thr Ala Glu Glu Ala Leu Lys Arg Phe
22 1                      5                      10
23
24 <210> 214
25 <211> 10
26 <212> PRT
27 <213> Synthetic
28
29 <400> 214
30
31 Glu Thr Tyr Ala Glu Ala Leu Lys Arg Phe
32 1                      5                      10
33
34 <210> 215
35 <211> 10
36 <212> PRT
37 <213> Synthetic
38
39 <400> 215
40
41 Glu Thr Tyr Glu Ala Ala Leu Lys Arg Phe
42 1                      5                      10
43
44 <210> 216
45 <211> 10
46 <212> PRT
47 <213> Synthetic
48
49 <400> 216
50
51 Glu Thr Tyr Glu Glu Ala Ala Lys Arg Phe
52 1                      5                      10
53
54 <210> 217
55 <211> 10
56 <212> PRT
57 <213> Synthetic

```

```

1
2 <400> 217
3
4 Glu Thr Tyr Glu Glu Ala Leu Ala Arg Phe
5 1 5 10
6
7 <210> 218
8 <211> 10
9 <212> PRT
10 <213> Synthetic
11
12 <400> 218
13
14 Glu Thr Tyr Glu Glu Ala Leu Lys Ala Phe
15 1 5 10
16
17 <210> 219
18 <211> 10
19 <212> PRT
20 <213> Synthetic
21
22 <400> 219
23
24 Glu Thr Tyr Glu Glu Ala Leu Lys Arg Ala
25 1 5 10
26
27 <210> 220
28 <211> 10
29 <212> PRT
30 <213> Synthetic
31
32 <400> 220
33
34 Gly Thr Asn Ala Gln Ser Leu Ala Asp Phe
35 1 5 10
36
37 <210> 221
38 <211> 10
39 <212> PRT
40 <213> Synthetic
41
42 <400> 221
43
44 Glu Thr Tyr Glu Ala Ala Leu Ala Ala Phe
45 1 5 10
46
47 <210> 222
48 <211> 10
49 <212> PRT
50 <213> Synthetic
51
52 <400> 222
53
54 Glu Thr Phe Glu Glu Ala Leu Lys Arg Phe
55 1 5 10
56
57 <210> 223

```

```

1  <211> 10
2  <212> PRT
3  <213> Synthetic
4
5  <400> 223
6
7  Tyr Glu Glu Ala Leu Lys Thr Phe Ala Lys
8  1          5          10
9
10 <210> 224
11 <211> 10
12 <212> PRT
13 <213> Synthetic
14
15 <400> 224
16
17 Phe Glu Glu Ala Leu Lys Arg Phe Ala Lys
18 1          5          10
19
20 <210> 225
21 <211> 10
22 <212> PRT
23 <213> Synthetic
24
25 <400> 225
26
27 Ala Ala Leu Lys Arg Phe Ala Lys Leu Leu
28 1          5          10
29
30 <210> 226
31 <211> 10
32 <212> PRT
33 <213> Synthetic
34
35 <400> 226
36
37 Glu Ala Ala Lys Arg Phe Ala Lys Leu Leu
38 1          5          10
39
40 <210> 227
41 <211> 10
42 <212> PRT
43 <213> Synthetic
44
45 <400> 227
46
47 Glu Ala Leu Ala Arg Phe Ala Lys Leu Leu
48 1          5          10
49
50 <210> 228
51 <211> 10
52 <212> PRT
53 <213> Synthetic
54
55 <400> 228
56
57 Glu Ala Leu Lys Ala Phe Ala Lys Leu Leu

```

```

1      1              5              10
2
3      <210>  229
4      <211>  10
5      <212>  PRT
6      <213>  Synthetic
7
8      <400>  229
9
10     Glu Ala Leu Lys Arg Ala Ala Lys Leu Leu
11     1              5              10
12
13     <210>  230
14     <211>  10
15     <212>  PRT
16     <213>  Synthetic
17
18     <400>  230
19
20     Glu Ala Leu Lys Arg Phe Ala Ala Leu Leu
21     1              5              10
22
23     <210>  231
24     <211>  10
25     <212>  PRT
26     <213>  Synthetic
27
28     <400>  231
29
30     Glu Ala Leu Lys Arg Phe Ala Lys Ala Leu
31     1              5              10
32
33     <210>  232
34     <211>  10
35     <212>  PRT
36     <213>  Synthetic
37
38     <400>  232
39
40     Glu Ala Leu Lys Arg Phe Ala Lys Leu Ala
41     1              5              10
42
43     <210>  233
44     <211>  10
45     <212>  PRT
46     <213>  Synthetic
47
48     <400>  233
49
50     Gln Ser Leu Ala Asp Phe Ala Lys Gln Leu
51     1              5              10
52
53     <210>  234
54     <211>  10
55     <212>  PRT
56     <213>  Synthetic
57

```

```

1  <400> 234
2
3  Ala Ala Leu Ala Ala Phe Ala Lys Leu Leu
4  1          5          10
5
6  <210> 235
7  <211> 10
8  <212> PRT
9  <213> Synthetic
10
11 <400> 235
12
13 Leu Ala Asp Phe Ala Lys Gln Leu Ser Asp
14 1          5          10
15
16 <210> 236
17 <211> 10
18 <212> PRT
19 <213> Synthetic
20
21 <400> 236
22
23 Asp Phe Ala Lys Gln Leu Ser Asp Glu Arg
24 1          5          10
25
26 <210> 237
27 <211> 10
28 <212> PRT
29 <213> Synthetic
30
31 <400> 237
32
33 Ala Phe Ala Ala Leu Leu Ser Asp Arg Lys
34 1          5          10
35
36 <210> 238
37 <211> 10
38 <212> PRT
39 <213> Synthetic
40
41 <400> 238
42
43 Ser Thr Ala Ala Ala Pro Thr Tyr Phe Pro
44 1          5          10
45
46 <210> 239
47 <211> 10
48 <212> PRT
49 <213> Synthetic
50
51 <400> 239
52
53 Leu Lys Arg Phe Ala Lys Leu Leu Ser Asp
54 1          5          10
55
56 <210> 240
57 <211> 10

```

```

1  <212>  PRT
2  <213>  Synthetic
3
4  <400>  240
5
6  Gln Ala Leu Asp Ser Glu Asn Asn Phe Leu
7  1          5          10
8
9  <210>  241
10 <211>  10
11 <212>  PRT
12 <213>  Synthetic
13
14 <400>  241
15
16 Ser Asp Leu Ala Asp Phe Ala Lys Gln Leu
17 1          5          10
18
19 <210>  242
20 <211>  55
21 <212>  DNA
22 <213>  Synthetic
23
24 <400>  242
25 ggagctcgag aaaagagagg ctgaagcttc attgaattac aaaaaaatgc tgttg      55
26
27
28 <210>  243
29 <211>  42
30 <212>  DNA
31 <213>  Synthetic
32
33 <400>  243
34 tccaactgt cctggtccat aagaagcttt gtttgctcgg ag      42
35
36
37 <210>  244
38 <211>  36
39 <212>  DNA
40 <213>  Synthetic
41
42 <400>  244
43 gcttcttatg gaccaggaca gttgggagaa atggtg      36
44
45 <210>  245
46 <211>  39
47 <212>  DNA
48 <213>  Synthetic
49
50 <400>  245
51 ggtctagagg aattctcatt acctaattga agcaaatgc      39
52
53 <210>  246
54 <211>  1128
55 <212>  DNA
56 <213>  Synthetic
57

```

1 <400> 246
 2 tcgagaaaag agagggctgaa gcttcattga attacaaaaa aatgctgttg ctctcattag 60
 3
 4 gcactggcac tacttcagag tttgataaaa catatacagc aaaagaggga gctacctgga 120
 5
 6 ctgctgtaca ttggatgtta gttatacaga aaatgactga tgcagcaagt tcttacatga 180
 7
 8 ctgattatta cctttctact gcttttcaag ctcttgattc aaaaaacaat tacctcaggg 240
 9
 10 ttcaagaaaa tgcattaaca ggcacaaact ctgaaatgga tgatgcttct gaggctaata 300
 11
 12 tgggaattatt agtacaagtt ggtgaaaact tattgaagaa accagtttcc gaagacaatc 360
 13
 14 ctgaaaccta tgaggaaact ctaaagaggt ttgcaaaatt gctctctgat aggaagaaac 420
 15
 16 tccgagcaaa caaagcttct tatggaccag gacagttggg agaaatggtg actgttctta 480
 17
 18 gtattgatgg aggtggaatt agagggatca ttccggctac cattctcgaa tttcttgaag 540
 19
 20 gacaacttca ggaatggac aataatgcag atgcaagact tgcagattac tttgatgtaa 600
 21
 22 ttggaggaac aagtacagga ggtttattga ctgctatgat aagtactcca aatgaaaaca 660
 23
 24 atcgaccctt tgctgtctgc aaagaaattg taccttttta ctctgaacat ggcctcaga 720
 25
 26 tttttaatcc tagtggtaaa atttttaggcc caaaatatga tggaaaatat cttatgcaa 780
 27
 28 ttcttcaaga aaaacttgga gaaactcgtg tgcatacaagc ttgacagaa gttgtcatct 840
 29
 30 caagctttga catcaaaaca aataagccag taatattcac taagtcaa ttagcaaaact 900
 31
 32 ctccagaatt ggaatgctaag atgtatgaca taagtattc cacagcagca gctccaacat 960
 33
 34 attttctcc gcattacttt gttactaata ctagtaattg agatgaatat gagttcaatc 1020
 35
 36 ttgttgatgg tgctgttgct actgttgctg atccggcggtt attatccatt agcgttgcaa 1080
 37
 38
 39 cgagacttgc aaaaaggat ccagcatttg cttcaattag gtaatgag 1128
 40
 41 <210> 247
 42 <211> 366
 43 <212> PRT
 44 <213> Synthetic
 45
 46 <400> 247
 47
 48 Ser Leu Asn Tyr Lys Lys Met Leu Leu Leu Ser Leu Gly Thr Gly Thr
 49 1 5 10 15
 50
 51 Thr Ser Glu Phe Asp Lys Thr Tyr Thr Ala Lys Glu Ala Ala Thr Trp
 52 20 25 30
 53
 54 Thr Ala Val His Trp Met Leu Val Ile Gln Lys Met Thr Asp Ala Ala
 55 35 40 45
 56
 57 Ser Ser Tyr Met Thr Asp Tyr Tyr Leu Ser Thr Ala Phe Gln Ala Leu

1	50	55	60
2			
3	Asp Ser Lys Asn Asn Tyr Leu Arg Val Gln Glu Asn Ala Leu Thr Gly		
4	65	70	75
5			
6	Thr Thr Thr Glu Met Asp Asp Ala Ser Glu Ala Asn Met Glu Leu Leu		
7		85	90
8			95
9	Val Gln Val Gly Glu Asn Leu Leu Lys Lys Pro Val Ser Glu Asp Asn		
10		100	105
11			110
12	Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg Phe Ala Lys Leu Leu Ser		
13		115	120
14			125
15	Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala Ser Tyr Gly Pro Gly Gln		
16		130	135
17			140
18	Leu Gly Glu Met Val Thr Val Leu Ser Ile Asp Gly Gly Gly Ile Arg		
19		145	150
20			155
21	Gly Ile Ile Pro Ala Thr Ile Leu Glu Phe Leu Glu Gly Gln Leu Gln		
22		165	170
23			175
24	Glu Met Asp Asn Asn Ala Asp Ala Arg Leu Ala Asp Tyr Phe Asp Val		
25		180	185
26			190
27	Ile Gly Gly Thr Ser Thr Gly Gly Leu Leu Thr Ala Met Ile Ser Thr		
28		195	200
29			205
30	Pro Asn Glu Asn Asn Arg Pro Phe Ala Ala Ala Lys Glu Ile Val Pro		
31		210	215
32			220
33	Phe Tyr Phe Glu His Gly Pro Gln Ile Phe Asn Pro Ser Gly Gln Ile		
34		225	230
35			235
36	Leu Gly Pro Lys Tyr Asp Gly Lys Tyr Leu Met Gln Val Leu Gln Glu		
37		245	250
38			255
39	Lys Leu Gly Glu Thr Arg Val His Gln Ala Leu Thr Glu Val Val Ile		
40		260	265
41			270
42	Ser Ser Phe Asp Ile Lys Thr Asn Lys Pro Val Ile Phe Thr Lys Ser		
43		275	280
44			285
45	Asn Leu Ala Asn Ser Pro Glu Leu Asp Ala Lys Met Tyr Asp Ile Ser		
46		290	295
47			300
48	Tyr Ser Thr Ala Ala Ala Pro Thr Tyr Phe Pro Pro His Tyr Phe Val		
49		305	310
50			315
51	Thr Asn Thr Ser Asn Gly Asp Glu Tyr Glu Phe Asn Leu Val Asp Gly		
52		325	330
53			335
54	Ala Val Ala Thr Val Ala Asp Pro Ala Leu Leu Ser Ile Ser Val Ala		
55		340	345
56			350
57	Thr Arg Leu Ala Gln Lys Asp Pro Ala Phe Ala Ser Ile Arg		

1		355	360	365	
2	<210>	248			
3	<211>	55			
4	<212>	DNA			
5	<213>	Synthetic			
6					
7	<400>	248			
8	ggagctcgag	aaaagagagg	ctgaagctaa	tactagtaat	ggagatgaat atgag 55
9					
10	<210>	249			
11	<211>	39			
12	<212>	DNA			
13	<213>	Synthetic			
14					
15	<400>	249			
16	ggtctagagg	aattctcatt	aagtaacaaa	gtaatgcgg	39
17					
18	<210>	250			
19	<211>	1128			
20	<212>	DNA			
21	<213>	Synthetic			
22					
23	<400>	250			
24	tcgagaaaag	agaggctgaa	gctaatacta	gtaatggaga	tgaatatgag ttcaatcttg 60
25					
26	ttgatggtgc	tggtgctact	gttgctgac	cggcgttatt	atccattagc gttgcaacga 120
27					
28	gacttgcaca	aaaggatcca	gcatttgctt	caattaggtc	attgaattac aaaaaaatgc 180
29					
30	tggtgctctc	attaggcact	ggcactactt	cagagtttga	taaaacatat acagcaaaaag 240
31					
32	aggcagctac	ctggactgct	gtacattgga	tgtagtttat	acagaaaaatg actgatgcag 300
33					
34	caagttctta	catgactgat	tattaccttt	ctactgcttt	tcaagctctt gattcaaaaa 360
35					
36	acaattacct	cagggttcaa	gaaaatgcat	taacaggcac	aactactgaa atggatgatg 420
37					
38	cttctgaggc	taatatggaa	ttattagtac	aagttggtga	aaacttattg aagaaaccag 480
39					
40	tttcgaaga	caatcctgaa	acctatgagg	aagctctaaa	gaggtttgca aaattgctct 540
41					
42	ctgataggaa	gaaactccga	gcaaacaaa	cttcttatgg	accaggacag ttgggagaaa 600
43					
44	tggtgactgt	tcttagtatt	gatggagggt	gaattagagg	gatcattccg gctaccattc 660
45					
46	tcgaatttct	tgaaggacaa	cttcaggaaa	tggacaataa	tgccagatga agacttgcag 720
47					
48	attactttga	tgtaattgga	ggaacaagta	caggagggtt	attgactgct atgataagta 780
49					
50	ctccaaatga	aaacaatcga	ccctttgctg	ctgccaaa	ga aattgtacct ttttacttcg 840
51					
52	aacatggccc	tcagattttt	aatcctatgt	gtcaaatttt	aggcccaaaa tatgatggaa 900
53					
54	aatatcttat	gcaagttctt	caagaaaaac	ttggagaaac	tcgtgtgcat caagctttga 960
55					
56	cagaagttgt	catctcaagc	tttgacatca	aaacaaataa	gccagtaata ttcactaagt 1020
57					

```

1  caaatttagc aaactctcca gaattggatg ctaagatgta tgacataagt tattccacag 1080
2
3  cagcagctcc aacatatattt cctccgcatt actttgttac ttaatgag 1128
4
5  <210> 251
6  <211> 366
7  <212> PRT
8  <213> Synthetic
9
10 <400> 251
11
12 Asn Thr Ser Asn Gly Asp Glu Tyr Glu Phe Asn Leu Val Asp Gly Ala
13 1 5 10 15
14
15 Val Ala Thr Val Ala Asp Pro Ala Leu Leu Ser Ile Ser Val Ala Thr
16 20 25 30
17
18 Arg Leu Ala Gln Lys Asp Pro Ala Phe Ala Ser Ile Arg Ser Leu Asn
19 35 40 45
20
21 Tyr Lys Lys Met Leu Leu Leu Ser Leu Gly Thr Gly Thr Thr Ser Glu
22 50 55 60
23
24 Phe Asp Lys Thr Tyr Thr Ala Lys Glu Ala Ala Thr Trp Thr Ala Val
25 65 70 75 80
26
27 His Trp Met Leu Val Ile Gln Lys Met Thr Asp Ala Ala Ser Ser Tyr
28 85 90 95
29
30 Met Thr Asp Tyr Tyr Leu Ser Thr Ala Phe Gln Ala Leu Asp Ser Lys
31 100 105 110
32
33 Asn Asn Tyr Leu Arg Val Gln Glu Asn Ala Leu Thr Gly Thr Thr Thr
34 115 120 125
35
36 Glu Met Asp Asp Ala Ser Glu Ala Asn Met Glu Leu Leu Val Gln Val
37 130 135 140
38
39 Gly Glu Asn Leu Leu Lys Lys Pro Val Ser Glu Asp Asn Pro Glu Thr
40 145 150 155 160
41
42 Tyr Glu Glu Ala Leu Lys Arg Phe Ala Lys Leu Leu Ser Asp Arg Lys
43 165 170 175
44
45 Lys Leu Arg Ala Asn Lys Ala Ser Tyr Gly Pro Gly Gln Leu Gly Glu
46 180 185 190
47
48 Met Val Thr Val Leu Ser Ile Asp Gly Gly Gly Ile Arg Gly Ile Ile
49 195 200 205
50
51 Pro Ala Thr Ile Leu Glu Phe Leu Glu Gly Gln Leu Gln Glu Met Asp
52 210 215 220
53
54 Asn Asn Ala Asp Ala Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly
55 225 230 235 240
56
57 Thr Ser Thr Gly Gly Leu Leu Thr Ala Met Ile Ser Thr Pro Asn Glu

```

	245	250	255	
1				
2				
3	Asn Asn Arg Pro Phe Ala Ala Ala	Lys Glu Ile Val Pro Phe Tyr Phe		
4	260	265	270	
5				
6	Glu His Gly Pro Gln Ile Phe Asn	Pro Ser Gly Gln Ile Leu Gly Pro		
7	275	280	285	
8				
9	Lys Tyr Asp Gly Lys Tyr Leu Met	Gln Val Leu Gln Glu Lys Leu Gly		
10	290	295	300	
11				
12	Glu Thr Arg Val His Gln Ala Leu Thr	Glu Val Val Ile Ser Ser Phe		
13	305	310	315	320
14				
15	Asp Ile Lys Thr Asn Lys Pro Val Ile	Phe Thr Lys Ser Asn Leu Ala		
16	325	330	335	
17				
18	Asn Ser Pro Glu Leu Asp Ala Lys	Met Tyr Asp Ile Ser Tyr Ser Thr		
19	340	345	350	
20				
21	Ala Ala Ala Pro Thr Tyr Phe Pro	Pro His Tyr Phe Val Thr		
22	355	360	365	
23				
24	<210>	252		
25	<211>	55		
26	<212>	DNA		
27	<213>	Synthetic		
28				
29	<400>	252		
30	ggagctcgag aaaagagagg ctgaagctag ttattccaca gcagcagctc caaca		55	
31				
32	<210>	253		
33	<211>	39		
34	<212>	DNA		
35	<213>	Synthetic		
36				
37	<400>	253		
38	ggtctagagg aattctcatt atatgtcata catcttagc		39	
39				
40	<210>	254		
41	<211>	1128		
42	<212>	DNA		
43	<213>	Synthetic		
44				
45	<400>	254		
46	tcgagaaaag agaggctgaa gctagttatt ccacagcagc agctccaaca tatattcttc		60	
47				
48	cgcattactt tggtactaat actagtaatg gagatgaata tgagttcaat ctgtgtgatg		120	
49				
50	gtgctgttgc tactgttgct gatccggcgt tattatccat tagcgttgca acgagacttg		180	
51				
52	cacaaaaagga tccagcattt ggttcaatta ggctattgaa ttacaaaaaa atgctgttgc		240	
53				
54	tctcattagg cactggcact acttcagagt ttgataaaac atatacagca aaagaggcag		300	
55				
56	ctacctggac tgctgtacat tggatgttag ttatacagaa aatgactgat gcagcaagtt		360	
57				

1 cttacatgac tgattattac ctttctactg cttttcaagc tcttgattca aaaaacaatt 420
 2
 3 acctcagggg tcaagaaaat gcattaacag gcacaactac tgaaatggat gatgcttctg 480
 4
 5 aggctaatat ggaattatta gtacaagttg gtgaaaactt attgaagaaa ccagtttccg 540
 6
 7 aagacaatcc tgaacctat gaggaagctc taaagaggtt tgcaaaattg ctctctgata 600
 8
 9 ggaagaaaact ccgagcaaac aaagcttctt atggaccagg acagttggga gaaatggtga 660
 10
 11 ctgttcttag tattgatgga ggtggaatta gagggatcat tccggctacc attctcgaat 720
 12
 13 ttcttgaagg acaacttcag gaaatggaca ataatgcaga tgcaagactt gcagattact 780
 14
 15 ttgatgtaat tggaggaaca agtacaggag gtttattgac tgctatgata agtactccaa 840
 16
 17 atgaaaacaa tcgacccttt gctgctgcca aagaattgt acctttttac ttcgaacatg 900
 18
 19 gccctcagat ttttaactct agtgggtcaaa ttttaggcc aaatatgat ggaaatatc 960
 20
 21 ttatgcaagt tcttcaagaa aaacttgagg aaactcgtgt gcataagct ttgcagagaag 1020
 22
 23 ttgtcatctc aagctttgac atcaaaaacaa ataagccagt aatattcact aagtcaaatt 1080
 24
 25 tagcaaacct tccagaattg gatgctaaga tgtatgacat ataatgag 1128
 26
 27 <210> 255
 28 <211> 366
 29 <212> PRT
 30 <213> Synthetic
 31
 32 <400> 255
 33
 34 Ser Tyr Ser Thr Ala Ala Ala Pro Thr Tyr Phe Pro Pro His Tyr Phe
 35 1 5 10 15
 36
 37 Val Thr Asn Thr Ser Asn Gly Asp Glu Tyr Glu Phe Asn Leu Val Asp
 38 20 25 30
 39
 40 Gly Ala Val Ala Thr Val Ala Asp Pro Ala Leu Leu Ser Ile Ser Val
 41 35 40 45
 42
 43 Ala Thr Arg Leu Ala Gln Lys Asp Pro Ala Phe Ala Ser Ile Arg Ser
 44 50 55 60
 45
 46 Leu Asn Tyr Lys Lys Met Leu Leu Leu Ser Leu Gly Thr Gly Thr Thr
 47 65 70 75 80
 48
 49 Ser Glu Phe Asp Lys Thr Tyr Thr Ala Lys Glu Ala Ala Thr Trp Thr
 50 85 90 95
 51
 52 Ala Val His Trp Met Leu Val Ile Gln Lys Met Thr Asp Ala Ala Ser
 53 100 105 110
 54
 55 Ser Tyr Met Thr Asp Tyr Tyr Leu Ser Thr Ala Phe Gln Ala Leu Asp
 56 115 120 125
 57

1 Ser Lys Asn Asn Tyr Leu Arg Val Gln Glu Asn Ala Leu Thr Gly Thr
 2 130 135 140
 3
 4 Thr Thr Glu Met Asp Asp Ala Ser Glu Ala Asn Met Glu Leu Leu Val
 5 145 150 155 160
 6
 7 Gln Val Gly Glu Asn Leu Leu Lys Lys Pro Val Ser Glu Asp Asn Pro
 8 165 170 175
 9
 10 Glu Thr Tyr Glu Glu Ala Leu Lys Arg Phe Ala Lys Leu Leu Ser Asp
 11 180 185 190
 12
 13 Arg Lys Lys Leu Arg Ala Asn Lys Ala Ser Tyr Gly Pro Gly Gln Leu
 14 195 200 205
 15
 16 Gly Glu Met Val Thr Val Leu Ser Ile Asp Gly Gly Gly Ile Arg Gly
 17 210 215 220
 18
 19 Ile Ile Pro Ala Thr Ile Leu Glu Phe Leu Glu Gly Gln Leu Gln Glu
 20 225 230 235 240
 21
 22 Met Asp Asn Asn Ala Asp Ala Arg Leu Ala Asp Tyr Phe Asp Val Ile
 23 245 250 255
 24
 25 Gly Gly Thr Ser Thr Gly Gly Leu Leu Thr Ala Met Ile Ser Thr Pro
 26 260 265 270
 27
 28 Asn Glu Asn Asn Arg Pro Phe Ala Ala Ala Lys Glu Ile Val Pro Phe
 29 275 280 285
 30
 31 Tyr Phe Glu His Gly Pro Gln Ile Phe Asn Pro Ser Gly Gln Ile Leu
 32 290 295 300
 33
 34 Gly Pro Lys Tyr Asp Gly Lys Tyr Leu Met Gln Val Leu Gln Glu Lys
 35 305 310 315 320
 36
 37 Leu Gly Glu Thr Arg Val His Gln Ala Leu Thr Glu Val Val Ile Ser
 38 325 330 335
 39
 40 Ser Phe Asp Ile Lys Thr Asn Lys Pro Val Ile Phe Thr Lys Ser Asn
 41 340 345 350
 42
 43 Leu Ala Asn Ser Pro Glu Leu Asp Ala Lys Met Tyr Asp Ile
 44 355 360 365
 45
 46 <210> 256
 47 <211> 55
 48 <212> DNA
 49 <213> Synthetic
 50
 51
 52 <400> 256
 53 ggagctcgag aaaagagagg ctgaagctac atatacagca aaagaggcag ctacc 55
 54
 55 <210> 257
 56 <211> 39
 57 <212> DNA
 58 <213> Synthetic

1
2 <400> 257
3 ggtctagagg aattctcatt atttatcaaa ctctgaagt 39
4
5 <210> 258
6 <211> 1128
7 <212> DNA
8 <213> Synthetic
9
10 <400> 258
11 tcgagaaaag agaggetgaa gctacatata cagcaaaaaga ggcagctacc tggactgctg 60
12 tacattggat gttagtata cagaaaaata ctgatgcagc aagttcttac atgactgatt 120
13 attacctttc tactgctttt caagctcttg attcaaaaaa caattacctc aggggtcaag 180
14
15
16
17 aaaatgcatt aacaggcaca actactgaaa tggatgatgc ttctgaggct aatatggaat 240
18
19 tattagtaca agttgggtgaa aacttattga agaaaccagt ttccgaagac aatcctgaaa 300
20
21 cctatgagga agctctaaag aggtttgcaa aattgctctc tgataggaag aaactccgat 360
22
23 caaacaagaag ttcttatgga ccaggacagt tgggagaaat ggtgactgtt cttagtattg 420
24
25 atggagggtg aattagaggg atcattccgg ctaccattct cgaatttctt gaaggacaac 480
26
27 ttcaggaagt ggacaataat gcagatgcaa gacttgacga ttactttgat gtaattggag 540
28
29 gaacaagtag aggaggttta ttgactgcta tgataagtag tccaaatgaa aacaatcgac 600
30
31 cctttgtctg tgccaaagaa attgtacctt ttaacttcga acatggccct cagattttta 660
32
33 atcctagtgg tcaaatttta ggcccaaaat atgatggaaa atatcttatg caagttcttc 720
34
35 aagaaaaact tggagaaact cgtgtgcacg aagctttgac agaagttgtc atctcaagct 780
36
37 ttgacatcaa aacaataaag ccagtaatat tcaactaagtc aaatttagca aactctccag 840
38
39 aattggatgc taagatgtat gacataagtt attccacagc agcagctcca acatattttc 900
40
41 ctccgcatta ctttgttact aatactagta atggagatga atatgagttc aatcttgttg 960
42
43 atggtgtctg tgctactgtt gctgatccgg cggtattatc cattagcggt gcaacgagac 1020
44
45 ttgcacaaaa ggatccagca ttgtcttcaa tttaggtcatt gaattacaaa aaaatgctgt 1080
46
47 tgctctcatt aggcactggc actacttcag agtttgataa ataatgag 1128
48
49 <210> 259
50 <211> 366
51 <212> PRT
52 <213> Synthetic
53
54 <400> 259
55
56 Thr Tyr Thr Ala Lys Glu Ala Ala Thr Trp Thr Ala Val His Trp Met
57 1 5 10 15

1
2 Leu Val Ile Gln Lys Met Thr Asp Ala Ala Ser Ser Tyr Met Thr Asp
3 20 25 30
4
5 Tyr Tyr Leu Ser Thr Ala Phe Gln Ala Leu Asp Ser Lys Asn Asn Tyr
6 35 40 45
7
8 Leu Arg Val Gln Glu Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp
9 50 55 60
10
11 Asp Ala Ser Glu Ala Asn Met Glu Leu Leu Val Gln Val Gly Glu Asn
12 65 70 75 80
13
14 Leu Leu Lys Lys Pro Val Ser Glu Asp Asn Pro Glu Thr Tyr Glu Glu
15 85 90 95
16
17 Ala Leu Lys Arg Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg
18 100 105 110
19
20 Ser Asn Lys Ala Ser Tyr Gly Pro Gly Gln Leu Gly Glu Met Val Thr
21 115 120 125
22
23 Val Leu Ser Ile Asp Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala Thr
24 130 135 140
25
26 Ile Leu Glu Phe Leu Glu Gly Gln Leu Gln Glu Met Asp Asn Asn Ala
27 145 150 155 160
28
29 Asp Ala Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr
30 165 170 175
31
32 Gly Gly Leu Leu Thr Ala Met Ile Ser Thr Pro Asn Glu Asn Asn Arg
33 180 185 190
34
35 Pro Phe Ala Ala Ala Lys Glu Ile Val Pro Phe Tyr Phe Glu His Gly
36 195 200 205
37
38 Pro Gln Ile Phe Asn Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr Asp
39 210 215 220
40
41 Gly Lys Tyr Leu Met Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg
42 225 230 235 240
43
44 Val His Gln Ala Leu Thr Glu Val Val Ile Ser Ser Phe Asp Ile Lys
45 245 250 255
46
47 Thr Asn Lys Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Asn Ser Pro
48 260 265 270
49
50 Glu Leu Asp Ala Lys Met Tyr Asp Ile Ser Tyr Ser Thr Ala Ala Ala
51 275 280 285
52
53 Pro Thr Tyr Phe Pro Pro His Tyr Phe Val Thr Asn Thr Ser Asn Gly
54 290 295 300
55
56 Asp Glu Tyr Glu Phe Asn Leu Val Asp Gly Ala Val Ala Thr Val Ala
57 305 310 315 320

1
 2 Asp Pro Ala Leu Leu Ser Ile Ser Val Ala Thr Arg Leu Ala Gln Lys
 3 325 330 335
 4
 5 Asp Pro Ala Phe Ala Ser Ile Arg Ser Leu Asn Tyr Lys Lys Met Leu
 6 340 345 350
 7
 8 Leu Leu Ser Leu Gly Thr Gly Thr Thr Ser Glu Phe Asp Lys
 9 355 360 365
 10
 11 <210> 260
 12 <211> 55
 13 <212> DNA
 14 <213> Synthetic
 15
 16 <400> 260
 17 ggagctcgag aaaagagagg ctgaagctaa tgcattaaca ggcacaacta ctgaa 55
 18
 19 <210> 261
 20 <211> 39
 21 <212> DNA
 22 <213> Synthetic
 23
 24 <400> 261
 25 ggtctagagg aattctcatt attcttgaac cctgaggta 39
 26
 27 <210> 262
 28 <211> 1128
 29 <212> DNA
 30 <213> Synthetic
 31
 32 <400> 262
 33 tcgagaaaag agaggctgaa gctaattgcat taacaggcac aactactgaa atggatgatg 60
 34
 35 cttctgaggc taatatggaa ttattagtagc aagttgggtga aaacttattg aagaaaccag 120
 36
 37 ttccgaaga caatcctgaa acctatgagg aagctctaaa gaggtttgca aaattgtctt 180
 38
 39 ctgataggaa gaaactccga gcaacaaaag cttcttatgg accaggacag ttgggagaaa 240
 40
 41 tggtgactgt tcttagtatt gatggagggtg gaattagagg gatcattccg gctaccattc 300
 42
 43 tcgaatttct tgaaggacaa cttcaggaaa tggacaataa tgcagatgca agacttgcag 360
 44
 45 attactttga tgtaattlga ggaacaagta caggaggttt attgactgtc atgataagta 420
 46
 47 ctccaaatga aaacaatcga ccctttgctg ctgccaaaga aattgtacct ttttactctg 480
 48
 49 aacatggccc tcagattttt aatcctagtgt gtcaaatattt aggcccaaaa tatgatggaa 540
 50
 51 aatatcttat gcaagttctt caagaaaaac ttggagaaac tcgtgtgcat caagctttga 600
 52
 53 cagaagtgtg catctcaagc ttgacatca aaacaaataa gccagtaata ttactaagt 660
 54
 55 caaatattagc aaactctcca gaattggatg ctaagatgta tgacataagt tattccacag 720
 56
 57 cagcagctcc aacatatattt cctccgcatt actttgttac taatactagt aatggagatg 780


```

1
2 aatatgagtt caatcttggt gatgggtgctg ttgctactgt tgctgatccg gcgttattat 840
3
4 ccattagcgt tgcaacgaga cttgcacaaa aggatccagc atttgcctca attaggtcat 900
5
6 tgaattacaa aaaaatgctg ttgctctcat taggcactgg cactacttca gagtttgata 960
7
8 aaacatatat agcaaaaagag gcagctacct ggactgctgt acattggatg ttagttatac 1020
9
10 agaaaaatgac tgatgcagca agttctttaca tgactgatta ttacctttct actgcttttc 1080
11
12 aagctcttga ttcaaaaaac aattacctca ggggttcaaga ataatgag 1128
13
14 <210> 263
15 <211> 366
16 <212> PRT
17 <213> Synthetic
18
19 <400> 263
20
21 Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu Ala
22 1 5 10 15
23
24 Asn Met Glu Leu Leu Val Gln Val Gly Glu Asn Leu Leu Lys Lys Pro
25 20 25 30
26
27 Val Ser Glu Asp Asn Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg Phe
28 35 40 45
29
30 Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala Ser
31 50 55 60
32
33 Tyr Gly Pro Gly Gln Leu Gly Glu Met Val Thr Val Leu Ser Ile Asp
34 65 70 75 80
35
36 Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala Thr Ile Leu Glu Phe Leu
37 85 90 95
38
39 Glu Gly Gln Leu Gln Glu Met Asp Asn Asn Ala Asp Ala Arg Leu Ala
40 100 105 110
41
42 Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly Leu Leu Thr
43 115 120 125
44
45 Ala Met Ile Ser Thr Pro Asn Glu Asn Asn Arg Pro Phe Ala Ala Ala
46 130 135 140
47
48 Lys Glu Ile Val Pro Phe Tyr Phe Glu His Gly Pro Gln Ile Phe Asn
49 145 150 155 160
50
51 Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr Asp Gly Lys Tyr Leu Met
52 165 170 175
53
54 Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His Gln Ala Leu
55 180 185 190
56
57 Thr Glu Val Val Ile Ser Ser Phe Asp Ile Lys Thr Asn Lys Pro Val

```

1		195		200		205	
2							
3	Ile	Phe	Thr	Lys	Ser	Asn	Leu
4		210				215	
5							
6	Met	Tyr	Asp	Ile	Ser	Tyr	Ser
7		225				230	
8							
9	Pro	His	Tyr	Phe	Val	Thr	Asn
10					245		
11							
12	Asn	Leu	Val	Asp	Gly	Ala	Val
13					260		
14							
15	Ser	Ile	Ser	Val	Ala	Thr	Arg
16					275		
17							
18	Ser	Ile	Arg	Ser	Leu	Asn	Tyr
19					290		
20							
21	Thr	Gly	Thr	Thr	Ser	Glu	Phe
22		305				310	
23							
24	Ala	Thr	Trp	Thr	Ala	Val	His
25					325		
26							
27	Asp	Ala	Ala	Ser	Ser	Tyr	Met
28					340		
29							
30	Gln	Ala	Leu	Asp	Ser	Lys	Asn
31					355		
32							
33	<210>	264					
34	<211>	1158					
35	<212>	DNA					
36	<213>	Synthetic					
37							
38	<400>	264					
39	atggccacca	ccaagagctt	cctcatcctg	atcttcatga	tcctggccac	caccagcagc	60
40							
41	accttcgcc	agctcggcga	gatggtgacc	gtgctctcca	tcgacggcgg	tggcatcagg	120
42							
43	ggcatcatcc	cggccacat	cctggagttc	ctggagggcc	aactccagga	gatggacaac	180
44							
45	aacgccgacg	cccgcctggc	cgactacttc	gacgtgatcg	gtggcaccag	caccggcggt	240
46							
47	ctcctgacgc	ccatgatctc	cactccgaac	gagaacaacc	gccctctcgc	cgctgcgaag	300
48							
49	gagatcgtcc	cgcttctaett	cgaacacggc	ectcagattt	tcaacccttc	gggtcaaate	360
50							
51	ctggggcccca	agtacgacgg	caagtacctt	atgcaagtgc	ttcaggagaa	gctgggcgag	420
52							
53	actagggtgc	accaggcgct	gaccgaggtc	gtcatctcca	gcttcgacat	caagaccaac	480
54							
55	aagccagtc	tcttcaccaa	gtccaacctg	gccaacagcc	cggagctgga	cgctaagatg	540
56							
57	tacgacatct	cctactccac	tgctgccgct	cccacgtact	tcctctccga	ctactctgtc	600

1
 2 accaacacca gcaacggcga cgagtagcag ttcaaccttg ttgacgggtgc ggtggctacg 660
 3
 4 gtggcgccgacc cggcgctect gtccatcagc gtgcgccagc gcctggccca gaagatcca 720
 5
 6 gccttcgcta gcattaggag cctcaactac aagaagatgc tgctgctcag cctgggcact 780
 7
 8 ggcaacgacct ccgagttcga caagacctac actgccaaag aggccgctac ctggaccgcc 840
 9
 10 gtccattgga tgctggtcat ccagaagatg acggacgccg cttccageta catgaccgac 900
 11
 12 tactacctct ccactgcgtt ccaggcgctt gactccaaga acaactacct ccgtgttcag 960
 13
 14 gagaatgcc tcaactggcac caccgaccgag atggacgatg cctccgaggc caacatggag 1020
 15
 16 ctgctcgtcc aggtgggtga gaacctcttg aagaagcccg tctccgaaga caatcccgag 1080
 17
 18 acctatgagg aagcgctcaa gcgcttgcc aagctgctct ctgataggaa gaaactccgc 1140
 19
 20 gctaacaagg ccagctac 1158
 21
 22 <210> 265
 23 <211> 386
 24 <212> PRT
 25 <213> Synthetic
 26
 27 <400> 265
 28
 29 Met Ala Thr Lys Ser Phe Leu Ile Leu Ile Phe Met Ile Leu Ala
 30 1 5 10 15
 31
 32 Thr Thr Ser Ser Thr Phe Ala Gln Leu Gly Glu Met Val Thr Val Leu
 33 20 25 30
 34
 35 Ser Ile Asp Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala Thr Ile Leu
 36 35 40 45
 37
 38 Glu Phe Leu Glu Gly Gln Leu Gln Glu Met Asp Asn Asn Ala Asp Ala
 39 50 55 60
 40
 41 Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly
 42 65 70 75 80
 43
 44 Leu Leu Thr Ala Met Ile Ser Thr Pro Asn Glu Asn Asn Arg Pro Phe
 45 85 90 95
 46
 47 Ala Ala Ala Lys Glu Ile Val Pro Phe Tyr Phe Glu His Gly Pro Gln
 48 100 105 110
 49
 50 Ile Phe Asn Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr Asp Gly Lys
 51 115 120 125
 52
 53 Tyr Leu Met Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His
 54 130 135 140
 55
 56 Gln Ala Leu Thr Glu Val Val Ile Ser Ser Phe Asp Ile Lys Thr Asn
 57 145 150 155 160

1
 2 Lys Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Asn Ser Pro Glu Leu
 3 165 170 175
 4
 5 Asp Ala Lys Met Tyr Asp Ile Ser Tyr Ser Thr Ala Ala Ala Pro Thr
 6 180 185 190
 7
 8 Tyr Phe Pro Pro His Tyr Phe Val Thr Asn Thr Ser Asn Gly Asp Glu
 9 195 200 205
 10
 11 Tyr Glu Phe Asn Leu Val Asp Gly Ala Val Ala Thr Val Ala Asp Pro
 12 210 215 220
 13
 14 Ala Leu Leu Ser Ile Ser Val Ala Thr Arg Leu Ala Gln Lys Asp Pro
 15 225 230 235 240
 16
 17 Ala Phe Ala Ser Ile Arg Ser Leu Asn Tyr Lys Lys Met Leu Leu Leu
 18 245 250 255
 19
 20 Ser Leu Gly Thr Gly Thr Thr Ser Glu Phe Asp Lys Thr Tyr Thr Ala
 21 260 265 270
 22
 23 Lys Glu Ala Ala Thr Trp Thr Ala Val His Trp Met Leu Val Ile Gln
 24 275 280 285
 25
 26 Lys Met Thr Asp Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Leu Ser
 27 290 295 300
 28
 29 Thr Ala Phe Gln Ala Leu Asp Ser Lys Asn Asn Tyr Leu Arg Val Gln
 30 305 310 315 320
 31
 32 Glu Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu
 33 325 330 335
 34
 35 Ala Asn Met Glu Leu Leu Val Gln Val Gly Glu Asn Leu Leu Lys Lys
 36 340 345 350
 37
 38 Pro Val Ser Glu Asp Asn Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg
 39 355 360 365
 40
 41 Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala
 42 370 375 380
 43
 44 Ser Tyr
 45 385
 46
 47 <210> 266
 48 <211> 55
 49 <212> DNA
 50 <213> Synthetic
 51
 52 <400> 266
 53 ggagctcgag aaaagagagg ctgaagctag cctcaactac aagaagatgc tgctg
 54
 55 <210> 267
 56 <211> 42
 57 <212> DNA

55

1	<213> Synthetic	
2		
3	<400> 267	
4	gccgagctgt cctggtccgt agctggcctt gttagcgcg ag	42
5		
6	<210> 268	
7	<211> 36	
8	<212> DNA	
9	<213> Synthetic	
10		
11	<400> 268	
12	gccagctacg gaccaggaca gctcggcgag atgggtg	36
13		
14	<210> 269	
15	<211> 39	
16	<212> DNA	
17	<213> Synthetic	
18		
19	<400> 269	
20	ggctagagg aattctcatt acctaagtct agcgaaggc	39
21		
22	<210> 270	
23	<211> 1167	
24	<212> DNA	
25	<213> Synthetic	
26		
27	<400> 270	
28	atggccacca ccaagagctt cctcctcctg atcttcctga tcttgccac caccagcagc	60
29		
30	accttcgcca gcctcaacta caagaagatg ctgctgctca gcctgggac tggcacgacc	120
31		
32	tccagattcg acaagaccta cactgccaa gaggccgcta cctggaccgc cgtccattgg	180
33		
34	atgctggtca tccagaagat gacggacgcc gcttcagct acatgaccga ctactacctc	240
35		
36	tccactgcgt tccaggcgct tgaactcaag aacaactacc tccgtgttca ggagaatgcc	300
37		
38	ctcactggca ccacgaccga gatggacgat gcctccgag ccaacatgga gctgctcgtc	360
39		
40	caggtgggtg agaacctcct gaagaagccc gtctccgaag acaatcccg gacctatgag	420
41		
42	gaagcgctca agcgctttgc caagctgctc tctgatagga agaaactccg cgctaacaag	480
43		
44	gccagctacg gaccaggaca gctcggcgag atggtagacc tgctctccat cgacggcggt	540
45		
46	ggcatcaggg gcatcctccc ggccaccatc ctggagttcc tggagggcca actccaggag	600
47		
48	atggacaaca acgccgacgc ccgcctggcc gactacttcg acgtgatcgg tggcaccagc	660
49		
50	accggcggtc tcctgaccgc catgatctcc actccgaacg agaacaaccg ccccttcgcc	720
51		
52	gctgcgaagg agatcgctcc gtctacttc gaacacggcc ctacagattt caaccctcg	780
53		
54	ggtaaatcc tgggccccta gtacacggc aagtacctta tgcaagtgtc tcaggagaag	840
55		
56	ctgggcgaga ctagggtgca ccaggcgctg accgaggtcg tcatctccag ctctcgacatc	900
57		

1 aagaccaaca agccagtcac cttcaccaag tccaacctgg ccaacagccc ggagctggac 960
 2
 3 gctaagatgt acgacatctc ctactccact gctgccgctc ccacgtactt ccctccgcac 1020
 4
 5 tactctgtca ccaacaccag caacggcgac gactacgagt tcaacctgtg tgacgggtgcg 1080
 6
 7 gtggctacgg tggcgacccc ggcgctcctg tccatcagcg tcgccacgcg cctggcccgag 1140
 8
 9 aaggatccag ccttcgctag cattagg 1167
 10
 11 <210> 271
 12 <211> 389
 13 <212> PRT
 14 <213> Synthetic
 15
 16 <400> 271
 17
 18 Met Ala Thr Thr Lys Ser Phe Leu Ile Leu Ile Phe Met Ile Leu Ala
 19 1 5 10 15
 20
 21 Thr Thr Ser Ser Thr Phe Ala Ser Leu Asn Tyr Lys Lys Met Leu Leu
 22 20 25 30
 23
 24 Leu Ser Leu Gly Thr Gly Thr Thr Ser Glu Phe Asp Lys Thr Tyr Thr
 25 35 40 45
 26
 27 Ala Lys Glu Ala Ala Thr Trp Thr Ala Val His Trp Met Leu Val Ile
 28 50 55 60
 29
 30 Gln Lys Met Thr Asp Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Leu
 31 65 70 75 80
 32
 33 Ser Thr Ala Phe Gln Ala Leu Asp Ser Lys Asn Asn Tyr Leu Arg Val
 34 85 90 95
 35
 36 Gln Glu Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala Ser
 37 100 105 110
 38
 39 Glu Ala Asn Met Glu Leu Leu Val Gln Val Gly Glu Asn Leu Leu Lys
 40 115 120 125
 41
 42 Lys Pro Val Ser Glu Asp Asn Pro Glu Thr Tyr Glu Glu Ala Leu Lys
 43 130 135 140
 44
 45 Arg Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys
 46 145 150 155 160
 47
 48 Ala Ser Tyr Gly Pro Gly Gln Leu Gly Glu Met Val Thr Val Leu Ser
 49 165 170 175
 50
 51 Ile Asp Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala Thr Ile Leu Glu
 52 180 185 190
 53
 54 Phe Leu Glu Gly Gln Leu Gln Glu Met Asp Asn Asn Ala Asp Ala Arg
 55 195 200 205
 56
 57 Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly Leu

```

1          210                      215                      220
2
3  Leu Thr Ala Met Ile Ser Thr Pro Asn Glu Asn Asn Arg Pro Phe Ala
4  225                      230                      235                      240
5
6  Ala Ala Lys Glu Ile Val Pro Phe Tyr Phe Glu His Gly Pro Gln Ile
7                      245                      250                      255
8
9  Phe Asn Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr Asp Gly Lys Tyr
10                      260                      265                      270
11
12  Leu Met Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His Gln
13                      275                      280                      285
14
15  Ala Leu Thr Glu Val Val Ile Ser Ser Phe Asp Ile Lys Thr Asn Lys
16                      290                      295                      300
17
18  Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Asn Ser Pro Glu Leu Asp
19  305                      310                      315
20
21  Ala Lys Met Tyr Asp Ile Ser Tyr Ser Thr Ala Ala Ala Pro Thr Tyr
22                      325                      330                      335
23
24  Phe Pro Pro His Tyr Phe Val Thr Asn Thr Ser Asn Gly Asp Glu Tyr
25                      340                      345                      350
26
27  Glu Phe Asn Leu Val Asp Gly Ala Val Ala Thr Val Ala Asp Pro Ala
28                      355                      360                      365
29
30  Leu Leu Ser Ile Ser Val Ala Thr Arg Leu Ala Gln Lys Asp Pro Ala
31                      370                      375                      380
32
33  Phe Ala Ser Ile Arg
34  385
35
36  <210> 272
37  <211> 55
38  <212> DNA
39  <213> Synthetic
40
41  <400> 272
42  ggagctcgag aaaagagagg ctgaagctac tgccaaggagg gccgtacct ggacc 55
43
44  <210> 273
45  <211> 39
46  <212> DNA
47  <213> Synthetic
48
49  <400> 273
50  ggtctagagg aattctcatt acttgtcgaa ctcgagagg 39
51
52  <210> 274
53  <211> 1167
54  <212> DNA
55  <213> Synthetic
56
57  <400> 274

```

1	atgccacca ccaagagctt cctcatcctg atcttcatga tcttggccac caccagcagc	60
2		
3	accttcgcca cctacactgc caaggaggcc gctacctgga ccgccgtcca ttggatgctg	120
4		
5	gtcatccaga agatgacgga cgccgcttcc agctacatga ccgactacta cctctccact	180
6		
7	gcgttcaggc cgcttgactc caagaacaac tacctccgtg ttcaggagaa tgccctcaact	240
8		
9	ggcaccacga ccgagatgga cgatgcctcc gaggccaaca tggagctgct cgccaggtg	300
10		
11	ggtgagaacc tcctgaagaa gcccgcttcc gaagacaatc ccgagaccta tgaggaagcg	360
12		
13	ctcaagcgct ttgccaagct gctctctgat aggaagaaac tccgcgctaa caaggccagc	420
14		
15	tacggaccag gacagctcgg cgagatgggt accgtgctct ccacgcagcg cggtggcatc	480
16		
17	aggggcatca tcccgccac catcctggag ttctggagg gccaaactcca ggagatggac	540
18		
19	aacaacccgc acgcccgcct ggccgactac ttgcagtgta tcggtggcac cagcaccggc	600
20		
21	ggtctcctga ccgccatgat ctccactccg aacgagaaca accgcccctt cgccgctgcg	660
22		
23	aaggagatcg tcccgttcta cttcgaacac gccctcaga ttttcaacct ctcggtcaa	720
24		
25	atcctggggc ccaagtacga cggcaagtac cttatgcaag tgcttcagga gaagctgggc	780
26		
27	gagactaggc tgaccaggc gctgaccgag gtgcgtcatct ccagcttcga catcaagacc	840
28		
29	aacaagccag tcattctcac caagtccaac ctggccaaca gcccgagct ggacgctaag	900
30		
31	atgtacgaca tctcctaact cactgctgcc gctcccagct acttccctcc gcaactcttc	960
32		
33	gtcaccaaca ccagcaacgg cgacgagtac gagttcaacc ttgttgacgg tgcggtggct	1020
34		
35	acggtggcgg acccggcgct cctgtccatc agcgtcgcca cgcgcctggc ccagaaggat	1080
36		
37	ccagccttcg ctagcattag gagcctcaac tacaagaaga tgctgctgct cagcctgggc	1140
38		
39	actggcacga cctccgagtt cgacaag	1167
40		
41	<210> 275	
42	<211> 389	
43	<212> PRT	
44	<213> Synthetic	
45		
46	<400> 275	
47		
48	Met Ala Thr Thr Lys Ser Phe Leu Ile Leu Ile Phe Met Ile Leu Ala	
49	1 5 10 15	
50		
51	Thr Thr Ser Ser Thr Phe Ala Thr Tyr Thr Ala Lys Glu Ala Ala Thr	
52	20 25 30	
53		
54	Trp Thr Ala Val His Trp Met Leu Val Ile Gln Lys Met Thr Asp Ala	
55	35 40 45	
56		
57	Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Leu Ser Thr Ala Phe Gln Ala	

1	50	55	60
2			
3	Leu Asp Ser Lys Asn Asn Tyr Leu Arg Val Gln Glu Asn Ala Leu Thr		
4	65	70	75 80
5			
6	Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu Ala Asn Met Glu Leu		
7		85	90 95
8			
9	Leu Val Gln Val Gly Glu Asn Leu Leu Lys Lys Pro Val Ser Glu Asp		
10		100	105 110
11			
12	Asn Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg Phe Ala Lys Leu Leu		
13		115	120 125
14			
15	Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala Ser Tyr Gly Pro Gly		
16		130	135 140
17			
18	Gln Leu Gly Glu Met Val Thr Val Leu Ser Ile Asp Gly Gly Gly Ile		
19		145	150 155 160
20			
21	Arg Gly Ile Ile Pro Ala Thr Ile Leu Glu Phe Leu Glu Gly Gln Leu		
22		165	170 175
23			
24	Gln Glu Met Asp Asn Asn Ala Asp Ala Arg Leu Ala Asp Tyr Phe Asp		
25		180	185 190
26			
27	Val Ile Gly Gly Thr Ser Thr Gly Glu Leu Leu Thr Ala Met Ile Ser		
28		195	200 205
29			
30	Thr Pro Asn Glu Asn Asn Arg Pro Phe Ala Ala Ala Lys Glu Ile Val		
31		210	215 220
32			
33	Pro Phe Tyr Phe Glu His Gly Pro Gln Ile Phe Asn Pro Ser Gly Gln		
34		225	230 235 240
35			
36	Ile Leu Gly Pro Lys Tyr Asp Gly Lys Tyr Leu Met Gln Val Leu Gln		
37		245	250 255
38			
39	Glu Lys Leu Gly Glu Thr Arg Val His Gln Ala Leu Thr Glu Val Val		
40		260	265 270
41			
42	Ile Ser Ser Phe Asp Ile Lys Thr Asn Lys Pro Val Ile Phe Thr Lys		
43		275	280 285
44			
45	Ser Asn Leu Ala Asn Ser Pro Glu Leu Asp Ala Lys Met Tyr Asp Ile		
46		290	295 300
47			
48	Ser Tyr Ser Thr Ala Ala Ala Pro Thr Tyr Phe Pro Pro His Tyr Phe		
49		305	310 315 320
50			
51	Val Thr Asn Thr Ser Asn Gly Asp Glu Tyr Glu Phe Asn Leu Val Asp		
52		325	330 335
53			
54	Gly Ala Val Ala Thr Val Ala Asp Pro Ala Leu Leu Ser Ile Ser Val		
55		340	345 350
56			
57	Ala Thr Arg Leu Ala Gln Lys Asp Pro Ala Phe Ala Ser Ile Arg Ser		

```

1          355          360          365
2
3 Leu Asn Tyr Lys Lys Met Leu Leu Leu Ser Leu Gly Thr Gly Thr Thr
4          370          375          380
5
6 Ser Glu Phe Asp Lys
7 385
8
9 <210> 276
10 <211> 7
11 <212> PRT
12 <213> Synthetic
13
14 <400> 276
15
16 Gly Gly Gly Ser Gly Gly Gly
17 1 5
18
19 <210> 277
20 <211> 3
21 <212> PRT
22 <213> Synthetic
23
24 <400> 277
25
26 Gly Pro Gly
27 1
28
29 <210> 278
30 <211> 386
31 <212> PRT
32 <213> Solanum tuberosum
33
34 <400> 278
35
36 Met Ala Thr Thr Lys Ser Phe Leu Ile Leu Phe Phe Met Ile Leu Ala
37 1 5 10 15
38
39 Thr Thr Ser Ser Thr Cys Ala Lys Leu Glu Glu Met Val Thr Val Leu
40 20 25 30
41
42 Ser Ile Asp Gly Gly Gly Ile Lys Gly Ile Ile Pro Ala Ile Ile Leu
43 35 40 45
44
45 Glu Phe Leu Glu Gly Gln Leu Gln Glu Val Asp Asn Asn Lys Asp Ala
46 50 55 60
47
48 Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly
49 65 70 75 80
50
51 Leu Leu Thr Ala Met Ile Thr Thr Pro Asn Glu Asn Asn Arg Pro Phe
52 85 90 95
53
54 Ala Ala Ala Lys Asp Ile Val Pro Phe Tyr Phe Glu His Gly Pro His
55 100 105 110
56
57 Ile Phe Asn Tyr Ser Gly Ser Ile Ile Gly Pro Met Tyr Asp Gly Lys

```

1 115 120 125
2
3 Tyr Leu Leu Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His
4 130 135 140
5
6 Gln Ala Leu Thr Glu Val Ala Ile Ser Ser Phe Asp Ile Lys Thr Asn
7 145 150 155 160
8
9 Lys Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Lys Ser Pro Glu Leu
10 165 170 175
11
12 Asp Ala Lys Met Tyr Asp Ile Cys Tyr Ser Thr Ala Ala Ala Pro Ile
13 180 185 190
14
15 Tyr Phe Pro Pro His Tyr Phe Ile Thr His Thr Ser Asn Gly Asp Ile
16 195 200 205
17
18 Tyr Glu Phe Asn Leu Val Asp Gly Gly Val Ala Thr Val Gly Asp Pro
19 210 215 220
20
21 Ala Leu Leu Ser Leu Ser Val Ala Thr Arg Leu Ala Gln Glu Asp Pro
22 225 230 235 240
23
24 Ala Phe Ser Ser Ile Lys Ser Leu Asp Tyr Lys Gln Met Leu Leu Leu
25 245 250 255
26
27 Ser Leu Gly Thr Gly Thr Asn Ser Glu Phe Asp Lys Thr Tyr Thr Ala
28 260 265 270
29
30 Gln Glu Ala Ala Lys Trp Gly Pro Leu Arg Trp Met Leu Ala Ile Gln
31 275 280 285
32
33 Gln Met Thr Asn Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Ile Ser
34 290 295 300
35
36 Thr Val Phe Gln Ala Arg His Ser Gln Asn Asn Tyr Leu Arg Val Gln
37 305 310 315 320
38
39 Glu Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu
40 325 330 335
41
42 Ala Asn Met Glu Leu Leu Val Gln Val Gly Glu Thr Leu Leu Lys Lys
43 340 345 350
44
45 Pro Val Ser Lys Asp Ser Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg
46 355 360 365
47
48 Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala
49 370 375 380
50
51 Ser Tyr
52 385
53
54 <210> 279
55 <211> 386
56 <212> PRT
57 <213> Solanum tuberosum

1
 2 <400> 279
 3
 4 Met Ala Thr Thr Lys Ser Val Leu Val Leu Phe Phe Met Ile Leu Ala
 5 1 5 10 15
 6
 7 Thr Thr Ser Ser Thr Cys Ala Thr Leu Gly Glu Met Val Thr Val Leu
 8 20 25 30
 9
 10 Ser Ile Asp Gly Gly Gly Ile Lys Gly Ile Ile Pro Ala Thr Ile Leu
 11 35 40 45
 12
 13 Glu Phe Leu Glu Gly Gln Leu Gln Glu Val Asp Asn Asn Lys Asp Ala
 14 50 55 60
 15
 16 Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly
 17 65 70 75 80
 18
 19 Leu Leu Thr Ala Met Ile Thr Thr Pro Asn Glu Asn Asn Arg Pro Phe
 20 85 90 95
 21
 22 Ala Ala Ala Lys Asp Ile Val Pro Phe Tyr Phe Glu His Gly Pro His
 23 100 105 110
 24
 25 Ile Phe Asn Ser Ser Gly Ser Ile Phe Gly Pro Met Tyr Asp Gly Lys
 26 115 120 125
 27
 28 Tyr Phe Leu Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His
 29 130 135 140
 30
 31 Gln Ala Leu Thr Glu Val Ala Ile Ser Ser Phe Asp Ile Lys Thr Asn
 32 145 150 155 160
 33
 34 Lys Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Lys Ser Pro Glu Leu
 35 165 170 175
 36
 37 Asp Ala Lys Met Asn Asp Ile Cys Tyr Ser Thr Ala Ala Ala Pro Thr
 38 180 185 190
 39
 40 Tyr Phe Pro Pro His Tyr Phe Val Thr His Thr Ser Asn Gly Asp Lys
 41 195 200 205
 42
 43 Tyr Glu Phe Asn Leu Val Asp Gly Ala Val Ala Thr Val Gly Asp Pro
 44 210 215 220
 45
 46 Ala Leu Leu Ser Leu Ser Val Arg Thr Lys Leu Ala Gln Val Asp Pro
 47 225 230 235 240
 48
 49 Lys Phe Ala Ser Ile Lys Ser Leu Asn Tyr Asn Glu Met Leu Leu Leu
 50 245 250 255
 51
 52 Ser Leu Gly Thr Gly Thr Asn Ser Glu Phe Asp Lys Thr Tyr Thr Ala
 53 260 265 270
 54
 55 Glu Glu Ala Ala Lys Trp Gly Pro Leu Arg Trp Ile Leu Ala Ile Gln
 56 275 280 285
 57

1 Gln Met Thr Asn Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Leu Ser
 2 290 295 300
 3
 4 Thr Val Phe Gln Ala Arg His Ser Gln Asn Asn Tyr Leu Arg Val Gln
 5 305 310 315 320
 6
 7 Glu Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu
 8 325 330 335
 9
 10 Ala Asn Met Glu Leu Leu Val Gln Val Gly Glu Lys Leu Leu Lys Lys
 11 340 345 350
 12
 13 Pro Val Ser Lys Asp Ser Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg
 14 355 360 365
 15
 16 Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala
 17 370 375 380
 18
 19 Ser Tyr
 20 385
 21
 22 <210> 280
 23 <211> 365
 24 <212> PRT
 25 <213> Solanum tuberosum
 26
 27 <400> 280
 28
 29 Met Ala Leu Glu Glu Met Val Ala Val Leu Ser Ile Asp Gly Gly Gly
 30 1 5 10 15
 31
 32 Ile Lys Gly Ile Ile Pro Gly Thr Ile Leu Glu Phe Leu Glu Gly Gln
 33 20 25 30
 34
 35 Leu Gln Lys Met Asp Asn Asn Ala Asp Ala Arg Leu Ala Asp Tyr Phe
 36 35 40 45
 37
 38 Asp Val Ile Gly Gly Thr Ser Thr Gly Gly Leu Leu Thr Ala Met Ile
 39 50 55 60
 40
 41 Thr Thr Pro Asn Glu Asn Asn Arg Pro Phe Ala Ala Ala Asn Glu Ile
 42 65 70 75 80
 43
 44 Val Pro Phe Tyr Phe Glu His Gly Pro His Ile Phe Asn Ser Arg Tyr
 45 85 90 95
 46
 47 Trp Pro Ile Phe Trp Pro Lys Tyr Asp Gly Lys Tyr Leu Met Gln Val
 48 100 105 110
 49
 50 Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His Gln Ala Leu Thr Glu
 51 115 120 125
 52
 53 Val Ala Ile Ser Ser Phe Asp Ile Lys Thr Asn Lys Pro Val Ile Phe
 54 130 135 140
 55
 56 Thr Lys Ser Asn Leu Ala Lys Ser Pro Glu Leu Asp Ala Lys Thr Tyr
 57 145 150 155 160

1
 2 Asp Ile Cys Tyr Ser Thr Ala Ala Ala Pro Thr Tyr Phe Pro Pro His
 3 165 170 175
 4
 5 Tyr Phe Ala Thr Asn Thr Ile Asn Gly Asp Lys Tyr Glu Phe Asn Leu
 6 180 185 190
 7
 8 Val Asp Gly Ala Val Ala Thr Val Ala Asp Pro Ala Leu Leu Ser Val
 9 195 200 205
 10
 11 Ser Val Ala Thr Arg Arg Ala Gln Glu Asp Pro Ala Phe Ala Ser Ile
 12 210 215 220
 13
 14 Arg Ser Leu Asn Tyr Lys Lys Met Leu Leu Leu Ser Leu Gly Thr Gly
 15 225 230 235 240
 16
 17 Thr Thr Ser Glu Phe Asp Lys Thr His Thr Ala Glu Glu Thr Ala Lys
 18 245 250 255
 19
 20 Trp Gly Ala Leu Gln Trp Met Leu Val Ile Gln Gln Met Thr Glu Ala
 21 260 265 270
 22
 23 Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Leu Ser Thr Val Phe Gln Asp
 24 275 280 285
 25
 26 Leu His Ser Gln Asn Asn Tyr Leu Arg Val Gln Glu Asn Ala Leu Thr
 27 290 295 300
 28
 29 Gly Thr Thr Thr Lys Ala Asp Asp Ala Ser Glu Ala Asn Met Glu Leu
 30 305 310 315 320
 31
 32 Leu Ala Gln Val Gly Glu Asn Leu Leu Lys Lys Pro Val Ser Lys Asp
 33 325 330 335
 34
 35 Asn Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg Phe Ala Lys Leu Leu
 36 340 345 350
 37
 38 Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala Ser Tyr
 39 355 360 365
 40
 41 <210> 281
 42 <211> 364
 43 <212> PRT
 44 <213> Solanum tuberosum
 45
 46 <400> 281
 47
 48 Pro Trp Leu Glu Glu Met Val Thr Val Leu Ser Ile Asp Gly Gly Gly
 49 1 5 10 15
 50
 51 Ile Lys Gly Ile Ile Pro Ala Ile Ile Leu Glu Phe Leu Glu Gly Gln
 52 20 25 30
 53
 54 Leu Gln Glu Val Asp Asn Asn Lys Asp Ala Arg Leu Ala Asp Tyr Phe
 55 35 40 45
 56
 57 Asp Val Ile Gly Gly Thr Ser Thr Gly Gly Leu Leu Thr Ala Met Ile

1	50	55	60
2			
3	Thr Thr Pro Asn Glu Asn Asn Arg Pro Phe Ala Ala Ala Lys Asp Ile		
4	65	70	75 80
5			
6	Val Pro Phe Tyr Phe Glu His Gly Pro His Ile Phe Asn Tyr Ser Gly		
7		85	90 95
8			
9	Ser Ile Leu Gly Pro Met Tyr Asp Gly Lys Tyr Leu Leu Gln Val Leu		
10		100	105 110
11			
12	Gln Glu Lys Leu Gly Glu Thr Arg Val His Gln Ala Leu Thr Glu Val		
13		115	120 125
14			
15	Ala Ile Ser Ser Phe Asp Ile Lys Thr Asn Lys Pro Val Ile Phe Thr		
16		130	135 140
17			
18	Lys Ser Asn Leu Ala Lys Ser Pro Glu Leu Asp Ala Lys Met Tyr Asp		
19		145	150 155 160
20			
21	Ile Cys Tyr Ser Thr Ala Ala Ala Pro Ile Tyr Phe Pro Pro His His		
22		165	170 175
23			
24	Phe Val Thr His Thr Ser Asn Gly Ala Arg Tyr Glu Phe Asn Leu Val		
25		180	185 190
26			
27	Asp Gly Ala Val Ala Thr Val Gly Asp Pro Ala Leu Ser Leu Ser		
28		195	200 205
29			
30	Val Ala Thr Arg Leu Ala Gln Glu Asp Pro Ala Phe Ser Ser Ile Lys		
31		210	215 220
32			
33	Ser Leu Asp Tyr Lys Gln Met Leu Leu Leu Ser Leu Gly Thr Gly Thr		
34		225	230 235 240
35			
36	Asn Ser Glu Phe Asp Lys Thr Tyr Thr Ala Glu Glu Ala Ala Lys Trp		
37		245	250 255
38			
39	Gly Pro Leu Arg Trp Met Leu Ala Ile Gln Gln Met Thr Asn Ala Ala		
40		260	265 270
41			
42	Ser Phe Tyr Met Thr Asp Tyr Tyr Ile Ser Thr Val Phe Gln Ala Arg		
43		275	280 285
44			
45	His Ser Gln Asn Asn Tyr Leu Arg Val Gln Glu Asn Ala Leu Asn Gly		
46		290	295 300
47			
48	Thr Thr Thr Glu Met Asp Asp Ala Ser Glu Ala Asn Met Glu Leu Leu		
49		305	310 315 320
50			
51	Val Gln Val Gly Glu Thr Leu Leu Lys Lys Pro Val Ser Arg Asp Ser		
52		325	330 335
53			
54	Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg Phe Ala Lys Leu Leu Ser		
55		340	345 350
56			
57	Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala Ser Tyr		

```

1           355           360
2
3 <210> 282
4 <211> 386
5 <212> PRT
6 <213> Solanum tuberosum
7
8 <400> 282
9
10 Met Ala Thr Thr Lys Ser Phe Leu Ile Leu Phe Phe Met Ile Leu Ala
11 1 5 10 15
12
13 Thr Thr Ser Ser Thr Cys Ala Lys Leu Glu Glu Met Val Thr Val Leu
14 20 25 30
15
16 Ser Ile Asp Gly Gly Gly Ile Lys Gly Ile Ile Pro Ala Ile Ile Leu
17 35 40 45
18
19 Glu Phe Leu Glu Gly Gln Leu Gln Glu Val Asp Asn Asn Lys Asp Ala
20 50 55 60
21
22 Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly
23 65 70 75 80
24
25 Leu Leu Thr Ala Met Ile Thr Thr Pro Asn Glu Asn Asn Arg Pro Phe
26 85 90 95
27
28 Ala Ala Ala Lys Asp Ile Val Pro Phe Tyr Phe Glu His Gly Pro His
29 100 105 110
30
31 Ile Phe Asn Tyr Ser Gly Ser Ile Leu Gly Pro Met Tyr Asp Gly Lys
32 115 120 125
33
34 Tyr Leu Leu Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His
35 130 135 140
36
37 Gln Ala Leu Thr Glu Val Ala Ile Ser Ser Phe Asp Ile Lys Thr Asn
38 145 150 155 160
39
40 Lys Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Lys Ser Pro Glu Leu
41 165 170 175
42
43 Asp Ala Lys Met Tyr Asp Ile Cys Tyr Ser Thr Ala Ala Ala Pro Ile
44 180 185 190
45
46 Tyr Phe Pro Pro His His Phe Val Thr His Thr Ser Asn Gly Ala Arg
47 195 200 205
48
49 Tyr Glu Phe Asn Leu Val Asp Gly Ala Val Ala Thr Val Gly Asp Pro
50 210 215 220
51
52 Ala Leu Leu Ser Leu Ser Val Ala Thr Arg Leu Ala Gln Glu Asp Pro
53 225 230 235 240
54
55 Ala Phe Ser Ser Ile Lys Ser Leu Asp Tyr Lys Gln Met Leu Leu Leu
56 245 250 255
57

```


1 Ser Leu Gly Thr Gly Thr Asn Ser Glu Phe Asp Lys Thr Tyr Thr Ala
 2 260 265 270
 3
 4 Glu Glu Ala Ala Lys Trp Gly Pro Leu Arg Trp Met Leu Ala Ile Gln
 5 275 280 285
 6
 7 Gln Met Thr Asn Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Ile Ser
 8 290 295 300
 9
 10 Thr Val Phe Gln Ala Arg His Ser Gln Asn Asn Tyr Leu Arg Val Gln
 11 305 310 315 320
 12
 13 Glu Asn Ala Leu Asn Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu
 14 325 330 335
 15
 16 Ala Asn Met Glu Leu Leu Val Gln Val Gly Ala Thr Leu Leu Lys Lys
 17 340 345 350
 18
 19 Pro Val Ser Lys Asp Ser Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg
 20 355 360 365
 21
 22 Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala
 23 370 375 380
 24
 25 Ser Tyr
 26 385
 27
 28 <210> 283
 29 <211> 10
 30 <212> PRT
 31 <213> Synthetic
 32
 33 <400> 283
 34
 35 Ala Phe Phe Asp Lys Thr Tyr Thr Ala Lys
 36 1 5 10
 37
 38 <210> 284
 39 <211> 10
 40 <212> PRT
 41 <213> Synthetic
 42
 43 <400> 284
 44
 45 Cys Ile Phe Asp Ser Thr Tyr Thr Ala Lys
 46 1 5 10
 47
 48 <210> 285
 49 <211> 1161
 50 <212> DNA
 51 <213> Solanum tuberosum
 52
 53 <400> 285
 54 atggcaacta ctaaatcttt ttttaatttta atatttatga tattagcaac tactagtcca 60
 55
 56 acatttgctc agttgggaga aatggtgact gttcttagta ttgatggagg tggaattaga 120
 57

1 gggtatcattc cggtaccat tctcgaattt ctgaaggac aacttcagga aatggacaat 180
2
3 aatgcagatg caagacttgc agattacttt gatgtaattg gaggaacaag tacaggaggt 240
4
5 ttattgactg ctatgataag tactccaaat gaaaacaatc gaccctttgc tgctgccaaa 300
6
7 gaaattgtac ctttttactt cgaacatggc cctcagattt ttaactctag tggtaaaatt 360
8
9 ttaggcccaa aatatgatgg aaaatatctt atgcaagttc ttcaagaaaa acttggagaa 420
10
11 actcgtgtgc atcaagcttt gacagaagtt gtcatctcaa gctttgacat caaaacaaat 480
12
13 aagccagtaa tattcactaa gtcaaattta gcaaaccttc cagaattgga tgctaagatg 540
14
15 tatgacataa gttattccac agcagcagct ccaacatatt ttctctcgca ttactttgtt 600
16
17 actaatacta gtaatggaga tgaatatgag ttcaatcttg ttgatgtgc tgttgctact 660
18
19 gttgctgac cggtgttatt atccattagc gttgcaacga gacttgcaca aaaggatcca 720
20
21 gcatttgcct caattaggtc attgaattac aaaaaaatgc tgttgctctc attaggcact 780
22
23 ggcactactt cacagtttga taaaacatat acagcaaaag aggcagctac ctggactgct 840
24
25 gtacattgga tgttagttat acagaaaatg actgatgcag caagttctta catgactgat 900
26
27 tattaccttt ctactgcttt tcaagctctt gattcaaaaa acaattacct caggggtcaa 960
28
29 gaaaaatgat taacaggcac aactactgaa atggatgatg cttctgagcg taatatggaa 1020
30
31 ttattagtac aagttggtga aaacttattg aagaaccag tttccgaaga caatcctgaa 1080
32
33 acctatgagg aagctctaaa gaggtttgca aaattgctct ctgataggaa gaaactccga 1140
34
35 gcaaacaaag cttcttatta a 1161
36
37 <210> 286
38 <211> 386
39 <212> PRT
40 <213> Solanum tuberosum
41
42 <400> 286
43
44 Met Ala Thr Thr Lys Ser Phe Leu Ile Leu Ile Phe Met Ile Leu Ala
45 1 5 10 15
46
47 Thr Thr Ser Ser Thr Phe Ala Gln Leu Gly Glu Met Val Thr Val Leu
48 20 25 30
49
50 Ser Ile Asp Gly Gly Gly Ile Arg Gly Ile Ile Pro Ala Thr Ile Leu
51 35 40 45
52
53 Glu Phe Leu Glu Gly Gln Leu Gln Glu Met Asp Asn Asn Ala Asp Ala
54 50 55 60
55
56 Arg Leu Ala Asp Tyr Phe Asp Val Ile Gly Gly Thr Ser Thr Gly Gly
57 65 70 75 80

1
2 Leu Leu Thr Ala Met Ile Ser Thr Pro Asn Glu Asn Asn Arg Pro Phe
3 85 90 95
4
5 Ala Ala Ala Lys Glu Ile Val Pro Phe Tyr Phe Glu His Gly Pro Gln
6 100 105 110
7
8 Ile Phe Asn Pro Ser Gly Gln Ile Leu Gly Pro Lys Tyr Asp Gly Lys
9 115 120 125
10
11 Tyr Leu Met Gln Val Leu Gln Glu Lys Leu Gly Glu Thr Arg Val His
12 130 135 140
13
14 Gln Ala Leu Thr Glu Val Val Ile Ser Ser Phe Asp Ile Lys Thr Asn
15 145 150 155 160
16
17 Lys Pro Val Ile Phe Thr Lys Ser Asn Leu Ala Asn Ser Pro Glu Leu
18 165 170 175
19
20 Asp Ala Lys Met Tyr Asp Ile Ser Tyr Ser Thr Ala Ala Pro Thr
21 180 185 190
22
23 Tyr Phe Pro Pro His Tyr Phe Val Thr Asn Thr Ser Asn Gly Asp Glu
24 195 200 205
25
26 Tyr Glu Phe Asn Leu Val Asp Gly Ala Val Ala Thr Val Ala Asp Pro
27 210 215 220
28
29 Ala Leu Leu Ser Ile Ser Val Ala Thr Arg Leu Ala Gln Lys Asp Pro
30 225 230 235 240
31
32 Ala Phe Ala Ser Ile Arg Ser Leu Asn Tyr Lys Lys Met Leu Leu Leu
33 245 250 255
34
35 Ser Leu Gly Thr Gly Thr Thr Ser Glu Phe Asp Lys Thr Tyr Thr Ala
36 260 265 270
37
38 Lys Glu Ala Ala Thr Trp Thr Ala Val His Trp Met Leu Val Ile Gln
39 275 280 285
40
41 Lys Met Thr Asp Ala Ala Ser Ser Tyr Met Thr Asp Tyr Tyr Leu Ser
42 290 295 300
43
44 Thr Ala Phe Gln Ala Leu Asp Ser Lys Asn Asn Tyr Leu Arg Val Gln
45 305 310 315 320
46
47 Glu Asn Ala Leu Thr Gly Thr Thr Thr Glu Met Asp Asp Ala Ser Glu
48 325 330 335
49
50 Ala Asn Met Glu Leu Leu Val Gln Val Gly Glu Asn Leu Leu Lys Lys
51 340 345 350
52
53 Pro Val Ser Glu Asp Asn Pro Glu Thr Tyr Glu Glu Ala Leu Lys Arg
54 355 360 365
55
56 Phe Ala Lys Leu Leu Ser Asp Arg Lys Lys Leu Arg Ala Asn Lys Ala
57 370 375 380

1
 2 Ser Tyr
 3 385
 4
 5 <210> 287
 6 <211> 408
 7 <212> PRT
 8 <213> W098154327 12/3/98 (Equinus (Genus) Pentaclethra species
 9 macrolota)
 10
 11 <400> 287
 12
 13 Met Lys Ser Lys Met Ala Met Leu Leu Leu Leu Phe Cys Val Leu Ser
 14 1 5 10 15
 15
 16 Asn Gln Leu Val Ala Ala Phe Ser Thr Gln Ala Lys Ala Ser Lys Asp
 17 20 25 30
 18
 19 Gly Asn Leu Val Thr Val Leu Ala Ile Asp Gly Gly Gly Ile Arg Gly
 20 35 40 45
 21
 22 Ile Ile Pro Gly Val Ile Leu Lys Gln Leu Glu Ala Thr Leu Gln Arg
 23 50 55 60
 24
 25 Trp Asp Ser Ser Ala Arg Leu Ala Glu Tyr Phe Asp Val Val Ala Gly
 26 65 70 75 80
 27
 28 Thr Ser Thr Gly Gly Ile Ile Thr Ala Ile Leu Thr Ala Pro Asp Pro
 29 85 90 95
 30
 31 Gln Asn Lys Asp Arg Pro Leu Tyr Ala Ala Glu Glu Ile Ile Asp Phe
 32 100 105 110
 33
 34 Tyr Ile Glu His Gly Pro Ser Ile Phe Asn Lys Ser Thr Ala Cys Ser
 35 115 120 125
 36
 37 Leu Pro Gly Ile Phe Cys Pro Lys Tyr Asp Gly Lys Tyr Leu Gln Glu
 38 130 135 140
 39
 40 Ile Ile Ser Gln Lys Leu Asn Glu Thr Leu Leu Asp Gln Thr Thr Thr
 41 145 150 155 160
 42
 43 Asn Val Val Ile Pro Ser Phe Asp Ile Lys Leu Leu Arg Pro Thr Ile
 44 165 170 175
 45
 46 Phe Ser Thr Phe Lys Leu Glu Glu Val Pro Glu Leu Asn Val Lys Leu
 47 180 185 190
 48
 49 Ser Asp Val Cys Met Gly Thr Ser Ala Ala Pro Ile Val Phe Pro Pro
 50 195 200 205
 51
 52 Tyr Tyr Phe Lys His Gly Asp Thr Glu Phe Asn Leu Val Asp Gly Ala
 53 210 215 220
 54
 55 Ile Ile Ala Asp Ile Pro Ala Pro Val Ala Leu Ser Glu Val Leu Gln
 56 225 230 235 240
 57

1 Gln Glu Lys Tyr Lys Asn Lys Glu Ile Leu Leu Ser Ile Gly Thr
 2 245 255
 3
 4 Gly Val Val Lys Pro Gly Glu Gly Tyr Ser Ala Asn Arg Thr Trp Thr
 5 260 265 270
 6
 7 Ile Phe Asp Trp Ser Ser Glu Thr Leu Ile Gly Leu Met Gly His Gly
 8 275 280 285
 9
 10 Thr Arg Ala Met Ser Asp Tyr Tyr Val Gly Ser His Phe Lys Ala Leu
 11 290 295 300
 12
 13 Gln Pro Gln Asn Asn Tyr Leu Arg Ile Gln Glu Tyr Asp Leu Asp Pro
 14 305 310 315 320
 15
 16 Ala Leu Glu Ser Ile Asp Asp Ala Ser Thr Glu Asn Met Glu Asn Leu
 17 325 330 335
 18
 19 Glu Lys Val Gly Gln Ser Leu Leu Asn Glu Pro Val Lys Arg Met Asn
 20 340 345 350
 21
 22 Leu Asn Thr Phe Val Val Glu Glu Thr Gly Glu Gly Thr Asn Ala Glu
 23 355 360 365
 24
 25 Ala Leu Asp Arg Leu Ala Gln Ile Leu Tyr Glu Glu Lys Ile Thr Arg
 26 370 375 380
 27
 28 Gly Leu Gly Lys Ile Ser Leu Glu Val Asp Asn Ile Asp Pro Tyr Thr
 29 385 390 395 400
 30
 31 Glu Arg Val Arg Lys Leu Leu Phe
 32 405
 33
 34 <210> 288
 35 <211> 410
 36 <212> PRT
 37 <213> Zea mays
 38
 39 <400> 288
 40
 41 Met Gly Ser Ile Gly Arg Gly Thr Ala Asn Cys Ala Thr Val Pro Gln
 42 1 5 10 15
 43
 44 Pro Pro Pro Ser Thr Gly Lys Leu Ile Thr Ile Leu Ser Ile Asp Gly
 45 20 25 30
 46
 47 Gly Gly Ile Arg Gly Leu Ile Pro Ala Thr Ile Ile Ala Tyr Leu Glu
 48 35 40 45
 49
 50 Ala Lys Leu Gln Glu Leu Asp Gly Pro Asp Ala Arg Ile Ala Asp Tyr
 51 50 55 60
 52
 53 Phe Asp Val Ile Ala Gly Thr Ser Thr Gly Ala Leu Leu Ala Ser Met
 54 65 70 75 80
 55
 56 Leu Ala Ala Pro Asp Glu Asn Asn Arg Pro Leu Phe Ala Ala Lys Asp
 57 85 90 95

1
2 Leu Thr Thr Phe Tyr Leu Glu Asn Gly Pro Lys Ile Phe Pro Gln Lys
3 100 105 110
4
5 Lys Ala Gly Leu Leu Thr Pro Leu Arg Asn Leu Leu Gly Leu Val Arg
6 115 120 125
7
8 Gly Pro Lys Tyr Asp Gly Val Phe Leu His Asp Lys Ile Lys Ser Leu
9 130 135 140
10
11 Thr His Asp Val Arg Val Ala Asp Thr Val Thr Asn Val Ile Val Pro
12 145 150 155 160
13
14 Ala Phe Asp Val Lys Tyr Leu Gln Pro Ile Ile Phe Ser Thr Tyr Glu
15 165 170 175
16
17 Ala Lys Thr Asp Thr Leu Lys Asn Ala His Leu Ser Asp Ile Cys Ile
18 180 185 190
19
20 Ser Thr Ser Ala Ala Pro Thr Tyr Phe Pro Ala His Phe Phe Lys Thr
21 195 200 205
22
23 Glu Ala Thr Asp Gly Arg Pro Pro Arg Glu Tyr His Leu Val Asp Gly
24 210 215 220
25
26 Gly Val Ala Ala Asn Asn Pro Thr Met Val Ala Met Ser Met Leu Thr
27 225 230 235 240
28
29 Lys Glu Val His Arg Arg Asn Pro Asn Phe Asn Ala Gly Ser Pro Thr
30 245 250 255
31
32 Glu Tyr Thr Asn Tyr Leu Ile Ile Ser Val Gly Thr Gly Ser Ala Lys
33 260 265 270
34
35 Gln Ala Glu Lys Tyr Thr Ala Glu Gln Cys Ala Lys Trp Gly Leu Ile
36 275 280 285
37
38 Gln Trp Leu Tyr Asn Gly Gly Phe Thr Pro Ile Ile Asp Ile Phe Ser
39 290 295 300
40
41 His Ala Ser Ser Asp Met Val Asp Ile His Ala Ser Ile Leu Phe Gln
42 305 310 315 320
43
44 Ala Leu His Cys Glu Lys Lys Tyr Leu Arg Ile Gln Asp Asp Thr Leu
45 325 330 335
46
47 Thr Gly Asn Ala Ser Ser Val Asp Ile Ala Thr Lys Glu Asn Met Glu
48 340 345 350
49
50 Ser Leu Ile Ser Ile Gly Gln Glu Leu Leu Lys Lys Pro Val Ala Arg
51 355 360 365
52
53 Val Asn Ile Asp Thr Gly Val Tyr Glu Ser Cys Asp Gly Glu Gly Thr
54 370 375 380
55
56 Asn Ala Gln Ser Leu Ala Asp Phe Ala Lys Gln Leu Ser Asp Glu Arg
57 385 390 395 400

1
 2 Lys Leu Arg Lys Ser Asn Leu Asn Ser Asn
 3 405 410
 4
 5 <210> 289
 6 <211> 508
 7 <212> PRT
 8 <213> Zea mays
 9
 10 <400> 289
 11
 12 Arg Pro Thr Arg Pro Arg His Pro Arg Asn Thr Gln Lys Arg Gly Ala
 13 1 5 10 15
 14
 15 Leu Leu Val Gly Trp Ile Leu Phe Ser Leu Ala Ala Ser Pro Val Lys
 16 20 25 30
 17
 18 Phe Gln Thr His Met Gly Ser Ile Gly Arg Gly Thr Ala Asn Cys Ala
 19 35 40 45
 20
 21 Thr Val Pro Gln Pro Pro Pro Ser Thr Gly Lys Leu Ile Thr Ile Leu
 22 50 55 60
 23
 24 Ser Ile Asp Gly Gly Gly Ile Arg Gly Leu Ile Pro Ala Thr Ile Ile
 25 65 70 75 80
 26
 27 Ala Tyr Leu Glu Ala Lys Leu Gln Glu Leu Asp Gly Pro Asp Ala Arg
 28 85 90 95
 29
 30 Ile Ala Asp Tyr Phe Asp Val Ile Ala Gly Thr Ser Thr Gly Ala Leu
 31 100 105 110
 32
 33 Leu Ala Ser Met Leu Ala Ala Pro Asp Glu Asn Asn Arg Pro Leu Phe
 34 115 120 125
 35
 36 Ala Ala Lys Asp Leu Thr Thr Phe Tyr Leu Glu Asn Gly Pro Lys Ile
 37 130 135 140
 38
 39 Phe Pro Gln Lys Lys Ala Gly Leu Leu Thr Pro Leu Arg Asn Leu Leu
 40 145 150 155 160
 41
 42 Gly Leu Val Arg Gly Pro Lys Tyr Asp Gly Val Phe Leu His Asp Lys
 43 165 170 175
 44
 45 Ile Lys Ser Leu Thr His Asp Val Arg Val Ala Asp Thr Val Thr Asn
 46 180 185 190
 47
 48 Val Ile Val Pro Ala Phe Asp Val Lys Tyr Leu Gln Pro Ile Ile Phe
 49 195 200 205
 50
 51 Ser Thr Tyr Glu Ala Lys Thr Asp Ala Leu Lys Asn Ala His Leu Ser
 52 210 215 220
 53
 54 Asp Ile Cys Ile Ser Thr Ser Ala Ala Pro Thr Tyr Phe Pro Ala His
 55 225 230 235 240
 56
 57 Phe Phe Lys Thr Glu Ala Thr Asp Gly Arg Pro Pro Arg Glu Tyr His

1
 2
 3 Leu Val Asp Gly Gly Val Ala Ala Asn Asn Pro Thr Met Val Ala Met
 4 260 265 270
 5
 6 Ser Met Leu Thr Lys Glu Val His Arg Arg Asn Pro Asn Phe Asn Ala
 7 275 280 285
 8
 9 Gly Ser Pro Thr Glu Tyr Thr Asn Tyr Leu Ile Ile Ser Val Gly Thr
 10 290 295 300
 11
 12 Gly Ser Ala Lys Gln Ala Glu Lys Tyr Thr Ala Glu Gln Cys Ala Lys
 13 305 310 315 320
 14
 15 Trp Gly Leu Ile Gln Trp Leu Tyr Asn Gly Gly Phe Thr Pro Ile Ile
 16 325 330 335
 17
 18 Asp Ile Phe Ser His Ala Ser Ser Asp Met Val Asp Ile His Ala Ser
 19 340 345 350
 20
 21 Ile Leu Phe Gln Ala Leu His Cys Glu Lys Lys Tyr Leu Arg Ile Gln
 22 355 360 365
 23
 24 Leu Tyr Tyr Ala Gly Tyr Phe Asp Trp Glu Arg Ile Val Arg Gly His
 25 370 375 380
 26
 27 Arg His Gln Gly Glu His Gly Val Ser Asp Ile Asp Arg Pro Gly Ala
 28 385 390 395 400
 29
 30 Ala Gln Glu Ala Ser Gly Glu Ser Glu His Arg His Arg Ala Val Arg
 31 405 410 415
 32
 33 Val Leu Arg Arg Gly His Lys Cys Thr Val Ala Ser Leu Arg Gln Ala
 34 420 425 430
 35
 36 Thr Leu Arg Ala Gln Ala Thr Gln Glu Gln Ser Gln Leu Gln Leu Ile
 37 435 440 445
 38
 39 Asn Thr Ser Leu Ser His Ser Met Cys Ser Phe Arg Arg Phe Thr Val
 40 450 455 460
 41
 42 Ser Tyr Phe Phe Asn Phe Asn Ser Val Cys Val Leu Cys Val Leu Cys
 43 465 470 475 480
 44
 45 Val Tyr Gln Thr Phe Lys Phe Asn Gln Lys Lys Lys Lys Lys Lys
 46 485 490 495
 47
 48 Lys Lys Lys Lys Lys Lys Lys Lys Arg Ala Ala
 49 500 505
 50
 51 <210> 290
 52 <211> 410
 53 <212> PRT
 54 <213> Zea mays
 55
 56 <400> 290
 57

1 Met Gly Ser Ile Gly Arg Gly Thr Ala Asn Cys Ala Thr Val Pro Gln
 2 1 5 10 15
 3
 4 Pro Pro Pro Ser Thr Gly Lys Leu Ile Thr Ile Leu Ser Ile Asp Gly
 5 20 25 30
 6
 7 Gly Gly Ile Arg Gly Leu Ile Pro Ala Thr Ile Ile Ala Tyr Leu Glu
 8 35 40 45
 9
 10 Ala Lys Leu Gln Glu Leu Asp Gly Pro Asp Ala Arg Ile Ala Asp Tyr
 11 50 55 60
 12
 13 Phe Asp Val Ile Ala Gly Thr Ser Thr Gly Ala Leu Leu Ala Ser Met
 14 65 70 75 80
 15
 16 Leu Ala Ala Pro Asp Glu Asn Asn Arg Pro Leu Phe Ala Ala Lys Asp
 17 85 90 95
 18
 19 Leu Thr Thr Phe Tyr Leu Glu Asn Gly Pro Lys Ile Phe Pro Gln Lys
 20 100 105 110
 21
 22 Lys Ala Gly Leu Leu Thr Pro Leu Arg Asn Leu Leu Gly Leu Val Arg
 23 115 120 125
 24
 25 Gly Pro Lys Tyr Asp Gly Val Phe Leu His Asp Lys Ile Lys Ser Leu
 26 130 135 140
 27
 28 Thr His Asp Val Arg Val Ala Asp Thr Val Thr Asn Val Ile Val Pro
 29 145 150 155 160
 30
 31 Ala Phe Asp Val Lys Ser Leu Gln Pro Ile Ile Phe Ser Thr Tyr Glu
 32 165 170 175
 33
 34 Ala Lys Thr Asp Thr Leu Lys Asn Ala His Leu Ser Asp Ile Cys Ile
 35 180 185 190
 36
 37 Ser Thr Ser Ala Ala Pro Thr Tyr Phe Pro Ala His Phe Phe Lys Thr
 38 195 200 205
 39
 40 Glu Ala Thr Asp Gly Arg Pro Pro Arg Glu Tyr His Leu Val Asp Gly
 41 210 215 220
 42
 43 Gly Val Ala Ala Asn Asn Pro Thr Met Val Ala Met Ser Met Leu Thr
 44 225 230 235 240
 45
 46 Lys Glu Val His Arg Arg Asn Pro Asn Phe Asn Ala Gly Ser Pro Thr
 47 245 250 255
 48
 49 Glu Tyr Thr Asn Tyr Leu Ile Ile Ser Val Gly Thr Gly Ser Ala Lys
 50 260 265 270
 51
 52 Gln Ala Glu Lys Tyr Thr Ala Glu Gln Cys Ala Lys Trp Gly Leu Ile
 53 275 280 285
 54
 55 Gln Trp Leu Tyr Asn Gly Gly Phe Thr Pro Ile Ile Asp Ile Phe Ser
 56 290 295 300
 57

1 His Ala Ser Ser Asp Met Val Asp Ile His Ala Ser Ile Leu Phe Gln
 2 305 310 315 320
 3
 4 Ala Leu His Cys Glu Lys Lys Tyr Leu Arg Ile Gln Asp Asp Thr Leu
 5 325 330 335
 6
 7 Thr Gly Asn Ala Ser Ser Val Asp Ile Ala Thr Lys Glu Asn Met Glu
 8 340 345 350
 9
 10 Ser Leu Ile Ser Ile Gly Gln Glu Leu Leu Asn Lys Pro Val Ala Arg
 11 355 360 365
 12
 13 Val Asn Ile Asp Thr Gly Leu Tyr Glu Ser Cys Glu Gly Glu Gly Thr
 14 370 375 380
 15
 16 Asn Ala Gln Ser Leu Ala Asp Phe Ala Lys Gln Leu Ser Asp Glu Arg
 17 385 390 395 400
 18
 19 Lys Leu Arg Lys Ser Asn Leu Asn Ser Asn
 20 405 410
 21
 22 <210> 291
 23 <211> 410
 24 <212> PRT
 25 <213> Zea mays
 26
 27 <400> 291
 28
 29 Met Gly Ser Ile Gly Arg Gly Thr Ala Asn Cys Ala Thr Val Pro Gln
 30 1 5 10 15
 31
 32 Pro Pro Pro Ser Thr Gly Lys Leu Ile Thr Ile Leu Ser Ile Asp Gly
 33 20 25 30
 34
 35 Gly Gly Ile Arg Gly Leu Ile Pro Ala Thr Ile Ile Ala Tyr Leu Glu
 36 35 40 45
 37
 38 Ala Lys Leu Gln Glu Leu Asp Gly Pro Asp Ala Arg Ile Ala Asp Tyr
 39 50 55 60
 40
 41 Phe Asp Val Ile Ala Gly Thr Ser Thr Gly Ala Leu Leu Ala Ser Met
 42 65 70 75 80
 43
 44 Leu Ala Ala Pro Asp Glu Asn Asn Arg Pro Leu Phe Ala Ala Lys Asp
 45 85 90 95
 46
 47 Leu Thr Thr Phe Tyr Leu Glu Asn Gly Pro Lys Ile Phe Pro Gln Lys
 48 100 105 110
 49
 50 Lys Ala Gly Leu Leu Thr Pro Leu Arg Asn Leu Leu Gly Leu Val Arg
 51 115 120 125
 52
 53 Gly Pro Lys Tyr Asp Gly Val Phe Leu His Asp Lys Ile Lys Ser Leu
 54 130 135 140
 55
 56 Thr His Asp Val Arg Val Ala Asp Thr Val Thr Asn Val Ile Val Pro
 57 145 150 155 160

1
2 Ala Phe Asp Val Lys Tyr Leu Gln Pro Ile Ile Phe Ser Thr Tyr Glu
3 165 170 175
4
5 Ala Lys Thr Asp Ala Leu Lys Asn Ala His Leu Ser Asp Ile Cys Ile
6 180 185 190
7
8 Ser Thr Ser Ala Ala Pro Thr Tyr Phe Pro Ala His Phe Phe Lys Thr
9 195 200 205
10
11 Glu Ala Thr Asp Gly Arg Pro Pro Arg Glu Tyr His Leu Val Asp Gly
12 210 215 220
13
14 Gly Val Ala Ala Asn Asn Pro Thr Met Val Ala Met Ser Met Leu Thr
15 225 230 235 240
16
17 Lys Glu Val His Arg Arg Asn Pro Asn Phe Asn Ala Gly Ser Pro Thr
18 245 250 255
19
20 Glu Tyr Thr Asn Tyr Leu Ile Ile Ser Val Gly Thr Gly Ser Ala Lys
21 260 265 270
22
23 Gln Ala Glu Lys Tyr Thr Ala Glu Gln Cys Ala Lys Trp Gly Leu Ile
24 275 280 285
25
26 Gln Trp Leu Tyr Asn Gly Gly Phe Thr Pro Ile Ile Asp Ile Phe Ser
27 290 295 300
28
29 His Ala Ser Ser Asp Met Val Asp Ile His Ala Ser Ile Leu Phe Gln
30 305 310 315 320
31
32 Ala Leu His Cys Glu Lys Lys Tyr Leu Arg Ile Gln Asp Asp Thr Leu
33 325 330 335
34
35 Thr Gly Asn Ala Ser Ser Val Asp Ile Ala Thr Lys Glu Asn Met Glu
36 340 345 350
37
38 Ser Leu Ile Ser Ile Gly Gln Glu Leu Leu Lys Lys Pro Val Ala Arg
39 355 360 365
40
41 Val Asn Ile Asp Thr Gly Leu Tyr Glu Ser Cys Asp Gly Glu Gly Thr
42 370 375 380
43
44 Asn Ala Gln Ser Leu Ala Asp Phe Ala Lys Gln Leu Ser Asp Glu Arg
45 385 390 395 400
46
47 Lys Leu Arg Lys Ser Asn Leu Asn Ser Asn
48 405 410
49
50 <210> 292
51 <211> 410
52 <212> PRT
53 <213> Zea mays
54
55 <400> 292
56
57 Met Gly Ser Ile Gly Arg Gly Thr Ala Asn Cys Ala Thr Val Pro Gln

1	1	5	10	15
2				
3	Pro	Pro	Pro	Ser
4			Thr	Gly
5			Lys	Leu
6			Ile	Thr
7			Ile	Leu
8			Ser	Ile
9			Asp	Gly
10				
11				
12	Gly	Gly	Ile	Arg
13			Gly	Leu
14			Ile	Pro
15			Ala	Thr
16			Ile	Ile
17			Ala	Tyr
18			Leu	Glu
19				
20				
21				
22				
23				
24				
25				
26				
27				
28				
29				
30				
31				
32				
33				
34				
35				
36				
37				
38				
39				
40				
41				
42				
43				
44				
45				
46				
47				
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				

1 305 310 315 320
2
3 Ala Leu His Cys Glu Lys Lys Tyr Leu Arg Ile Gln Asp Asp Thr Leu
4 325 330 335
5
6 Thr Gly Asn Ala Ser Ser Val Asp Ile Ala Thr Lys Glu Asn Met Glu
7 340 345 350
8
9 Ser Leu Ile Ser Ile Gly Gln Glu Leu Leu Asn Lys Pro Val Ala Arg
10 355 360 365
11
12 Val Asn Ile Asp Thr Gly Leu Tyr Glu Ser Cys Glu Gly Glu Gly Thr
13 370 375 380
14
15 Asn Ala Gln Ser Leu Ala Asp Phe Ala Lys Gln Leu Ser Asp Glu Arg
16 385 390 395 400
17
18 Lys Leu Arg Lys Ser Asn Leu Asn Ser Asn
19 405 410
20
21 <210> 293
22 <211> 337
23 <212> PRT
24 <213> Zea mays
25
26 <400> 293
27
28 Met Gly Ser Ile Gly Arg Gly Thr Ala Asn Cys Ala Thr Val Pro Gln
29 1 5 10 15
30
31 Pro Pro Pro Ser Thr Gly Lys Leu Ile Thr Ile Leu Ser Ile Asp Gly
32 20 25 30
33
34 Gly Gly Ile Arg Gly Leu Ile Pro Ala Thr Ile Ile Ala Tyr Leu Glu
35 35 40 45
36
37 Ala Lys Leu Gln Glu Leu Asp Gly Pro Asp Ala Arg Ile Ala Asp Tyr
38 50 55 60
39
40 Phe Asp Val Ile Ala Gly Thr Ser Thr Gly Ala Leu Leu Ala Ser Met
41 65 70 75 80
42
43 Leu Ala Ala Pro Asp Glu Asn Asn Arg Pro Leu Phe Ala Ala Lys Asp
44 85 90 95
45
46 Leu Thr Thr Phe Tyr Leu Glu Asn Gly Pro Lys Ile Phe Pro Gln Lys
47 100 105 110
48
49 Lys Ala Gly Leu Leu Thr Pro Leu Arg Asn Leu Leu Gly Leu Val Arg
50 115 120 125
51
52 Gly Pro Lys Tyr Asp Gly Val Phe Leu His Asp Lys Ile Lys Ser Leu
53 130 135 140
54
55 Thr His Asp Val Arg Val Ala Asp Thr Val Thr Asn Val Ile Val Pro
56 145 150 155 160
57

1 Ala Phe Asp Val Lys Tyr Leu Gln Pro Ile Ile Phe Ser Thr Tyr Glu
2 165 170 175
3
4 Ala Lys Thr Asp Ala Leu Lys Asn Ala His Leu Ser Asp Ile Cys Ile
5 180 185 190
6
7 Ser Thr Ser Ala Ala Pro Thr Tyr Phe Pro Ala His Phe Phe Lys Thr
8 195 200 205
9
10 Glu Ala Thr Asp Gly Arg Pro Pro Arg Glu Tyr His Leu Val Asp Gly
11 210 215 220
12
13 Gly Val Ala Ala Asn Asn Pro Thr Met Val Ala Met Ser Met Leu Thr
14 225 230 235 240
15
16 Lys Glu Val His Arg Arg Asn Pro Asn Phe Asn Ala Gly Ser Pro Thr
17 245 250 255
18
19 Glu Tyr Thr Asn Tyr Leu Ile Ile Ser Val Gly Thr Gly Ser Ala Lys
20 260 265 270
21
22 Gln Ala Glu Lys Tyr Thr Ala Glu Gln Cys Ala Lys Trp Gly Leu Ile
23 275 280 285
24
25 Gln Trp Leu Tyr Asn Gly Gly Phe Thr Pro Ile Ile Asp Ile Phe Ser
26 290 295 300
27
28 His Ala Ser Ser Asp Met Val Asp Ile His Ala Ser Ile Leu Phe Gln
29 305 310 315 320
30
31 Ala Leu His Cys Glu Lys Lys Tyr Leu Arg Ile Gln Leu Tyr Tyr Ala
32 325 330 335
33
34 Gly
35
36
37
38
39